



Bookmarks

- ▶ [Unit 0: Overview](#)
- ▶ [Entrance Survey](#)
- ▶ [Unit 1: Probability models and axioms](#)
- ▼ [Unit 2: Conditioning and independence](#)

[Unit overview](#)

[Lec. 2: Conditioning and Bayes' rule](#)

[Exercises 2 due Feb 2, 2017 20:59 ART](#)

[Lec. 3: Independence](#)

[Exercises 3 due Feb 2, 2017 20:59 ART](#)

[Solved problems](#)[Problem Set 2](#)

[Problem Set 2 due Feb 2, 2017 20:59 ART](#)

- ▶ [Unit 3: Counting](#)

Unit 2: Conditioning and independence > Lec. 3: Independence > Exercise: Independence of event complements

Exercise: Independence of event complements

[Bookmark this page](#)

Exercise: Independence of event complements

1/1 point (graded)

Suppose that A and B are independent events. Are A^c and B^c independent?

Yes, they are independent ▼

✓ Answer: Yes, they are independent

Answer:

We saw in the previous segment that for any 2 generic events E_1 and E_2 , independence of E_1 and E_2 implies independence of E_1 and E_2^c . In the case of this particular problem, we can apply this result with $E_1 = A$ and $E_2 = B$ to conclude that since A and B are assumed to be independent, then A and B^c are also independent.

Independence is symmetric, so A and B^c being independent is the same as B^c and A being independent. If we now reuse the generic result with $E_1 = B^c$ and $E_2 = A$, we can conclude that B^c and A^c are also independent, which by symmetry is the same as A^c and B^c being independent.

To summarize:

A and B independent $\Rightarrow A$ and B^c independent $\Rightarrow B^c$ and A independent $\Rightarrow B^c$ and A^c independent $\Rightarrow A^c$ and B^c independent

Submit

You have used 1 of 1 attempt

✓ Correct (1/1 point)

© All Rights Reserved



© 2012-2017 edX Inc. All rights reserved except where noted. EdX, Open edX and the edX and Open EdX logos are registered trademarks or trademarks of edX Inc.

POWERED BY
OPENedX

