



Bookmarks

- ▶ [Unit 0: Overview](#)
- ▶ [Entrance Survey](#)
- ▶ [Unit 1: Probability models and axioms](#)
- ▼ [Unit 2: Conditioning and independence](#)

[Unit overview](#)[Lec. 2: Conditioning and Bayes' rule](#)
[Exercises 2 due Feb 2, 2017](#)
 20:59 ART
[Lec. 3: Independence](#)
[Exercises 3 due Feb 2, 2017](#)
 20:59 ART
[Solved problems](#)[Problem Set 2](#)
[Problem Set 2 due Feb 2, 2017](#)
 20:59 ART

- ▶ [Unit 3: Counting](#)

Unit 2: Conditioning and independence > Problem Set 2 > Problem 2 Vertical: A reliability problem


Problem 2 Vertical: A reliability problem

Bookmark this page

Problem 2: A reliability problem

4.0/4.0 points (graded)

Consider the communication network shown in the figure below and suppose that each link can fail with probability p . Assume that failures of different links are independent.



1. Assume that $p = 1/3$. Find the probability that there exists a path from **A** to **B** along which no link has failed. (Give a numerical answer.)

0.4609


✓ Answer: 0.46091

2. Given that exactly one link in the network has failed, find the probability that there exists a path from **A** to **B** along which no link has failed. (Give a numerical answer.)

0.8

✓ Answer: 0.8

Answer:



Let E be the event that there exists an operational path from **A** to **B**. Note that the probability that the Upper path from **A** to **C** is operational is $(1 - p)^2$. So the probability that the Upper path fails is $1 - (1 - p)^2$. Similarly, the Lower path fails with probability $1 - (1 - p)^2$ as well.

1. We can break the problem down into two parts. In order for there to be an operational path from **A** to **B**, there must be an operational path from **A** to **C**, and Link 5 must be operational.

$$\begin{aligned}\mathbf{P}(E) &= \mathbf{P}(\text{there exists a path from } A \text{ to } C \text{ and Link 5 is operational}) \\ &= (1 - \mathbf{P}(\text{Upper path fails and Lower path fails})) \cdot \mathbf{P}(\text{Link 5 is operational}) \\ &= \left\{1 - \left[1 - (1 - p)^2\right]^2\right\} \cdot (1 - p).\end{aligned}$$

When $p = 1/3$, this gives us $\mathbf{P}(E) \approx 0.46091$.

2. Since all links are equally likely to fail and since exactly one link has failed, each link has the same probability $1/5$ of being the one that failed. There will be no path from **A** to **B** only in the case where the link that failed is Link 5, which happens with probability $1/5$. Therefore, the desired probability is $1 - 1/5 = 4/5$.

You have used 1 of 2 attempts

DISCUSSION

Click "Show Discussion" below to see discussions on this problem.

Discussion

Topic: Unit 2/Problem Set 2 / A reliability problem

© All Rights Reserved



© 2012-2017 edX Inc. All rights reserved except where noted. EdX, Open edX and the edX and Open EdX logos are registered trademarks or trademarks of edX Inc.

POWERED BY
OPENedX

