

Quiz 2

[Help Center](#)

The **due date** for this quiz is **Sun 20 Sep 2015 8:30 PM ART**.

☐ In accordance with the Coursera Honor Code, I (Diego Dell'Era) certify that the answers here are my own work.

Question 1

Consider the following data with x as the predictor and y as the outcome.

```
x <- c(0.61, 0.93, 0.83, 0.35, 0.54, 0.16, 0.91, 0.62, 0.62)
y <- c(0.67, 0.84, 0.6, 0.18, 0.85, 0.47, 1.1, 0.65, 0.36)
```

Give a P-value for the two sided hypothesis test of whether β_1 from a linear regression model is 0 or not.

- ☐ 2.325
- ☐ 0.05296
- ☐ 0.391
- ☐ 0.025

Question 2

Consider the previous problem, give the estimate of the residual standard deviation.

- ☐ 0.223
- ☐ 0.4358
- ☐ 0.05296
- ☐ 0.3552

Question 3

In the `mtcars` data set, fit a linear regression model of weight (predictor) on mpg (outcome). Get a 95% confidence interval for the expected mpg at the average weight. What is the lower endpoint?

- ☐ 21.190
- ☐ 18.991
- ☐ -4.00
- ☐ -6.486

Question 4

Refer to the previous question. Read the help file for `mtcars`. What is the weight coefficient interpreted as?

- ☐ The estimated expected change in mpg per 1,000 lb increase in weight.
- ☐ The estimated 1,000 lb change in weight per 1 mpg increase.
- ☐ The estimated expected change in mpg per 1 lb increase in weight.
- ☐ It can't be interpreted without further information

Question 5

Consider again the `mtcars` data set and a linear regression model with mpg as predicted by weight (1,000 lbs). A new car is coming weighing 3000 pounds. Construct a 95% prediction interval for its mpg. What is the upper endpoint?

- ☐ 27.57
- ☐ -5.77
- ☐ 21.25
- ☐ 14.93

Question 6

Consider again the `mtcars` data set and a linear regression model with mpg as predicted by weight (in 1,000 lbs). A “short” ton is defined as 2,000 lbs. Construct a 95% confidence interval for the expected change in mpg per 1 short ton increase in weight. Give the lower endpoint.

- ☐ -12.973
- ☐ 4.2026
- ☐ -6.486
- ☐ -9.000

Question 7

If my X from a linear regression is measured in centimeters and I convert it to meters what would happen to the slope coefficient?

- ☐ It would get multiplied by 10
- ☐ It would get multiplied by 100.
- ☐ It would get divided by 100
- ☐ It would get divided by 10

Question 8

I have an outcome, Y , and a predictor, X and fit a linear regression model with

$Y = \beta_0 + \beta_1 X + \epsilon$ to obtain $\hat{\beta}_0$ and $\hat{\beta}_1$. What would be the consequence to the subsequent slope and intercept if I were to refit the model with a new regressor, $X + c$ for some constant, c ?

- ☐ The new slope would be $c\hat{\beta}_1$
- ☐ The new slope would be $\hat{\beta}_1 + c$

- ☐ The new intercept would be $\hat{\beta}_0 + c\hat{\beta}_1$
- ☐ The new intercept would be $\hat{\beta}_0 - c\hat{\beta}_1$

Question 9

Refer back to the mtcars data set with mpg as an outcome and weight (wt) as the predictor. About what is the ratio of the the sum of the squared errors, $\sum_{i=1}^n (Y_i - \hat{Y}_i)^2$ when comparing a model with just an intercept (denominator) to the model with the intercept and slope (numerator)?

- ☐ 0.25
- ☐ 0.75
- ☐ 0.50
- ☐ 4.00

Question 10

Do the residuals always have to sum to 0 in linear regression?

- ☐ If an intercept is included, then they will sum to 0.
- ☐ The residuals never sum to zero.
- ☐ If an intercept is included, the residuals most likely won't sum to zero.
- ☐ The residuals must always sum to zero.

☐ In accordance with the Coursera Honor Code, I (Diego Dell'Era) certify that the answers here are my own work.

Submit Answers

Save Answers

You cannot submit your work until you agree to the Honor Code. Thanks!

