The manuscript "Floquet Topological Frequency-Converting Amplifier" proposes a Floquet-modulated single-mode bosonic system in which both the mode frequency and the decay rate are periodically driven. In Floquet-Sambe space, this leads to an effective non-Hermitian synthetic lattice, which the authors argue supports topological frequency conversion and directional amplification. The manuscript looks solid, and the proposed implementation in superconducting circuits is potentially interesting. However, in its present form I find that the main physical message of the manuscript is not sufficiently clear.

My main concern is that the paper reads as somewhat incremental with respect to the existing literature on non-Hermitian topological amplification, Floquet synthetic dimensions, and singular-value topology of input-output response matrices. I do not doubt that the calculations may be correct, but the manuscript does not yet make sufficiently transparent what the central conceptual advance is. Is the main result the minimality of the setup, namely that a single driven oscillator can realize this physics? Is the novelty the use of simultaneous frequency and dissipation modulation? Or is there a genuinely new topological amplification mechanism? At present this point is somewhat hidden behind the formal development, and the authors should state it much more explicitly.

I would be open to recommending publication after revision, provided the authors address the previous and following points.

1. The authors should explain more clearly why the periodic modulations generate nearest-neighbor couplings in the synthetic lattice (page 3: *"This structure arises ... leading to nearest- neighbor couplings in the synthetic lattice."*)

2. The piecewise definition of the input-noise term around Eq. (11) should be clarified. The manuscript states that for $\bar{\omega} > \Omega/2$ the last term must be replaced by a different expression.

3. What do the authors mean with "zero–singular-value eigenstates"?

4. Since the manuscript emphasizes non-Hermitian topology, the authors should clarify the role, if any, of the non-Hermitian skin effect. Does the proposed synthetic-frequency lattice exhibit skin-mode accumulation, and can this be observed in an experimentally accessible quantity?

In summary, I find the work potentially publishable, but the manuscript should be revised to make the physical mechanism, novelty, and experimentally observable signatures much clearer. At present the main point is somewhat obscured by the formal calculations, and the authors should make a stronger case that the result is more than an incremental reformulation of known Floquet and non-Hermitian topological-amplifier physics.