


Figure 1. Five factor EPOCH model with 20 indicator items. See Table 1 for estimated confirmatory factor loadings, covariances, and model fit by sample.

This document is copyrighted by the American Psychological Association or one of its allied publishers.
This article is intended solely for the personal use of the individual user and is not to be disseminated broadly.

nia; Santa Ana, California; and Jefferson Parish (near New Orleans), Louisiana. All participants had been arrested for a relatively minor offense, such as theft or simple assault. Participants ranged in age from 13 to 17 ($M = 15.30$, $SD = 1.27$). As part of a larger longitudinal study, participants completed a series of measures at baseline and 6 months later. The 25-item version of the EPOCH measure was included at both occasions; of the 654 adolescents who completed the measure at baseline, 494 also completed the measure at the 6-month follow up. Other measures included IQ, exposure to violence, self-esteem, motivation to succeed, future orientation, psychosocial maturity, general anxiety disorder, major depressive disorder, aggression, school performance, teacher bonding, school misconduct, arrests, social consequences for using drugs or alcohol, and health care visits.

Sample 6: U.S. strengths-based study. Sample 7 consisted of 70 students (32 males, 38 females) from an urban middle school in Philadelphia, PA. Participants ranged in age from 12 to 14 ($M = 13.07$, $SD = 0.35$), were in Grade 8, and predominantly from low SES backgrounds. Students came from three classes (remedial, average, and advanced levels) in an urban charter school; two classes were randomly assigned to a 5-day strengths-based program, and one class completed the measures but did not receive the training. Measures were completed at baseline, at the end of the 5-day program, and 3 months later. At each of the three assessments, participants completed the 25-item version of the EPOCH measure. Age, gender, and treatment condition were also available.

Sample 7: U.S. inner city youth. Sample 7 was specifically targeted to test the structure and cross-time stability of the final 20-item EPOCH measure. The sample consisted of 291 inner-city adolescents (72 males, 80 females, 139 unknown) from a middle school and a high school in west Philadelphia, an economically disadvantaged area of the city. Participants ranged in age from 12 to 17 ($M = 14.48$, $SD = 0.97$), were in Grades 7 through 10, and predominantly from low SES backgrounds. Participants indicated their age, gender, and completed the 20 EPOCH items. Two to three weeks later, 164 youth completed the 20-item measure again. Of these, 82 individuals were successfully matched to their Time 1 score.

Sample 8: Australian girls school. Sample 8 consisted of 327 adolescent girls from a Catholic all-girls school in Victoria, Australia. Participants ranged in age from 13 to 17 ($M = 14.71$, $SD = 1.18$), were in Grades 8 through 11, and predominantly from middle to upper-middle SES backgrounds. The participants completed an online questionnaire, which included the 20-item EPOCH measure. Other measures included life satisfaction, psychological well-being (self-acceptance, purpose in life, environmental mastery, autonomy, and positive relationships), social well-being, hope, depression, anxiety, alcohol and illicit drug use, perceived stress, sleep habits, and exercise.

Sample 9: U.S. hospitalized youth. Sample 9 consisted of 146 adolescents (63 males, 79 females, 4 unknown) from several U.S. hospitals who were inpatients receiving treatment for various medical conditions. Participants ranged in age from 10 to 18 ($M = 13.73$, $SD = 2.52$). No SES information was available. Participants completed a baseline questionnaire, were randomly assigned to a positive intervention (receiving a drawing of a superhero and then drawing a superhero for another child) or control condition (receiving a drawing only), and then reported changes in emotion. The current analysis used the initial assessment from the full sample, regardless of treatment group, which included the 20-item EPOCH measure. Other measures included positive and negative emotion and meaning in life.

Sample 10: U.S. after school program. Sample 10 consisted of 89 students from a public school in Atlanta, New Jersey. Participants ranged in age from 13 to 15 ($M = 13.48$, $SD = 0.55$), were in 8th grade, and from low SES backgrounds. At the beginning of the term, as part of an in-class exercise, students completed the 20-item EPOCH measure and answered questions about perceived academic performance and the extent to which basic needs are met. The survey was repeated at the end of the term, with 62 participants successfully matched across the time points. In addition, objective language arts and math test scores were matched to