

SSIE 660: Applied Stochastic Processes

Homework assignment 9 – Key

Nov. 30, 2016

Due: Dec 7, 2016 Before class starts

1. Solve Chapter 6. Problem 4.

Let $N(t)$ denote the number of customers in the station at time t . Then $\{N(t)\}$ is a birth and death process. The entering rate is the arrival rate multiplied by the probability of entering the system with $\lambda_n = \lambda \alpha_n, \mu_n = \mu$.

2. Solve Chapter 6. Problem 8.

The number of failed machines is a birth and death process with

$$\lambda_0 = 2\lambda, \lambda_1 = \lambda,$$

$$\lambda_n = 0, n > 1$$

$$\mu_1 = \mu_2 = \mu,$$

$$\mu_n = 0, n \neq 1, 2$$

Find more parameters such as v_0, q_{01} , etc. Then write the Kolmogorov backward equations using such parameters. That is,

$$v_0 = q_{01} = 2\lambda$$

$$v_1 = (\lambda + \mu)$$

$$v_2 = q_{21} = \mu$$

$$q_{01} = 2\lambda$$

$$q_{10} = \mu$$

$$q_{12} = \lambda$$

$$q_{21} = \mu$$

$$\text{All other } v_j, q_{ij} = 0$$

The Kolmogorov backward equations are then:

$$\begin{aligned}
i = 0, j = 0; P'_{00}(t) &= 2\lambda[P_{10}(t) - P_{00}(t)] \\
i = 0, j = 1; P'_{01}(t) &= 2\lambda[P_{11}(t) - P_{01}(t)] \\
i = 0, j = 2; P'_{02}(t) &= 2\lambda[P_{12}(t) - P_{02}(t)] \\
i = 1, j = 0; P'_{10}(t) &= \mu P_{00}(t) + \lambda P_{20}(t) - (\lambda + \mu)P_{10}(t) \\
i = 1, j = 1; P'_{11}(t) &= \mu P_{01}(t) + \lambda P_{21}(t) - (\lambda + \mu)P_{11}(t) \\
i = 1, j = 2; P'_{12}(t) &= \mu P_{02}(t) + \lambda P_{22}(t) - (\lambda + \mu)P_{12}(t) \\
i = 1, j = 2; P'_{12}(t) &= \mu P_{02}(t) + \lambda P_{22}(t) - (\lambda + \mu)P_{12}(t) \\
i = 2, j = 0; P'_{20}(t) &= \mu[P_{10}(t) - P_{20}(t)] \\
i = 2, j = 1; P'_{21}(t) &= \mu[P_{11}(t) - P_{21}(t)] \\
i = 2, j = 2; P'_{22}(t) &= \mu[P_{12}(t) - P_{22}(t)]
\end{aligned}$$

3. Solve Chapter 6. Problem 13.

With the number of customers in the shop as the state, we get a birth and death process with

$$\lambda_0 = \lambda_1 = 3, \mu_1 = \mu_2 = 4$$

Therefore

$$P_1 = \frac{3}{4}P_0, P_2 = \frac{3}{4}P_1 = \left(\frac{3}{4}\right)^2 P_0$$

And since $\sum_0^2 P_i = 1$, we get

$$P_0 = \left(1 + \frac{3}{4} + \left(\frac{3}{4}\right)^2\right)^{-1} = \frac{16}{37}$$

(a) The average number of customers in the shop is

$$\begin{aligned}
P_1 + 2P_2 &= \left(\frac{3}{4} + 2\left(\frac{3}{4}\right)^2\right) P_0 \\
&= \frac{30}{16} \left(1 + \frac{3}{4} + 2\left(\frac{3}{4}\right)^2\right) P_0 = \frac{30}{37}
\end{aligned}$$

(b) The proportion of customers that enter the shop is

$$\frac{\lambda(1 - P_2)}{\lambda} = 1 - P_2 = 1 - \frac{9}{16} \cdot \frac{16}{37} = \frac{28}{37}$$

(c) Now $\mu = 8$, and so

$$P_0 = \left(1 + \frac{3}{8} + \left(\frac{3}{8}\right)^2\right)^{-1} = \frac{64}{97}$$

So the proportion of customers who now enter the shop is

$$1 - P_2 = 1 - \left(\frac{3}{8}\right)^2 \frac{264}{97} = 1 - \frac{9}{97} = \frac{88}{97}$$

The rate of added customers is therefore

$$\lambda \left(\frac{88}{97}\right) - \lambda \left(\frac{28}{37}\right) = 3 \left(\frac{88}{97} - \frac{28}{37}\right) = 0.45$$

The business he does would improve by .45 customers per hour.

4. Solve Chapter 6. Problem 14.

Letting the number of cars in the station be the states, we have a birth and death process with

$$\lambda_0 = \lambda_1 = \lambda_2 = 20$$

$$\lambda_i = 0, i > 2$$

$$\mu_1 = \mu_2 = 12 \text{ (12 cars per hour)}$$

Hence,

$$P_1 = \frac{5}{3} P_0$$

$$P_2 = \frac{5}{3} P_1 = \left(\frac{5}{3}\right)^2 P_0$$

$$P_3 = \frac{5}{3} P_2 = \left(\frac{5}{3}\right)^3 P_0$$

In addition,

$$\sum_{i=0}^3 P_i = 1$$

Therefore,

$$P_0 = \left(1 + \frac{5}{3} + \left(\frac{5}{3}\right)^2 + \left(\frac{5}{3}\right)^3\right)^{-1} = \frac{27}{272}$$

- (a) The fraction of the attendant's time spent servicing cars is equal to the fraction of time there are cars in the system and is therefore $1 - P_0 = 245/272$.
- (b) The fraction of potential customers that are lost is equal to the fraction of customers that arrive when there are three cars in the station and is therefore

$$P_3 = \left(\frac{5}{3}\right)^3 P_0 = \frac{125}{272}$$

5. Solve Chapter 6. Problem 15.

With the number of customers in the system as the state, we get a birth and death process with

$$\lambda_0 = \lambda_1 = \lambda_2 = 3$$

$$\lambda_i = 0, i > 2$$

$$\mu_1 = 2, \mu_2 = \mu_3 = 4$$

Therefore, the balance equations reduce to

$$P_1 = \frac{3}{2}P_0, P_2 = \frac{3}{4}P_1 = \frac{9}{8}P_0, P_3 = \frac{3}{4}P_2 = \frac{27}{32}P_0$$

And therefore,

$$P_0 = \left(1 + \frac{3}{2} + \frac{9}{8} + \frac{27}{32}\right)^{-1} = \frac{32}{143}$$

(a) The fraction of potential customers that enter the system is

$$1 - P_3 = 1 - \frac{27}{32} * \frac{32}{143} = \frac{116}{143}$$

(b) With a server working twice as fast we would get

$$P_1 = \frac{3}{4}P_0, P_2 = \frac{3}{4}P_1 = \left(\frac{3}{4}\right)^2 P_0, P_3 = \left(\frac{3}{4}\right)^3 P_0$$

And therefore,

$$P_0 = \left(1 + \frac{3}{4} + \left(\frac{3}{4}\right)^2 + \left(\frac{3}{4}\right)^3\right)^{-1} = \frac{64}{175}$$

so that now

$$1 - P_3 = 1 - \frac{27}{64} * \frac{64}{175} = \frac{148}{175}$$

6. Solve Chapter 6. Problem 19.

There are four states. Let state 0 mean that no machines are down, state 1 that machine 1 is down and 2 is up, state 2 that machine 1 is up and 2 is down, and 3 that both machines are down (machine 1 is in service).

The balance equations are as follows:

$$(\lambda_1 + \lambda_2)P_0 = \mu_1P_1 + \mu_2P_2$$

$$(\mu_1 + \lambda_2)P_1 = \lambda_1P_0$$

$$(\lambda_1 + \mu_2)P_2 = \lambda_2P_0 + \mu_1P_3$$

$$\mu_1P_3 = \lambda_2P_1 + \lambda_1P_2$$

$$\sum_{i=0}^3 P_i = 1$$

These equations are easily solved and the proportion of time machine 2 is down is $P_2 + P_3$.