

# CSCI 4830 / 5722

# Computer Vision



University of Colorado **Boulder**

# CSCI 4830 / 5722

# Computer Vision



Dr. Ioana Fleming  
Spring 2019  
Lecture 2



University of Colorado **Boulder**

# Reminders

## Submissions:

- Homework 1: Sat 1/26 at 6 pm

Moodle, Matlab, Piazza

Disabilities forms

## Readings:

- Szeliski 2.1 before Friday



University of Colorado **Boulder**

# What is Computer Vision?

Extracting information from images

Methods for: acquiring, processing, analyzing and interpreting images.

Digital images

- from a camera (usually)
- divided into smaller elements (pixels)
- stored as a matrix of numbers, usually positive (0-255)
- 8-bit, 12-bit, 16-bit, or float
- color image has 3 channels: R, G, B



University of Colorado  
Boulder

# What is Computer Vision?

Extracting information from images

Besides pixel information, what else can we use?

- information about the camera parameters
  - if known
  - if not, derive them from information from multiple images
  - use differences between images




University of Colorado  
Boulder

# Image Transformations: 2D

Motivation:

- mosaics



University of Colorado  
Boulder

# Image Transformations: 2D


Motivation:

- medical imaging: multi-modality registration /fusion

**Image registration:**


“the determination of a one-to-one mapping between the coordinates in one space and those in another, such that points in the two spaces that correspond to the same anatomical point are mapped to each other.”

Calvin Maurer '93



University of Colorado  
Boulder


# Image Transformations: 2D



University of Colorado **Boulder**

# Image Transformations: 2D

Solution: fit the parameters of some transformation according to a set of corresponding/matching “feature” pairs




University of Colorado  
Boulder

# Image Transformations: 2D

Transformations: use “location” of each corresponding feature pair.

Location: image coordinates vs world coordinates



University of Colorado  
Boulder

# Image coordinates, Points, Vectors

$$\mathbf{x} = \begin{pmatrix} x_1 \\ x_2 \\ \vdots \\ x_n \end{pmatrix}$$

e.g.


$$\mathbf{x} = \begin{pmatrix} 3 \\ 2 \\ 5 \end{pmatrix}$$

The **length** of  $x$ , a.k.a. the **norm** or **2-norm** of  $x$ , is

e.g.,

$$\|\mathbf{x}\| = \sqrt{x_1^2 + x_2^2 + \cdots + x_n^2}$$

$$\|\mathbf{x}\| = \sqrt{3^2 + 2^2 + 5^2} = \sqrt{38}$$



University of Colorado  
Boulder

## Vector Arithmetic

**Addition** To add two vectors together, add them componentwise

```
>> v1 = [3 2 5]'
```

```
v1 =
```

```
3  
2  
5
```

```
>> v2 = [1 3 2]'
```

```
v2 =
```

```
1  
3  
2
```

```
>> v1+v2
```

```
ans =
```


```
4  
5  
7
```



University of Colorado  
Boulder

# Vector Addition


$$u + v = \begin{bmatrix} u_1 \\ u_2 \end{bmatrix} + \begin{bmatrix} v_1 \\ v_2 \end{bmatrix} = \begin{bmatrix} u_1 + v_1 \\ u_2 + v_2 \end{bmatrix}$$



University of Colorado  
Boulder

# Vector Subtraction

$$u - v = \begin{bmatrix} u_1 \\ u_2 \end{bmatrix} - \begin{bmatrix} v_1 \\ v_2 \end{bmatrix} = \begin{bmatrix} u_1 - v_1 \\ u_2 - v_2 \end{bmatrix}$$



## Scaling a vector

$$\mathbf{z} = \alpha \mathbf{x}$$

for a scalar  $\alpha$  then

$$\mathbf{z} = \alpha \begin{pmatrix} 3 \\ 2 \\ 5 \end{pmatrix} = \begin{pmatrix} 3\alpha \\ 5\alpha \\ 2\alpha \end{pmatrix}$$

(This is just like stretching/shrinking the vector by a factor  $\alpha$ )




University of Colorado  
Boulder

# Scaling an object


**Scaling** a coordinate means multiplying each of its components by a scalar.

**Uniform scaling** means this scalar is the same for all components:



# Scaling an object

**Non-uniform scaling** means different scalar per component



University of Colorado  
Boulder

#

# Inner product of two vectors

$$a = \begin{bmatrix} 6 \\ 2 \\ -3 \end{bmatrix} \quad b = \begin{bmatrix} 4 \\ 1 \\ 5 \end{bmatrix}$$

$$a \cdot b = a^T b = [6 \quad 2 \quad -3] \begin{bmatrix} 4 \\ 1 \\ 5 \end{bmatrix} = 6 \cdot 4 + 2 \cdot 1 + (-3) \cdot 5 = 11$$




University of Colorado  
Boulder

# Inner (dot) Product

$$u^T v = \begin{bmatrix} u_1 & u_2 \end{bmatrix} \begin{bmatrix} v_1 \\ v_2 \end{bmatrix} = u_1 \cdot v_1 + u_2 \cdot v_2$$

The inner product is a **SCALAR**.



$$u^T v = \begin{bmatrix} u_1 & u_2 \end{bmatrix} \begin{bmatrix} v_1 \\ v_2 \end{bmatrix} = \|u\| \|v\| \cos \alpha$$

$$u^T v = 0 \Leftrightarrow u \perp v$$

---

In matlab use `sum(a.*b)`

`>> a = [1 2 3]`

`a =`

1            2            3

`>> b = [2 3 4]`

`b =`

2            3            4

`>> sum(a.*b)`

`ans =`

20

---



Univ  
Bou

## Matrices

A matrix is an  $n \times M$  element array

$$A = \begin{bmatrix} a_{11} & a_{12} & \dots & a_{1m} \\ a_{21} \\ \vdots \\ \vdots \\ a_{N1} & \dots & \dots & a_{NM} \end{bmatrix}$$

**transpose** of matrix A (written  $A^T$ ) is  $a_{ji}$  (A with rows and columns flipped)



University of Colorado  
Boulder

# Transpose of a Matrix

Transpose:

$$C_{m \times n} = A^T \quad n \times m \qquad (A + B)^T = A^T + B^T$$

$$c_{ij} = a_{ji} \qquad (AB)^T = B^T A^T$$

Examples:

$$\begin{bmatrix} 6 & 2 \\ 1 & 5 \end{bmatrix}^T = \begin{bmatrix} 6 & 1 \\ 2 & 5 \end{bmatrix}$$

$$\begin{bmatrix} 6 & 2 \\ 1 & 5 \\ 3 & 8 \end{bmatrix}^T = \begin{bmatrix} 6 & 1 & 3 \\ 2 & 5 & 8 \end{bmatrix}$$

If  $A^T = A$ , we say  $A$  is symmetric.



University of Colorado  
Boulder

# Transpose of a Matrix: properties

$$(A^T)^T = A$$

$$(A + B)^T = A^T + B^T$$

$$(AB)^T = B^T A^T$$

$$(rA^T) = rA^T$$



University of Colorado  
Boulder

## Transpose in Matlab

In Matlab use A'

```
>> A = [1 2 ; 3 4]
```

```
A =
```

```
1 2  
3 4
```

```
>> A'
```

```
ans =
```

```
1 3  
2 4
```



University of Colorado  
Boulder

## Matrix Multiplication Example

$$\begin{bmatrix} 1 & 2 & 3 \\ 3 & 2 & 1 \end{bmatrix} \begin{bmatrix} 4 & 5 \\ 10 & 2 \\ 2 & 10 \end{bmatrix} = \begin{bmatrix} a_{11} & a_{12} \\ a_{21} & a_{22} \end{bmatrix}$$



University of Colorado  
Boulder

## Matrix Multiplication Example

$$\begin{bmatrix} 1 & 2 & 3 \\ 3 & 2 & 1 \end{bmatrix} \begin{bmatrix} 4 & 5 \\ 10 & 2 \\ 2 & 10 \end{bmatrix} = \begin{bmatrix} a_{11} & a_{12} \\ a_{21} & a_{22} \end{bmatrix}$$

$$a_{11} = 1 * 4 +$$



University of Colorado  
Boulder

## Matrix Multiplication Example

$$\begin{bmatrix} 1 & 2 & 3 \\ 3 & 2 & 1 \end{bmatrix} \begin{bmatrix} 4 & 5 \\ 10 & 2 \\ 2 & 10 \end{bmatrix} = \begin{bmatrix} a_{11} & a_{12} \\ a_{21} & a_{22} \end{bmatrix}$$

$$a_{11} = 1 * 4 + 2 * 10 +$$



University of Colorado  
Boulder

## Matrix Multiplication Example

$$\begin{bmatrix} 1 & 2 & 3 \\ 3 & 2 & 1 \end{bmatrix} \begin{bmatrix} 4 & 5 \\ 10 & 2 \\ 2 & 10 \end{bmatrix} = \begin{bmatrix} 30 & a_{12} \\ a_{21} & a_{22} \end{bmatrix}$$

$$a_{11} = 1 * 4 + 2 * 10 + 3 * 2 = 30$$



University of Colorado  
Boulder

## Matrix Multiplication Example

$$\begin{bmatrix} 1 & 2 & 3 \\ 3 & 2 & 1 \end{bmatrix} \begin{bmatrix} 4 & 5 \\ 10 & 2 \\ 2 & 10 \end{bmatrix} = \begin{bmatrix} 30 & 39 \\ a_{21} & a_{22} \end{bmatrix}$$

$$a_{12} = 1 * 5 + 2 * 2 + 3 * 10 = 39$$



University of Colorado  
Boulder

## Matrix Multiplication Example

$$\begin{bmatrix} 1 & 2 & 3 \\ 3 & 2 & 1 \end{bmatrix} \begin{bmatrix} 4 & 5 \\ 10 & 2 \\ 2 & 10 \end{bmatrix} = \begin{bmatrix} 30 & 39 \\ 34 & a_{22} \end{bmatrix}$$

$$a_{12} = 3 * 4 + 2 * 10 + 1 * 2 = 34$$



University of Colorado  
Boulder

## Matrix Multiplication Example

$$\begin{bmatrix} 1 & 2 & 3 \\ 3 & 2 & 1 \end{bmatrix} \begin{bmatrix} 4 & 5 \\ 10 & 2 \\ 2 & 10 \end{bmatrix} = \begin{bmatrix} 30 & 39 \\ 34 & 29 \end{bmatrix}$$

$$a_{12} = 3 * 5 + 2 * 2 + 1 * 10 = 29$$



University of Colorado  
Boulder

# Matrix Product

Product:

$$C_{n \times p} = A_{n \times m} B_{m \times p}$$

A and B must have compatible dimensions

$$c_{ij} = \sum_{k=1}^m a_{ik} b_{kj}$$

In Matlab: >> A\*B

Example:

$$\begin{bmatrix} 2 & 5 \\ 3 & 1 \end{bmatrix} \cdot \begin{bmatrix} 6 & 2 \\ 1 & 5 \end{bmatrix} = \begin{bmatrix} 17 & 29 \\ 19 & 11 \end{bmatrix}$$

$$\begin{bmatrix} 6 & 2 \\ 1 & 5 \end{bmatrix} \cdot \begin{bmatrix} 2 & 5 \\ 3 & 1 \end{bmatrix} = \begin{bmatrix} \quad & \quad \\ \quad & \quad \end{bmatrix}$$

Matrix Multiplication is not commutative:

$$A_{n \times n} B_{n \times n} \neq B_{n \times n} A_{n \times n}$$



University of Colorado  
Boulder

# Matrix Addition

Sum:

$$A_{n \times m} = \begin{bmatrix} a_{11} & a_{12} & \dots & a_{1m} \\ a_{21} & a_{22} & \dots & a_{2m} \\ \dots & & & \\ a_{n1} & a_{n2} & \dots & a_{nm} \end{bmatrix}$$

$$\begin{aligned} C_{n \times m} &= A_{n \times m} + B_{n \times m} \\ c_{ij} &= a_{ij} + b_{ij} \end{aligned}$$

Example:

A and B must have the same dimensions

$$\begin{bmatrix} 2 & 5 \\ 3 & 1 \end{bmatrix} + \begin{bmatrix} 6 & 2 \\ 1 & 5 \end{bmatrix} = \begin{bmatrix} 8 & 7 \\ 4 & 6 \end{bmatrix}$$



University of Colorado  
Boulder

# Determinant of a Matrix

Determinant:

A must be square

$$\det \begin{bmatrix} a_{11} & a_{12} \\ a_{21} & a_{22} \end{bmatrix} = \begin{vmatrix} a_{11} & a_{12} \\ a_{21} & a_{22} \end{vmatrix} = a_{11}a_{22} - a_{21}a_{12}$$

Example:

$$\det \begin{bmatrix} 2 & 5 \\ 3 & 1 \end{bmatrix} = 2 - 15 = -13$$

$$\det \begin{bmatrix} a_{11} & a_{12} & a_{13} \\ a_{21} & a_{22} & a_{23} \\ a_{31} & a_{32} & a_{33} \end{bmatrix} = a_{11} \begin{vmatrix} a_{22} & a_{23} \\ a_{32} & a_{33} \end{vmatrix} - a_{12} \begin{vmatrix} a_{21} & a_{23} \\ a_{31} & a_{33} \end{vmatrix} + a_{13} \begin{vmatrix} a_{21} & a_{22} \\ a_{31} & a_{32} \end{vmatrix}$$



University of Colorado  
Boulder

# Determinant in Matlab

```
In matlab use det(A)

>> A = [ 1 2 ; 3 4]

A =

    1     2
    3     4

>> det(A)

ans =

    -2
```



# Inverse of a Matrix

- If  $A$  is a square matrix, the **inverse** of  $A$ , called  $A^{-1}$ , satisfies

$$AA^{-1} = I \quad \text{and} \quad A^{-1}A = I,$$

- Where  $I$ , the **identity matrix**, is a diagonal matrix with all 1's on the diagonal.

$$I_2 = \begin{bmatrix} 1 & 0 \\ 0 & 1 \end{bmatrix}$$

$$I_3 = \begin{bmatrix} 1 & 0 & 0 \\ 0 & 1 & 0 \\ 0 & 0 & 1 \end{bmatrix}$$



University of Colorado  
Boulder

# Inverse of a 2D Matrix

For a 2-D matrix

$$A = \begin{bmatrix} a & b \\ c & d \end{bmatrix}$$

$$A^{-1} = \frac{\begin{bmatrix} d & -b \\ -c & a \end{bmatrix}}{|A|}$$

Example:

$$\begin{bmatrix} 6 & 2 \\ 1 & 5 \end{bmatrix}^{-1} = \frac{1}{28} \begin{bmatrix} 5 & -2 \\ -1 & 6 \end{bmatrix}$$

$$\begin{bmatrix} 6 & 2 \\ 1 & 5 \end{bmatrix}^{-1} \cdot \begin{bmatrix} 6 & 2 \\ 1 & 5 \end{bmatrix} = \frac{1}{28} \begin{bmatrix} 5 & -2 \\ -1 & 6 \end{bmatrix} \cdot \begin{bmatrix} 6 & 2 \\ 1 & 5 \end{bmatrix} = \frac{1}{28} \begin{bmatrix} 28 & 0 \\ 0 & 28 \end{bmatrix} = \begin{bmatrix} 1 & 0 \\ 0 & 1 \end{bmatrix}$$



University of Colorado  
Boulder

# Inverses in Matlab

in Matlab use `inv(A)`

```
>> A = [1 2 ; 3 4]
```

```
A =
```

```
1 2  
3 4
```

```
>> inv(A)
```

```
ans =
```

```
-2.0000 1.0000  
1.5000 -0.5000
```



University of Colorado  
Boulder

# Trace of a matrix

**trace** of a Matrix is  $\text{Tr}(A) =$

$$\sum_{i=1}^N a_{ii}$$

(the sum of the diagonal entries)

In matlab use `trace(A)`



University of Colorado  
Boulder

# Next lecture

- Continue image transformations in 2D and Linear Algebra Review
- Readings: Szeliski 2.1



University of Colorado  
Boulder