

# CSCI 4830 / 5722

# Computer Vision



University of Colorado **Boulder**

# CSCI 4830 / 5722

# Computer Vision



Dr. Ioana Fleming  
Spring 2019  
Lecture 4



University of Colorado **Boulder**

# Reminders

## Submissions:

- Homework 1: Sat 1/26 at 6 pm

## Readings:

- Szeliski Ch. 3
- P&F Ch. 1 (or 1&2 in 1st edition)

Office hours calendar - on Moodle

Moodle (password: '*vision*')

---



University of Colorado **Boulder**


# Linear transformations

A linear transformation from a linear vector space  $\mathbf{R}^n$  to  $\mathbf{R}^m$  is defined as a map  $L: \mathbf{R}^n \rightarrow \mathbf{R}^m$  such that

- $L(x + y) = L(x) + L(y)$
- $L(\alpha * x) = \alpha * L(x)$

$L$  can be represented as a matrix  $A$ , such that

$$L(x) = A^*x$$

  
vector

$$\begin{bmatrix} x \\ y \end{bmatrix}$$



# Linear transformations

Examples:

Identity:

- identity operation:

$$x' = x$$

$$y' = y$$

- With a linear transformation matrix:

$$\begin{bmatrix} x' \\ y' \end{bmatrix} = \begin{bmatrix} 1 & 0 \\ 0 & 1 \end{bmatrix} * \begin{bmatrix} x \\ y \end{bmatrix}$$

**identity  
matrix**



University of Colorado **Boulder**

# Linear transformations

Examples:

Inversion:

- inversion operation:

$$x' = -x$$

$$y' = -y$$

- With a linear transformation matrix:

$$\begin{bmatrix} x' \\ y' \end{bmatrix} = \begin{bmatrix} -1 & 0 \\ 0 & -1 \end{bmatrix} * \begin{bmatrix} x \\ y \end{bmatrix}$$

**inversion**

**matrix**



University of Colorado **Boulder**

# Linear transformations

## Scaling:

- Scaling operation:

$$x' = a * x$$

$$y' = b * y$$

- With a linear transformation matrix:

$$\begin{bmatrix} x' \\ y' \end{bmatrix} = \begin{bmatrix} a & 0 \\ 0 & b \end{bmatrix} * \begin{bmatrix} x \\ y \end{bmatrix}$$

**scaling  
matrix**



University of Colorado **Boulder**

# Other Linear transformations matrices

## Mirror / reflection:

- About y axis:

$$\begin{bmatrix} x \\ y \end{bmatrix} = \begin{bmatrix} -1 & 0 \\ 0 & 1 \end{bmatrix} * \begin{bmatrix} x \\ y \end{bmatrix}$$

- Over (0,0):

$$\begin{bmatrix} x \\ y \end{bmatrix} = \begin{bmatrix} -1 & 0 \\ 0 & -1 \end{bmatrix} * \begin{bmatrix} x \\ y \end{bmatrix}$$




University of Colorado **Boulder**

# Other Linear transformations matrices

## Rotation:

- 2D rotation about (0,0):

$$\begin{bmatrix} x' \\ y' \end{bmatrix} = \begin{bmatrix} \cos\theta & \sin\theta \\ -\sin\theta & \cos\theta \end{bmatrix} * \begin{bmatrix} x \\ y \end{bmatrix}$$




University of Colorado **Boulder**

# What about ...?

## Translation:

- 2D translation operations:

$$x' = x + a \quad \leftarrow$$

$$y' = y + b \quad \leftarrow$$

$$\begin{bmatrix} x' \\ y' \end{bmatrix} = \begin{bmatrix} x \\ y \end{bmatrix} + \begin{bmatrix} t_x \\ t_y \end{bmatrix}$$

- Is it possible to have a linear transformation?

$$\begin{bmatrix} x' \\ y' \end{bmatrix} = \begin{bmatrix} a & b \\ c & d \end{bmatrix} * \begin{bmatrix} x \\ y \end{bmatrix}$$

~~$\begin{bmatrix} a & b \\ c & d \end{bmatrix}$~~



University of Colorado **Boulder**

# 2D Linear transformations

- Is it possible to have a 2D linear transformation for translation?

$$\begin{bmatrix} x' \\ y' \end{bmatrix} = \begin{bmatrix} a & b \\ c & d \end{bmatrix} * \begin{bmatrix} x \\ y \end{bmatrix}$$

- Only linear 2D transformations can be represented with a 2x2 transformation matrix.
- Linear transformations are combinations of: scale, rotation, mirror and shear



# Homogeneous coordinates

## Conversions for 2D and 3D

Converting to *homogeneous* coordinates

$$(x, y) \Rightarrow \begin{bmatrix} x \\ y \\ 1 \end{bmatrix}$$

homogeneous image  
coordinates

$$(x, y, z) \Rightarrow \begin{bmatrix} x \\ y \\ z \\ 1 \end{bmatrix}$$

homogeneous scene  
coordinates

Converting *from* homogeneous coordinates

$$\begin{bmatrix} x \\ y \\ w \end{bmatrix} \Rightarrow (x/w, y/w)$$

$$\begin{bmatrix} x \\ y \\ z \\ w \end{bmatrix} \Rightarrow (x/w, y/w, z/w)$$



University of Colorado **Boulder**

# Homogeneous coordinates

$$\begin{bmatrix} 3 \\ 5 \\ 1 \end{bmatrix} \rightarrow \begin{bmatrix} 3 \\ 5 \\ 1 \end{bmatrix} \begin{bmatrix} 6 \\ 10 \\ 2 \end{bmatrix} \begin{bmatrix} 9 \\ 15 \\ 3 \end{bmatrix} \dots$$

Invariant to scaling

$$k \begin{bmatrix} x \\ y \\ w \end{bmatrix} = \begin{bmatrix} kx \\ ky \\ kw \end{bmatrix} \Rightarrow \begin{bmatrix} \frac{kx}{kw} \\ \frac{ky}{kw} \end{bmatrix} = \begin{bmatrix} \frac{x}{w} \\ \frac{y}{w} \end{bmatrix}$$

Homogeneous  
Coordinates

Cartesian  
Coordinates




University of Colorado **Boulder**

# Translation

$$\begin{bmatrix} x' \\ y' \\ 1 \end{bmatrix} = \begin{bmatrix} 1 & 0 & t_x \\ 0 & 1 & t_y \\ 0 & 0 & 1 \end{bmatrix} \begin{bmatrix} x \\ y \\ 1 \end{bmatrix}$$

$3 \times 1$        $3 \times 3$        $3 \times 1$

$$x' = x + t_x$$
$$y' = y + t_y$$



University of Colorado **Boulder**

# Other 2D Transformations

Basic 2D Transformations at 3x3 transformation matrices:  
translation, rotation, scale, shear



$$\begin{bmatrix} x' \\ y' \\ 1 \end{bmatrix} = \begin{bmatrix} \cos\theta & -\sin\theta & 0 \\ \sin\theta & \cos\theta & 0 \\ 0 & 0 & 1 \end{bmatrix} \begin{bmatrix} x \\ y \\ 1 \end{bmatrix} \quad - \text{Rot}$$

$$\begin{bmatrix} \cos\theta & -\sin\theta & tx \\ \sin\theta & \cos\theta & ty \\ 0 & 0 & 1 \end{bmatrix} \begin{bmatrix} x \\ y \\ 1 \end{bmatrix}$$

$$\begin{bmatrix} x' \\ y' \\ 1 \end{bmatrix} = \begin{bmatrix} s_x & 0 & 0 \\ 0 & s_y & 0 \\ 0 & 0 & 1 \end{bmatrix} \begin{bmatrix} x \\ y \\ 1 \end{bmatrix} \quad \text{Scale} \rightarrow$$

$$\begin{aligned} R+T \\ x' &= s_x * x \\ y' &= s_y * y \end{aligned}$$

Shear




University of Colorado **Boulder**

3D Rotation of Points

$$R' = \frac{\begin{bmatrix} \cos \theta & \sin \theta \\ -\sin \theta & \cos \theta \end{bmatrix}}{\det(R)} = \begin{bmatrix} \cos \theta & \sin \theta \\ -\sin \theta & \cos \theta \end{bmatrix}$$

Slide Credit: Saverese

Rotation around the coordinate axes, counter-clockwise:



$$(yz) \quad R_x(\alpha) = \begin{bmatrix} 1 & 0 & 0 \\ 0 & \cos \alpha & -\sin \alpha \\ 0 & \sin \alpha & \cos \alpha \end{bmatrix}$$

$$(xz) \quad R_y(\beta) = \begin{bmatrix} \cos \beta & 0 & \sin \beta \\ 0 & 1 & 0 \\ -\sin \beta & 0 & \cos \beta \end{bmatrix}$$

$$(xy) \quad R_z(\gamma) = \begin{bmatrix} \cos \gamma & -\sin \gamma & 0 \\ \sin \gamma & \cos \gamma & 0 \\ 0 & 0 & 1 \end{bmatrix}$$

Enter angles



University of Colorado **Boulder**

# Rigid transformations

Transformation between two coordinate systems includes a rotation and a translation

Before:

$$\begin{bmatrix} x' \\ y' \end{bmatrix} = R \begin{bmatrix} x \\ y \end{bmatrix} + T$$

2x2      2x1


$$\begin{bmatrix} t_x \\ t_y \end{bmatrix}$$

With homogeneous coordinates:

$$\begin{bmatrix} x \\ y \\ 1 \end{bmatrix} = \begin{bmatrix} R & T \\ 0 & 0 & 1 \end{bmatrix} \begin{bmatrix} x \\ y \\ 1 \end{bmatrix}$$



# A camera creates an image ...



The image intensity  $I(x,y)$  measures how much light is captured at pixel  $(x,y)$ .

- Where does a point  $(X,Y,Z)$  in the world get imaged?
- What is the brightness at the resulting point  $(x,y)$ ?



University of Colorado **Boulder**

# A camera creates an image ...




1. Geometry:
  - Geometry of images captured by cameras
2. Radiometry:
  - What do the intensities in the image tell us about the light in the real world?
3. Simplified models



University of Colorado **Boulder**

# Image formation



Let's design a camera


- Idea 1: put a piece of film in front of an object
- Do we get a reasonable image?



University of Colorado **Boulder**

Slide source: Seitz

# Pinhole camera



Idea 2: add a barrier to block off most of the rays


- This reduces blurring
- The opening known as the **aperture**



University of Colorado **Boulder**


Slide source: Seitz

# Vignetting Effect



University of Colorado **Boulder**

# Pinhole camera



$f$  = focal length

c = center of the camera



University of Colorado **Boulder**  
Figure from Forsyth

# Camera obscura: the pre-camera

- Known during classical period in China (Mozi, China, 470BC to 390BC) and Greece (Aristotle, 384BC to 322BC)




Illustration of Camera Obscura



Freestanding camera obscura at UNC Chapel Hill

Photo by Seth Ilys



University of Colorado **Boulder**

# Camera Obscura




Fig. 434.

Lens Based Camera Obscura, 1568

A 19th-century artist using a camera obscura to outline his subject



University of Colorado **Boulder**

# First Photograph



Joseph Niepce, 1826-27

Oldest surviving photograph  
– Took 8 hours on pewter plate



University of Colorado **Boulder**

# Multiple pinholes



University of Colorado **Boulder**

# Multiple pinholes



University of Colorado **Boulder**

# Multiple pinholes



University of Colorado **Boulder**


# Natural Pinhole Phenomenon



University of Colorado **Boulder**


# Dimensionality Reduction Machine (3D to 2D)

*3D world*



Point of observation

*2D image*



University of Colorado **Boulder**

Figures © Stephen E. Palmer, 2002

# The Pinhole Camera

$(0, i, j, k)$  - camera coord system

image plane -  $\parallel (i, j)$




FIGURE 1.4: The perspective projection equations are derived in this section from the collinearity of the point  $P$ , its image  $p$ , and the pinhole  $O$ .


$d$  - distance between  $\Pi$  and  $O$   
 $c$  - image center

$Oc$  - optical axis  $\perp \Pi$



University of Colorado **Boulder**

# Perspective (pinhole) Projection



$$P = (X, Y, Z)$$

$$\begin{cases} x = \lambda X \\ y = \lambda Y \\ d = \lambda Z \end{cases} \iff \lambda = \frac{x}{X} = \frac{y}{Y} = \frac{d}{Z}$$

$$p = (x, y, z)$$

$\downarrow z = d$

$$\begin{cases} x = d \frac{X}{Z}, \\ y = d \frac{Y}{Z}. \end{cases}$$

$O, P, p$  colinear


$$\overrightarrow{O_p} = \lambda \overrightarrow{OP}$$

$f$



University of Colorado **Boulder**

# The Pinhole Camera




$$x = -\frac{fX}{Z} \quad y = -\frac{fY}{Z}$$

$$\begin{aligned} d < 0 && f = -d \\ f > 0 && \end{aligned}$$



University of Colorado **Boulder**

# The image is inverted



This was pointed out by Kepler in 1604

*d is negative!*

*Image plane at distance = focal length from the pinhole*

But this is no big deal. The brain can interpret it the right way. And for a camera, software can simply flip the image top-down and right-left. After this trick, we get


$$x = \frac{fX}{Z} \quad y = \frac{fY}{Z}$$



University of Colorado  
**Boulder**

From Descartes(1637), La Dioptrique

# Human retina vs camera retina



From Descartes(1637), *La Dioptrique*




FIGURE 1.14: Physical and normalized image coordinate systems.

- Digital images (and retinas) - spatially discrete, not continuous
- Perspective equation – only valid when all distances are measured in the camera’s reference frame, and when image coordinates have their origin at the image center where the axis of symmetry of the camera pierces its retina



University of Colorado **Boulder**

# Camera retina





FIGURE 1.14: Physical and normalized image coordinate systems.

In practice, world and camera coordinates are related by a series of factors:

- focal length
- size of pixels
- position of the image center
- position and orientation of the camera



**Intrinsic Parameters**




**Extrinsic Parameters**



University of Colorado **Boulder**

# Some perspective phenomena...



University of Colorado **Boulder**


# Projective Geometry

What is lost?



University of Colorado **Boulder**

# Length is not preserved



University of Colorado **Boulder**

Figure by David Forsyth

# Projective Geometry

What is lost?

- Length
- Angles



University of Colorado **Boulder**

# Projective Geometry


What is preserved?

- Straight lines are still straight



University of Colorado **Boulder**

# Parallel lines




The projections of two parallel lines lying in some plane  $\Phi$  appear to converge on a horizon line  $h$  formed by the intersection of the image plane  $\Pi$  with the plane parallel to  $\Phi$  and passing through the pinhole. Note that the line  $L$  parallel to  $\Pi$  in  $\Phi$  has no image at all.



# Vanishing points and lines

Parallel lines in the world  
intersect in the image at a  
“vanishing point”



University of Colorado **Boulder**