


Improved Differentially Private Analysis of Variance


Marika Swanberg Ira Globus-Harris Iris Griffith
Anna Ritz Andrew Bray Adam Groce

Reed College

Observed Data, $n = 30$



Observed Data, $n = 30$



Are gene allele and tumor size dependent?

Metric for dependency

$$F\text{-test} = \frac{\text{Variation between groups}}{\text{In-group variation}}$$


Metric for dependency

$$F\text{-test} = \frac{\text{Variation between groups}}{\text{In-group variation}}$$


$$SSA(D) = \sum_{j=1}^k n_j (\bar{y}_j - \bar{y})^2 / (k - 1)$$

$$SSE(D) = \sum_{i=1}^n (y_i - \bar{y}_{c_i})^2 / (n - k)$$

Simulate Random Data



Reference Distribution of F



Now do we think our data supports independence of gene allele and tumor size?

Why is F -test optimal?

Why is F -test optimal?

High probability of indicating dependence when variables are dependent . . .

Why is F -test optimal?

High probability of indicating dependence when variables are dependent . . .
. . . even when dataset is small.

Why is F -test optimal?

High probability of indicating dependence when variables are dependent . . .
. . . even when dataset is small.

Definition (Power)

The **power** of a hypothesis test is the probability it rejects H_0 . It depends on the alternate distribution H_A and n .

Why is F -test optimal?


High probability of indicating dependence when variables are dependent . . .
. . . even when dataset is small.

Definition (Power)

The **power** of a hypothesis test is the probability it rejects H_0 . It depends on the alternate distribution H_A and n .

Goal of any test statistic is achieving high power*.

Observed Data, $n = 30$



What if we want to keep this data private?

Differential privacy [DMNS06]

Definition

Two databases are **neighboring** if they differ only in the data of one individual.

Differential privacy [DMNS06]

Definition

Two databases are **neighboring** if they differ only in the data of one individual.

Definition

A query f is ϵ -**differentially private** if for all neighboring databases D, D' and all output sets S

$$\Pr[f(D) \in S] \leq e^\epsilon \Pr[f(D') \in S].$$

Properties of differential privacy [DMNS06]

Theorem (Post-processing)

If f is ϵ -differentially private then for any (randomized) function g , then if $h(D) = g(f(D))$, h is also ϵ -differentially private.

Properties of differential privacy [DMNS06]

Theorem (Post-processing)

If f is ε -differentially private then for any (randomized) function g , then if $h(D) = g(f(D))$, h is also ε -differentially private.

Theorem (Composition)

If f is ε_1 -differentially private and g is ε_2 -differentially private then if $h(D) = (g(D), f(D))$, h is $(\varepsilon_1 + \varepsilon_2)$ -differentially private.

Laplace mechanism

Definition (Sensitivity)

The sensitivity Δf of a deterministic, real-valued function f on databases is the maximum over all pairs of neighboring D, D' of $|f(D) - f(D')|$.

Laplace mechanism

Definition (Sensitivity)

The sensitivity Δf of a deterministic, real-valued function f on databases is the maximum over all pairs of neighboring D, D' of $|f(D) - f(D')|$.

Theorem (Laplace Mechanism)

Given any deterministic, real-valued function f on databases, define \hat{f} as

$$\hat{f}(D) = f(D) + Y,$$

where $Y \leftarrow \text{Lap}(\Delta f / \epsilon)$. The Laplace mechanism is ϵ -differentially private.

Related Works

Other work on private hypothesis testing:

Related Works

Other work on private hypothesis testing:

- Asymptotic analysis [WZ10, Smith11, CKMSU19]

Related Works

Other work on private hypothesis testing:

- Asymptotic analysis [WZ10, Smith11, CKMSU19]
- Chi-squared test (difference of discrete distributions) [VS09, FSU11, JS13, USF13, WLK15, GLRV16, RK17]

Related Works

Other work on private hypothesis testing:

- Asymptotic analysis [WZ10, Smith11, CKMSU19]
- Chi-squared test (difference of discrete distributions) [VS09, FSU11, JS13, USF13, WLK15, GLRV16, RK17]
- Other tests:

Related Works

Other work on private hypothesis testing:

- Asymptotic analysis [WZ10, Smith11, CKMSU19]
- Chi-squared test (difference of discrete distributions) [VS09, FSU11, JS13, USF13, WLK15, GLRV16, RK17]
- Other tests:
 - ▶ Binomial data [AS18] (Proven optimal!)

Related Works

Other work on private hypothesis testing:

- Asymptotic analysis [WZ10, Smith11, CKMSU19]
- Chi-squared test (difference of discrete distributions) [VS09, FSU11, JS13, USF13, WLK15, GLRV16, RK17]
- Other tests:
 - ▶ Binomial data [AS18] (Proven optimal!)
 - ▶ Difference of two means [OHK15, DNL18]

Related Works

Other work on private hypothesis testing:

- Asymptotic analysis [WZ10, Smith11, CKMSU19]
- Chi-squared test (difference of discrete distributions) [VS09, FSU11, JS13, USF13, WLK15, GLRV16, RK17]
- Other tests:
 - ▶ Binomial data [AS18] (Proven optimal!)
 - ▶ Difference of two means [OHK15, DNL18]
 - ▶ Linear regression [BRMC17, Sheffet17]

Related Works

Other work on private hypothesis testing:

- Asymptotic analysis [WZ10, Smith11, CKMSU19]
- Chi-squared test (difference of discrete distributions) [VS09, FSU11, JS13, USF13, WLK15, GLRV16, RK17]
- Other tests:
 - ▶ Binomial data [AS18] (Proven optimal!)
 - ▶ Difference of two means [OHK15, DNL18]
 - ▶ Linear regression [BRMC17, Sheffet17]

Earlier work is often missing:

Related Works

Other work on private hypothesis testing:

- Asymptotic analysis [WZ10, Smith11, CKMSU19]
- Chi-squared test (difference of discrete distributions) [VS09, FSU11, JS13, USF13, WLK15, GLRV16, RK17]
- Other tests:
 - ▶ Binomial data [AS18] (Proven optimal!)
 - ▶ Difference of two means [OHK15, DNL18]
 - ▶ Linear regression [BRMC17, Sheffet17]

Earlier work is often missing:

- Rigorous p-value computations

Related Works

Other work on private hypothesis testing:

- Asymptotic analysis [WZ10, Smith11, CKMSU19]
- Chi-squared test (difference of discrete distributions) [VS09, FSU11, JS13, USF13, WLK15, GLRV16, RK17]
- Other tests:
 - ▶ Binomial data [AS18] (Proven optimal!)
 - ▶ Difference of two means [OHK15, DNL18]
 - ▶ Linear regression [BRMC17, Sheffet17]

Earlier work is often missing:

- Rigorous p-value computations
- Power analysis

Private F -statistic [CBRG18]

Assume data is on the $[0, 1]$ interval.

Private F -statistic [CBRG18]

Assume data is on the $[0, 1]$ interval.

Theorem

SSE has sensitivity bounded by 7.

Private F -statistic [CBRG18]

Assume data is on the $[0, 1]$ interval.

Theorem

SSE has sensitivity bounded by 7.

$$\widehat{SSE}(D) = SSE(D) + \text{Lap}(7/\epsilon)$$

Private F -statistic [CBRG18]

Assume data is on the $[0, 1]$ interval.

Theorem

SSE has sensitivity bounded by 7.

$$\widehat{SSE}(D) = SSE(D) + \text{Lap}(7/\epsilon)$$

Theorem

SSA has sensitivity bounded by $9 + 5/n$.

Private F -statistic [CBRG18]

Assume data is on the $[0, 1]$ interval.

Theorem

SSE has sensitivity bounded by 7.

$$\widehat{SSE}(D) = SSE(D) + \text{Lap}(7/\varepsilon)$$

Theorem

SSA has sensitivity bounded by $9 + 5/n$.

$$\widehat{SSA}(D) = SSA(D) + \text{Lap}\left(\frac{9 + 5/n}{\varepsilon}\right)$$

Private ANOVA [CBRG18]


$$\hat{F}(D) = \frac{\widehat{SSA}(D)/(k-1)}{\widehat{SSE}(D)/(n-k)}$$

Private ANOVA [CBRG18]

$$\hat{F}(D) = \frac{\widehat{SSA}(D)/(k-1)}{\widehat{SSE}(D)/(n-k)}$$


Problem: What is the reference distribution now?

Private ANOVA [CBRG18]



Public reference distribution gives inaccurate p -values.

Private ANOVA [CBRG18]



Definition (Power)

The **power** of a hypothesis test is the probability it rejects H_0 . It depends on the alternate distribution H_A and n .

Improving ANOVA [SHGRGB19]

Are there other ways of measuring “dispersion” (analogous to variance)?

Improving ANOVA [SHGRGB19]

Are there other ways of measuring “dispersion” (analogous to variance)?

$$(x_1 - x_2)^2, \quad |x_1 - x_2|, \text{ or maybe } |x_1 - x_2|?$$

$$SSA(D) = \sum_{j=1}^k n_j (\bar{y}_j - \bar{y})^2 \implies SA(D) = \sum_{j=1}^k n_j |\bar{y}_j - \bar{y}|$$

$$SSE(D) = \sum_{i=1}^n (y_i - \bar{y}_{c_i})^2 \implies SE(D) = \sum_{i=1}^n |y_i - \bar{y}_{c_i}|$$

$$F(D) = \frac{SSA(D)/(k-1)}{SSE(D)/(n-k)} \implies F_1(D) = \frac{SA(D)/(k-1)}{SE(D)/(n-k)}$$

Improving ANOVA [SHGRGB19]

Are there other ways of measuring “dispersion” (analogous to variance)?

$$(x_1 - x_2)^2, \quad |x_1 - x_2|, \text{ or maybe } |x_1 - x_2|?$$

$$SSA(D) = \sum_{j=1}^k n_j (\bar{y}_j - \bar{y})^2 \implies SA(D) = \sum_{j=1}^k n_j |\bar{y}_j - \bar{y}|$$

$$SSE(D) = \sum_{i=1}^n (y_i - \bar{y}_{c_i})^2 \implies SE(D) = \sum_{i=1}^n |y_i - \bar{y}_{c_i}|$$

$$F(D) = \frac{SSA(D)/(k-1)}{SSE(D)/(n-k)} \implies F_1(D) = \frac{SA(D)/(k-1)}{SE(D)/(n-k)}$$

The new F_1 statistic has:

Improving ANOVA [SHGRGB19]

Are there other ways of measuring “dispersion” (analogous to variance)?

$$(x_1 - x_2)^2, \quad |x_1 - x_2|, \text{ or maybe } |x_1 - x_2|?$$

$$SSA(D) = \sum_{j=1}^k n_j (\bar{y}_j - \bar{y})^2 \implies SA(D) = \sum_{j=1}^k n_j |\bar{y}_j - \bar{y}|$$

$$SSE(D) = \sum_{i=1}^n (y_i - \bar{y}_{c_i})^2 \implies SE(D) = \sum_{i=1}^n |y_i - \bar{y}_{c_i}|$$

$$F(D) = \frac{SSA(D)/(k-1)}{SSE(D)/(n-k)} \implies F_1(D) = \frac{SA(D)/(k-1)}{SE(D)/(n-k)}$$

The new F_1 statistic has:

- Lower sensitivity (3 for SE , 4 for SA)

Improving ANOVA [SHGRGB19]

Are there other ways of measuring “dispersion” (analogous to variance)?

$$(x_1 - x_2)^2, \quad |x_1 - x_2|, \text{ or maybe } |x_1 - x_2|?$$

$$SSA(D) = \sum_{j=1}^k n_j (\bar{y}_j - \bar{y})^2 \implies SA(D) = \sum_{j=1}^k n_j |\bar{y}_j - \bar{y}|$$

$$SSE(D) = \sum_{i=1}^n (y_i - \bar{y}_{c_i})^2 \implies SE(D) = \sum_{i=1}^n |y_i - \bar{y}_{c_i}|$$

$$F(D) = \frac{SSA(D)/(k-1)}{SSE(D)/(n-k)} \implies F_1(D) = \frac{SA(D)/(k-1)}{SE(D)/(n-k)}$$

The new F_1 statistic has:

- Lower sensitivity (3 for SE , 4 for SA)
- Much higher typical value

Improving ANOVA [SHGRGB19]

Making F_1 private

Improving ANOVA [SHGRGB19]

Making F_1 private

- ① Use Laplace mechanism on SE and SA
 - ▶ $\widehat{SA} = F_1 = SA + \text{Lap}(4/\rho\varepsilon)$
 - ▶ $\widehat{SE} = F_1 = SA + \text{Lap}(3/(1-\rho)\varepsilon)$
 - ▶ $\widehat{F}_1 = \frac{\widehat{SA}/(k-1)}{\widehat{SE}/(n-k)}$
- ② Use simulation for reference distribution
 - ▶ Problem: need standard deviation

Improving ANOVA [SHGRGB19]

Making F_1 private

- ① Use Laplace mechanism on SE and SA
 - ▶ $\widehat{SA} = F_1 = SA + \text{Lap}(4/\rho\varepsilon)$
 - ▶ $\widehat{SE} = F_1 = SA + \text{Lap}(3/(1-\rho)\varepsilon)$
 - ▶ $\widehat{F}_1 = \frac{\widehat{SA}/(k-1)}{\widehat{SE}/(n-k)}$
- ② Use simulation for reference distribution
 - ▶ Problem: need standard deviation
- ③ Private estimate of standard deviation:
 - ▶ Allocate some of epsilon budget?

Improving ANOVA [SHGRGB19]

Making F_1 private

① Use Laplace mechanism on SE and SA

- ▶ $\widehat{SA} = F_1 = SA + \text{Lap}(4/\rho\varepsilon)$
- ▶ $\widehat{SE} = F_1 = SA + \text{Lap}(3/(1-\rho)\varepsilon)$
- ▶ $\widehat{F}_1 = \frac{\widehat{SA}/(k-1)}{\widehat{SE}/(n-k)}$

② Use simulation for reference distribution

- ▶ Problem: need standard deviation

③ Private estimate of standard deviation:

- ▶ Allocate some of epsilon budget? Makes power worse

Improving ANOVA [SHGRGB19]

Making F_1 private

- ① Use Laplace mechanism on SE and SA
 - ▶ $\widehat{SA} = F_1 = SA + \text{Lap}(4/\rho\varepsilon)$
 - ▶ $\widehat{SE} = F_1 = SA + \text{Lap}(3/(1-\rho)\varepsilon)$
 - ▶ $\widehat{F}_1 = \frac{\widehat{SA}/(k-1)}{\widehat{SE}/(n-k)}$
- ② Use simulation for reference distribution
 - ▶ Problem: need standard deviation
- ③ Private estimate of standard deviation:
 - ▶ Allocate some of epsilon budget? Makes power worse
 - ▶ Solution: derive an unbiased estimator for σ

Improving ANOVA [SHGRGB19]

Making F_1 private

- ① Use Laplace mechanism on SE and SA
 - ▶ $\widehat{SA} = F_1 = SA + \text{Lap}(4/\rho\varepsilon)$
 - ▶ $\widehat{SE} = F_1 = SA + \text{Lap}(3/(1-\rho)\varepsilon)$
 - ▶ $\widehat{F}_1 = \frac{\widehat{SA}/(k-1)}{\widehat{SE}/(n-k)}$
- ② Use simulation for reference distribution
 - ▶ Problem: need standard deviation
- ③ Private estimate of standard deviation:
 - ▶ Allocate some of epsilon budget? Makes power worse
 - ▶ Solution: derive an unbiased estimator for σ

$$\hat{\sigma} = \sqrt{\pi/2} \cdot \frac{\widehat{SE}}{(N-k)}$$

Improving ANOVA [SHGRGB19]


Making F_1 private

- ① Use Laplace mechanism on SE and SA
 - ▶ $\widehat{SA} = F_1 = SA + \text{Lap}(4/\rho\varepsilon)$
 - ▶ $\widehat{SE} = F_1 = SA + \text{Lap}(3/(1-\rho)\varepsilon)$
 - ▶ $\widehat{F}_1 = \frac{\widehat{SA}/(k-1)}{\widehat{SE}/(n-k)}$
- ② Use simulation for reference distribution
 - ▶ Problem: need standard deviation
- ③ Private estimate of standard deviation:
 - ▶ Allocate some of epsilon budget? Makes power worse
 - ▶ Solution: derive an unbiased estimator for σ

$$\hat{\sigma} = \sqrt{\pi/2} \cdot \frac{\widehat{SE}}{(N-k)}$$


Note: empirically verified to have valid p -values.

Power comparison of F_1 and F_2 statistics



¹In figure: $\varepsilon = 1$, $\mu = [0.35, 0.5, 0.65]$, and $\sigma = 0.15$

Power comparison of F_1 and F_2 statistics



Optimal public test $\not\Rightarrow$ optimal private test ¹.

¹In figure: $\varepsilon = 1$, $\mu = [0.35, 0.5, 0.65]$, and $\sigma = 0.15$

Further Optimization

New Developments [CKSBG19]

Kruskal-Wallis test analogous to F-test

New Developments [CKSBG19]


Kruskal-Wallis test analogous to F-test

Modified KW test, similar methods as [SHGRGB19]

New Developments [CKSBG19]

Kruskal-Wallis test analogous to F-test

Modified KW test, similar methods as [SHGRGB19]




For 80% power, need only 23% as much data as F_1 ([SHGRGB19]) ...

New Developments [CKSBG19]

Kruskal-Wallis test analogous to F-test

Modified KW test, similar methods as [SHGRGB19]



For 80% power, need only 23% as much data as F_1 ([SHGRGB19]) ... and about 1-2% as much data as F_2 ([CBRG18])

Thank you