

Improved Differentially Private Analysis of Variance

Marika Swanberg Ira Globus-Harris Iris Griffith
Anna Ritz Andrew Bray Adam Groce

Reed College

Hypothesis Tests

Is observed data D consistent with a proposed model, H_0 (null hypothesis)?

Hypothesis Tests

Is observed data D consistent with a proposed model, H_0 (null hypothesis)?

To carry out a hypothesis test:

Hypothesis Tests

Is observed data D consistent with a proposed model, H_0 (null hypothesis)?

To carry out a hypothesis test:

- 1 Compute a *test statistic*, $t = T(D)$.

Hypothesis Tests

Is observed data D consistent with a proposed model, H_0 (null hypothesis)?

To carry out a hypothesis test:


- 1 Compute a *test statistic*, $t = T(D)$.
- 2 Compute a p -value, $p = \Pr[T(D) \geq t \mid D \leftarrow H_0]$.
 - Compute where t falls on the reference distribution for T

Hypothesis Tests

Is observed data D consistent with a proposed model, H_0 (null hypothesis)?

To carry out a hypothesis test:

- 1 Compute a *test statistic*, $t = T(D)$.
- 2 Compute a *p-value*, $p = \Pr[T(D) \geq t \mid D \leftarrow H_0]$.
 - Compute where t falls on the reference distribution for T
- 3 Compare p to a preset value α (usually .05). Reject H_0 if $p < \alpha$.



Hypothesis Tests

A good hypothesis test:

Hypothesis Tests

A good hypothesis test:

- 1 Has a known reference distribution

Hypothesis Tests

A good hypothesis test:

- 1 Has a known reference distribution
- 2 Quickly diverges from the reference distribution if H_0 is false

Hypothesis Tests

A good hypothesis test:

- ① Has a known reference distribution
- ② Quickly diverges from the reference distribution if H_0 is false

Definition (Power)

The **power** of a hypothesis test is the probability it rejects H_0 . It depends on the alternate distribution H_A and n .

Differential privacy [DMNS06]

Definition

Two databases are **neighboring** if they differ only in the data of one individual.

Differential privacy [DMNS06]

Definition

Two databases are **neighboring** if they differ only in the data of one individual.

Definition

A query f is ϵ -**differentially private** if for all neighboring databases D, D' and all output sets S

$$\Pr[f(D) \in S] \leq e^\epsilon \Pr[f(D') \in S].$$

Properties of differential privacy [DMNS06]

Theorem (Post-processing)

If f is ϵ -differentially private then for any (randomized) function g , then if $h(D) = g(f(D))$, h is also ϵ -differentially private.

Properties of differential privacy [DMNS06]

Theorem (Post-processing)

If f is ε -differentially private then for any (randomized) function g , then if $h(D) = g(f(D))$, h is also ε -differentially private.

Theorem (Composition)

If f is ε_1 -differentially private and g is ε_2 -differentially private then if $h(D) = (g(D), f(D))$, h is $(\varepsilon_1 + \varepsilon_2)$ -differentially private.

Laplace mechanism

Definition (Sensitivity)

The sensitivity Δf of a deterministic, real-valued function f on databases is the maximum over all pairs of neighboring D, D' of $|f(D) - f(D')|$.

Laplace mechanism

Definition (Sensitivity)

The sensitivity Δf of a deterministic, real-valued function f on databases is the maximum over all pairs of neighboring D, D' of $|f(D) - f(D')|$.

Theorem (Laplace Mechanism)

Given any deterministic, real-valued function f on databases, define \hat{f} as

$$\hat{f}(D) = f(D) + Y,$$

where $Y \leftarrow \text{Lap}(\Delta f / \epsilon)$. The Laplace mechanism is ϵ -differentially private.

Related Works

Other work on private hypothesis testing:

Related Works

Other work on private hypothesis testing:

- Asymptotic analysis [WZ10, Smith11, CKMSU19]

Related Works

Other work on private hypothesis testing:

- Asymptotic analysis [WZ10, Smith11, CKMSU19]
- Chi-squared test (difference of discrete distributions) [VS09, FSU11, JS13, USF13, WLK15, GLRV16, RK17]

Related Works

Other work on private hypothesis testing:

- Asymptotic analysis [WZ10, Smith11, CKMSU19]
- Chi-squared test (difference of discrete distributions) [VS09, FSU11, JS13, USF13, WLK15, GLRV16, RK17]
- Other tests:

Related Works

Other work on private hypothesis testing:

- Asymptotic analysis [WZ10, Smith11, CKMSU19]
- Chi-squared test (difference of discrete distributions) [VS09, FSU11, JS13, USF13, WLK15, GLRV16, RK17]
- Other tests:
 - ▶ Binomial data [AS18] (Proven optimal!)

Related Works

Other work on private hypothesis testing:

- Asymptotic analysis [WZ10, Smith11, CKMSU19]
- Chi-squared test (difference of discrete distributions) [VS09, FSU11, JS13, USF13, WLK15, GLRV16, RK17]
- Other tests:
 - ▶ Binomial data [AS18] (Proven optimal!)
 - ▶ Difference of two means [OHK15, DNL18]

Related Works

Other work on private hypothesis testing:

- Asymptotic analysis [WZ10, Smith11, CKMSU19]
- Chi-squared test (difference of discrete distributions) [VS09, FSU11, JS13, USF13, WLK15, GLRV16, RK17]
- Other tests:
 - ▶ Binomial data [AS18] (Proven optimal!)
 - ▶ Difference of two means [OHK15, DNL18]
 - ▶ Survival analysis [NH17]

Related Works

Other work on private hypothesis testing:

- Asymptotic analysis [WZ10, Smith11, CKMSU19]
- Chi-squared test (difference of discrete distributions) [VS09, FSU11, JS13, USF13, WLK15, GLRV16, RK17]
- Other tests:
 - ▶ Binomial data [AS18] (Proven optimal!)
 - ▶ Difference of two means [OHK15, DNL18]
 - ▶ Survival analysis [NH17]
 - ▶ Linear regression [BRMC17, Sheffet17]

Related Works

Other work on private hypothesis testing:

- Asymptotic analysis [WZ10, Smith11, CKMSU19]
- Chi-squared test (difference of discrete distributions) [VS09, FSU11, JS13, USF13, WLK15, GLRV16, RK17]
- Other tests:
 - ▶ Binomial data [AS18] (Proven optimal!)
 - ▶ Difference of two means [OHK15, DNL18]
 - ▶ Survival analysis [NH17]
 - ▶ Linear regression [BRMC17, Sheffet17]

Earlier work is often missing:

Related Works

Other work on private hypothesis testing:

- Asymptotic analysis [WZ10, Smith11, CKMSU19]
- Chi-squared test (difference of discrete distributions) [VS09, FSU11, JS13, USF13, WLK15, GLRV16, RK17]
- Other tests:
 - ▶ Binomial data [AS18] (Proven optimal!)
 - ▶ Difference of two means [OHK15, DNL18]
 - ▶ Survival analysis [NH17]
 - ▶ Linear regression [BRMC17, Sheffet17]

Earlier work is often missing:

- Rigorous p-value computations

Related Works

Other work on private hypothesis testing:

- Asymptotic analysis [WZ10, Smith11, CKMSU19]
- Chi-squared test (difference of discrete distributions) [VS09, FSU11, JS13, USF13, WLK15, GLRV16, RK17]
- Other tests:
 - ▶ Binomial data [AS18] (Proven optimal!)
 - ▶ Difference of two means [OHK15, DNL18]
 - ▶ Survival analysis [NH17]
 - ▶ Linear regression [BRMC17, Sheffet17]

Earlier work is often missing:

- Rigorous p-value computations
- Power analysis

Unique Challenges of Private Hypothesis Testing

- How to compute p -value?

Unique Challenges of Private Hypothesis Testing

- How to compute p -value?
 - ▶ Reference distribution is no longer closed-form

Unique Challenges of Private Hypothesis Testing

- How to compute p -value?
 - ▶ Reference distribution is no longer closed-form
 - ▶ Solution: simulate it
- Lots of tricky details

Unique Challenges of Private Hypothesis Testing

- How to compute p -value?
 - ▶ Reference distribution is no longer closed-form
 - ▶ Solution: simulate it
- Lots of tricky details
 - ▶ Ex: negative private estimates of standard deviation because of negative noise?

Unique Challenges of Private Hypothesis Testing

- How to compute p -value?
 - ▶ Reference distribution is no longer closed-form
 - ▶ Solution: simulate it
- Lots of tricky details
 - ▶ Ex: negative private estimates of standard deviation because of negative noise?
 - ▶ Solution: ask the statistician

Unique Challenges of Private Hypothesis Testing

- How to compute p -value?
 - ▶ Reference distribution is no longer closed-form
 - ▶ Solution: simulate it
- Lots of tricky details
 - ▶ Ex: negative private estimates of standard deviation because of negative noise?
 - ▶ Solution: ask the statistician
- Low statistical power

Unique Challenges of Private Hypothesis Testing

- How to compute p -value?
 - ▶ Reference distribution is no longer closed-form
 - ▶ Solution: simulate it
- Lots of tricky details
 - ▶ Ex: negative private estimates of standard deviation because of negative noise?
 - ▶ Solution: ask the statistician
- Low statistical power
 - ▶ Need a lot of data/impractical

Unique Challenges of Private Hypothesis Testing

- How to compute p -value?
 - ▶ Reference distribution is no longer closed-form
 - ▶ Solution: simulate it
- Lots of tricky details
 - ▶ Ex: negative private estimates of standard deviation because of negative noise?
 - ▶ Solution: ask the statistician
- Low statistical power
 - ▶ Need a lot of data/impractical
 - ▶ Main point of improvement

Particular Hypothesis Test: ANOVA

ANOVA tests independence of two variables, one continuous and one categorical.

Particular Hypothesis Test: ANOVA

ANOVA tests independence of two variables, one continuous and one categorical.

H_0 = the within-category means are all equal

Particular Hypothesis Test: ANOVA

ANOVA tests independence of two variables, one continuous and one categorical.

H_0 = the within-category means are all equal

$$F(D) = \frac{SSA(D)/(k-1)}{SSE(D)/(n-k)}$$

Particular Hypothesis Test: ANOVA

ANOVA tests independence of two variables, one continuous and one categorical.

H_0 = the within-category means are all equal

$$F(D) = \frac{SSA(D)/(k-1)}{SSE(D)/(n-k)}$$

$$SSA(D) = \sum_{j=1}^k n_j (\bar{y}_j - \bar{y})^2.$$

Particular Hypothesis Test: ANOVA

ANOVA tests independence of two variables, one continuous and one categorical.

H_0 = the within-category means are all equal

$$F(D) = \frac{SSA(D)/(k-1)}{SSE(D)/(n-k)}$$

$$SSA(D) = \sum_{j=1}^k n_j (\bar{y}_j - \bar{y})^2.$$

$$SSE(D) = \sum_{i=1}^n (y_i - \bar{y}_{c_i})^2.$$

Private ANOVA [CBRG18]

Assume data is on the $[0, 1]$ interval.

Private ANOVA [CBRG18]

Assume data is on the $[0, 1]$ interval.

Theorem

SSE has sensitivity bounded by 7.

Private ANOVA [CBRG18]

Assume data is on the $[0, 1]$ interval.

Theorem

SSE has sensitivity bounded by 7.

$$\widehat{SSE}(D) = SSE(D) + \text{Lap}(7/\epsilon)$$

Private ANOVA [CBRG18]

Assume data is on the $[0, 1]$ interval.

Theorem

SSE has sensitivity bounded by 7.

$$\widehat{SSE}(D) = SSE(D) + \text{Lap}(7/\epsilon)$$

Theorem

SSA has sensitivity bounded by $9 + 5/n$.

Private ANOVA [CBRG18]

Assume data is on the $[0, 1]$ interval.

Theorem

SSE has sensitivity bounded by 7.

$$\widehat{SSE}(D) = SSE(D) + \text{Lap}(7/\varepsilon)$$

Theorem

SSA has sensitivity bounded by $9 + 5/n$.

$$\widehat{SSA}(D) = SSA(D) + \text{Lap}\left(\frac{9 + 5/n}{\varepsilon}\right)$$

Private ANOVA [CBRG18]

$$\hat{F}(D) = \frac{\widehat{SSA}(D)/(k-1)}{\widehat{SSE}(D)/(n-k)}$$

Private ANOVA [CBRG18]

$$\hat{F}(D) = \frac{\widehat{SSA}(D)/(k-1)}{\widehat{SSE}(D)/(n-k)}$$

Problem: The reference distribution is a mess.

Private ANOVA [CBRG18]

$$\hat{F}(D) = \frac{\widehat{SSA}(D)/(k-1)}{\widehat{SSE}(D)/(n-k)}$$

Problem: The reference distribution is a mess.

- Simulate it.

Private ANOVA [CBRG18]

$$\hat{F}(D) = \frac{\widehat{SSA}(D)/(k-1)}{\widehat{SSE}(D)/(n-k)}$$

Problem: The reference distribution is a mess.

- Simulate it.

Problem: The reference distribution is no longer scale free.

Private ANOVA [CBRG18]

$$\hat{F}(D) = \frac{\widehat{SSA}(D)/(k-1)}{\widehat{SSE}(D)/(n-k)}$$

Problem: The reference distribution is a mess.

- Simulate it.

Problem: The reference distribution is no longer scale free.

- Luckily, \widehat{SSE} is a variance estimate.

Private ANOVA [CBRG18]

$$\hat{F}(D) = \frac{\widehat{SSA}(D)/(k-1)}{\widehat{SSE}(D)/(n-k)}$$


Problem: The reference distribution is a mess.

- Simulate it.

Problem: The reference distribution is no longer scale free.

- Luckily, \widehat{SSE} is a variance estimate.
- Empirically checked: good enough, type 1 error rate bounded by α

Private ANOVA [CBRG18]



Improving ANOVA [SHGRGB19]

Create a new test statistic!

Improving ANOVA [SHGRGB19]

Create a new test statistic!

$$SSA(D) = \sum_{j=1}^k n_j (\bar{y}_j - \bar{y})^2 \implies SA(D) = \sum_{j=1}^k n_j |\bar{y}_j - \bar{y}|$$

$$SSE(D) = \sum_{i=1}^n (y_i - \bar{y}_{c_i})^2 \implies SE(D) = \sum_{i=1}^n |y_i - \bar{y}_{c_i}|$$

$$F(D) = \frac{SSA(D)/(k-1)}{SSE(D)/(n-k)} \implies F_1(D) = \frac{SA(D)/(k-1)}{SE(D)/(n-k)}$$

Improving ANOVA [SHGRGB19]

Create a new test statistic!

$$SSA(D) = \sum_{j=1}^k n_j (\bar{y}_j - \bar{y})^2 \implies SA(D) = \sum_{j=1}^k n_j |\bar{y}_j - \bar{y}|$$

$$SSE(D) = \sum_{i=1}^n (y_i - \bar{y}_{c_i})^2 \implies SE(D) = \sum_{i=1}^n |y_i - \bar{y}_{c_i}|$$

$$F(D) = \frac{SSA(D)/(k-1)}{SSE(D)/(n-k)} \implies F_1(D) = \frac{SA(D)/(k-1)}{SE(D)/(n-k)}$$

The new F_1 statistic has:

Improving ANOVA [SHGRGB19]

Create a new test statistic!

$$SSA(D) = \sum_{j=1}^k n_j (\bar{y}_j - \bar{y})^2 \implies SA(D) = \sum_{j=1}^k n_j |\bar{y}_j - \bar{y}|$$

$$SSE(D) = \sum_{i=1}^n (y_i - \bar{y}_{c_i})^2 \implies SE(D) = \sum_{i=1}^n |y_i - \bar{y}_{c_i}|$$

$$F(D) = \frac{SSA(D)/(k-1)}{SSE(D)/(n-k)} \implies F_1(D) = \frac{SA(D)/(k-1)}{SE(D)/(n-k)}$$

The new F_1 statistic has:

- Lower sensitivity (3 for SE , 4 for SA)

Improving ANOVA [SHGRGB19]

Create a new test statistic!

$$SSA(D) = \sum_{j=1}^k n_j (\bar{y}_j - \bar{y})^2 \implies SA(D) = \sum_{j=1}^k n_j |\bar{y}_j - \bar{y}|$$

$$SSE(D) = \sum_{i=1}^n (y_i - \bar{y}_{c_i})^2 \implies SE(D) = \sum_{i=1}^n |y_i - \bar{y}_{c_i}|$$

$$F(D) = \frac{SSA(D)/(k-1)}{SSE(D)/(n-k)} \implies F_1(D) = \frac{SA(D)/(k-1)}{SE(D)/(n-k)}$$

The new F_1 statistic has:

- Lower sensitivity (3 for SE , 4 for SA)
- Much higher typical value

Improving ANOVA [SHGRGB19]

Making F_1 private

Improving ANOVA [SHGRGB19]

Making F_1 private

- 1 Use Laplace mechanism as in [CBRG18]

Improving ANOVA [SHGRGB19]

Making F_1 private

- ① Use Laplace mechanism as in [CBRG18]
- ② Simulate reference distribution for computing p -value
 - ▶ Problem: need standard deviation

Improving ANOVA [SHGRGB19]

Making F_1 private

- ① Use Laplace mechanism as in [CBRG18]
- ② Simulate reference distribution for computing p -value
 - ▶ Problem: need standard deviation
- ③ Private estimate of standard deviation:
 - ▶ Allocate some of epsilon budget?

Improving ANOVA [SHGRGB19]

Making F_1 private

- ① Use Laplace mechanism as in [CBRG18]
- ② Simulate reference distribution for computing p -value
 - ▶ Problem: need standard deviation
- ③ Private estimate of standard deviation:
 - ▶ Allocate some of epsilon budget? Makes power worse

Improving ANOVA [SHGRGB19]

Making F_1 private

- ① Use Laplace mechanism as in [CBRG18]
- ② Simulate reference distribution for computing p -value
 - ▶ Problem: need standard deviation
- ③ Private estimate of standard deviation:
 - ▶ Allocate some of epsilon budget? Makes power worse
 - ▶ Solution: derive an unbiased estimator for σ


Improving ANOVA [SHGRGB19]

Making F_1 private


- 1 Use Laplace mechanism as in [CBRG18]
- 2 Simulate reference distribution for computing p -value
 - ▶ Problem: need standard deviation
- 3 Private estimate of standard deviation:
 - ▶ Allocate some of epsilon budget? Makes power worse
 - ▶ Solution: derive an unbiased estimator for σ

$$\hat{\sigma} = \sqrt{\pi/2} \cdot \frac{\widehat{SE}}{(N - k)}$$

Validity of p -values



Power of F_1 test



Power comparison at $\varepsilon = 1$. F achieves 80% power with 4500 observations. F_1 requires 300.

Further Optimization

New Developments [CKSBG19]

Kruskal-Wallis test analogous to F-test

New Developments [CKSBG19]


Kruskal-Wallis test analogous to F-test

Modified KW test, similar methods as [SHGRGB19]

New Developments [CKSBG19]

Kruskal-Wallis test analogous to F-test

Modified KW test, similar methods as [SHGRGB19]




For 80% power, need only 23% as much data as F_1 ([SHGRGB19]) ...

New Developments [CKSBG19]

Kruskal-Wallis test analogous to F-test

Modified KW test, similar methods as [SHGRGB19]



For 80% power, need only 23% as much data as F_1 ([SHGRGB19]) ... and about 1-2% as much data as F_2 ([CBRG18])

Thank you