

32. Spectral methods for boundary-value problems

Last time

- Recurrence relation for Chebyshev polynomials
- Clenshaw–Curtis integration

Goals for today

- Spectral methods
- Boundary-value problems for ODEs
- Eigenvalue problems
- Time evolution

(Pseudo)-spectral methods for boundary-value problems

- Want to solve **boundary-value problems**
- E.g. Poisson equation for potential due to electric charges

(Pseudo)-spectral methods for boundary-value problems

- Want to solve **boundary-value problems**
- E.g. Poisson equation for potential due to electric charges
- Example:

$$u_{xx} = f(x) \quad \text{for } -1 < x < 1$$

with **boundary conditions** $u(-1) = u(+1) = 0$

(Pseudo)-spectral methods for boundary-value problems

- Want to solve **boundary-value problems**
- E.g. Poisson equation for potential due to electric charges
- Example:

$$u_{xx} = f(x) \quad \text{for } -1 < x < 1$$

with **boundary conditions** $u(-1) = u(+1) = 0$

- Notation: derivative $u_x := u' := \frac{du}{dx}$
- u_{xx} = 2nd derivative

Boundary-value problems

- Note that there is no time
- Everything must be satisfied *simultaneously*

Boundary-value problems

- Note that there is no time
- Everything must be satisfied *simultaneously*
- Information from *both* boundaries influences value at intermediate point

Boundary-value problems

- Note that there is no time
- Everything must be satisfied *simultaneously*
- Information from *both* boundaries influences value at intermediate point
- There are stepping methods (**shooting**) but they do not work well

Boundary-value problems

- Note that there is no time
- Everything must be satisfied *simultaneously*
- Information from *both* boundaries influences value at intermediate point
- There are stepping methods (**shooting**) but they do not work well
- Try to reduce to a linear algebra problem

Boundary-value problems

- Note that there is no time
- Everything must be satisfied *simultaneously*
- Information from *both* boundaries influences value at intermediate point
- There are stepping methods (**shooting**) but they do not work well

- Try to reduce to a linear algebra problem
- Will follow Trefethen, *Spectral Methods in MATLAB* (Chaps. 6 + 7)

Collocation methods

- We will use **collocation method** to solve
- Sometimes called “pseudo-spectral”

Collocation methods

- We will use **collocation method** to solve
- Sometimes called “pseudo-spectral”
- Extension / inverse of methods used so far

Collocation methods

- We will use **collocation method** to solve
- Sometimes called “pseudo-spectral”
- Extension / inverse of methods used so far

- Suppose that equation satisfied by function u (unknown)

Collocation methods

- We will use **collocation method** to solve
 - Sometimes called “pseudo-spectral”
 - Extension / inverse of methods used so far
-
- Suppose that equation satisfied by function u (unknown)
 - Look at collection of $N + 1$ nodes t_j
 - With *unknown* values $u_j := u(t_j)$ at t_j

Collocation methods II

- Let $p(x)$ be degree- N polynomial interpolating u_j at t_j
- Note that u_j are the *unknowns*!
- Then we *impose*

$$p''(t_j) = u_j$$

Collocation methods II

- Let $p(x)$ be degree- N polynomial interpolating u_j at t_j
- Note that u_j are the *unknowns*!
- Then we *impose*

$$p''(t_j) = u_j$$

- So

$$\mathbf{M}\mathbf{u} = \mathbf{f}$$

where $\mathbf{M} := \mathbf{D}_N^2$

- \mathbf{D}_N is differentiation matrix

Boundary conditions

- Need to impose **boundary conditions**

Boundary conditions

- Need to impose **boundary conditions**
- e.g. Dirichlet:

$$u(-1) = u(1) = 0$$

Boundary conditions

- Need to impose **boundary conditions**
- e.g. Dirichlet:

$$u(-1) = u(1) = 0$$

- Various ways to implement this
- E.g. **boundary bordering**:
 set rows 1 and N of M to rows of identity matrix
 to impose $u_0 = u_N = 0$

Boundary conditions

- Need to impose **boundary conditions**
- e.g. Dirichlet:

$$u(-1) = u(1) = 0$$

- Various ways to implement this
- E.g. **boundary bordering**:
 set rows 1 and N of M to rows of identity matrix
 to impose $u_0 = u_N = 0$
- ODE is imposed at interior points

Chebyshev spectral methods

- Formulation in terms of differentiation matrices holds for finite differences – as in Problem Set

Chebyshev spectral methods

- Formulation in terms of differentiation matrices holds for finite differences – as in Problem Set
- But Chebyshev methods have usual advantage: **spectral convergence**

Chebyshev spectral methods

- Formulation in terms of differentiation matrices holds for finite differences – as in Problem Set
- But Chebyshev methods have usual advantage: **spectral convergence**
- Additional advantage: **interpolant** is output from method

Summary

- Chebyshev spectral methods
- Collocation: Assume ODE is satisfied at interior points