

25. Approximating functions: Fourier analysis

Last time

- Conditioning of solving linear systems
- Iterative methods for solving linear systems

Goals for today

- Approximating functions
- Fourier series for periodic functions
- Rate of decay of Fourier coefficients

Manipulating functions

- Suppose that we are given a complicated function f over an interval $[a, b]$

Manipulating functions

- Suppose that we are given a complicated function f over an interval $[a, b]$
- Want to do numerical calculations with f :
 - Addition: $g = f + f$,
i.e. $g(x) = f(x) + f(x) \quad \forall x \in [a, b]$
 - Multiplication $g = \alpha f^2$, i.e. $g(x) = \alpha * f(x) * f(x)$
 - Derivative: f'
 - Definite integral: $\int_a^b f(x) dx$
 - Find roots of f in $[a, b]$

Manipulating functions

- Suppose that we are given a complicated function f over an interval $[a, b]$
- Want to do numerical calculations with f :
 - Addition: $g = f + f$,
i.e. $g(x) = f(x) + f(x) \quad \forall x \in [a, b]$
 - Multiplication $g = \alpha f^2$, i.e. $g(x) = \alpha * f(x) * f(x)$
 - Derivative: f'
 - Definite integral: $\int_a^b f(x) dx$
 - Find roots of f in $[a, b]$
- Idea: Find **approximation** of f that can manipulate

Approximating functions

- Have seen some possibilities for approximating f :

Approximating functions

- Have seen some possibilities for approximating f :
- Taylor series:
 - Polynomial that reproduces f locally near a point
 - Works well only near that point

Approximating functions

- Have seen some possibilities for approximating f :
- Taylor series:
 - Polynomial that reproduces f locally near a point
 - Works well only near that point
- Lagrange interpolation:
 - Expensive to calculate and manipulate
 - Need to be careful about location of interpolation points

Approximating functions

- Have seen some possibilities for approximating f :
- Taylor series:
 - Polynomial that reproduces f locally near a point
 - Works well only near that point
- Lagrange interpolation:
 - Expensive to calculate and manipulate
 - Need to be careful about location of interpolation points
- So far we have not discussed **periodic** functions f :

$$f(t + T) = f(t) \quad \forall t$$

T is **period** of function

Where we are heading

- We will see in this and the next lectures that:

*For **smooth** functions f , approximate using sum of **basis functions***

Where we are heading

- We will see in this and the next lectures that:

*For **smooth** functions f , approximate using sum of **basis functions***

- Will start with the case of periodic functions, which is hopefully more familiar
- Same idea will apply in non-periodic setting to approximate a function f on interval $[-1, 1]$

Where we are heading

- We will see in this and the next lectures that:

*For **smooth** functions f , approximate using sum of **basis functions***

- Will start with the case of periodic functions, which is hopefully more familiar
- Same idea will apply in non-periodic setting to approximate a function f on interval $[-1, 1]$
- Will use “linear algebra with functions” – i.e. functional analysis

Fourier series

- Consider a **periodic** function f with period 2π
- We will assume f is not “nasty”, say continuous

Fourier series

- Consider a **periodic** function f with period 2π
- We will assume f is not “nasty”, say continuous
- Can show that f can be written as

$$f(x) = \sum_{n=-\infty}^{\infty} \hat{f}_n e^{inx}$$

- i.e. sum of **basis functions** ϕ_n with $\phi_n(x) := e^{inx}$
- Alternatively

$$f(x) = a_0 + \sum_{n=1}^{\infty} [a_n \cos(nx) + b_n \sin(nx)]$$

Infinite-dimensional vector space

- We have linear combination of *infinite* number of basis functions
- Does that even make sense?

Infinite-dimensional vector space

- We have linear combination of *infinite* number of basis functions
- Does that even make sense?
- I.e. *any* (sufficiently nice) periodic function can be so expressed

Infinite-dimensional vector space

- We have linear combination of *infinite* number of basis functions
- Does that even make sense?
- I.e. *any* (sufficiently nice) periodic function can be so expressed
- Need to specify *of which space*

Hilbert space

- $L^2 := \{f : \int_0^{2\pi} |f|^2 < \infty\}$ – **square-integrable functions**

Hilbert space

- $L^2 := \{f : \int_0^{2\pi} |f|^2 < \infty\}$ – **square-integrable functions**
- Is a **Hilbert space** – complete inner product space
- Infinite-dimensional generalization of Euclidean space

Hilbert space

- $L^2 := \{f : \int_0^{2\pi} |f|^2 < \infty\}$ – **square-integrable functions**
- Is a **Hilbert space** – complete inner product space
- Infinite-dimensional generalization of Euclidean space
- Can show ϕ_n form complete basis of L^2

Hilbert space

- $L^2 := \{f : \int_0^{2\pi} |f|^2 < \infty\}$ – **square-integrable functions**
- Is a **Hilbert space** – complete inner product space
- Infinite-dimensional generalization of Euclidean space
- Can show ϕ_n form complete basis of L^2
- Eigenfunctions of Laplacian operator $\frac{\partial^2}{\partial x^2}$
- Sturm–Liouville theory

Fourier coefficients

- \hat{f}_n are **Fourier coefficients** of f . How can they be calculated?

Fourier coefficients

- \hat{f}_n are **Fourier coefficients** of f . How can they be calculated?
- Expansion is a linear combination – but an *infinite* one
- And “vectors” are functions

Fourier coefficients

- \hat{f}_n are **Fourier coefficients** of f . How can they be calculated?
- Expansion is a linear combination – but an *infinite* one
- And “vectors” are functions
- Saw that coefficients of linear combination can be easy to find

Fourier coefficients

- \hat{f}_n are **Fourier coefficients** of f . How can they be calculated?
- Expansion is a linear combination – but an *infinite* one
- And “vectors” are functions
- Saw that coefficients of linear combination can be easy to find
- When have **orthogonal vectors**

Fourier coefficients

- \hat{f}_n are **Fourier coefficients** of f . How can they be calculated?
- Expansion is a linear combination – but an *infinite* one
- And “vectors” are functions
- Saw that coefficients of linear combination can be easy to find
- When have **orthogonal vectors**
- Need to be able to talk about orthogonality of *functions*

Inner product

- Inner product **generalizes** the dot product in Euclidean space
- Abstract the properties of the dot product
- Denote as

$$(f, g) := \int_0^{2\pi} \overline{f(x)} g(x) dx$$

- Where for $z = x + iy \in \mathbb{C}$, the complex conjugate \bar{z} denotes **complex conjugate** is $\bar{z} := x - iy$
- Properties: Linear in g and $(f, f) \geq 0$
- Two functions are **orthogonal** if $(f, g) = 0$

Orthogonality

- Let's look at inner products of our basis functions:

$$\begin{aligned}
 (\phi_n, \phi_m) &= \int_0^{2\pi} e^{-inx} e^{imx} dx \\
 &= \int_0^{2\pi} e^{i(m-n)x} dx \\
 &= \frac{1}{i(m-n)} \left[e^{i(m-n)x} \right]_{x=0}^{2\pi}
 \end{aligned}$$

- So $(\phi_n, \phi_m) = 0$ for $m \neq n$ and $= 2\pi$ for $m = n$
- Hence basis functions (ϕ_n) are orthogonal

Fourier coefficients II

- We have $f = \sum_{-\infty}^{\infty} \hat{f}_n \phi_n$
- Take inner product with ϕ_m :
- Obtain

$$\begin{aligned} (\phi_m, f) &= \sum_n \hat{f}_n (\phi_m, \phi_n) \\ &= 2\pi \hat{f}(m) \end{aligned}$$

- Thus

$$\hat{f}_m = \frac{1}{2\pi} \int_0^{2\pi} e^{-imx} f(x) dx$$

Representation on computer

- Natural idea to represent periodic function on computer:

sum finite number of terms

$$f \simeq f_N := \sum_{n=-N}^N \hat{f}(n) \phi_n$$

Representation on computer

- Natural idea to represent periodic function on computer:

sum finite number of terms

$$f \simeq f_N := \sum_{n=-N}^N \hat{f}(n) \phi_n$$

- Natural question: How *many* terms should we take?
- Want to take enough terms so that neglected terms are “small”
- What does this depend on?

Representation on computer

- Natural idea to represent periodic function on computer:

sum finite number of terms

$$f \simeq f_N := \sum_{n=-N}^N \hat{f}(n) \phi_n$$

- Natural question: How *many* terms should we take?
- Want to take enough terms so that neglected terms are “small”
- What does this depend on?
- Depends on how fast $\hat{f}(n)$ *decay*

Fourier coefficients of derivative

. . .

- Suppose f is periodic and differentiable
- f' is also periodic, so can expand in Fourier series:

$$f' = \sum \widehat{f'}_n \phi_n$$

$$\text{where } \widehat{f'}_n = \frac{1}{2\pi} \int_0^{2\pi} e^{-inx} f'(x) dx$$

- Integrate by parts to get

$$\widehat{f'}_n = in \widehat{f}_n$$

Rate of decay of Fourier coefficients

- Can show: If f integrable, e.g. continuous, on interval, then

$$\hat{f}_n \rightarrow 0 \quad \text{as } n \rightarrow \infty$$

- Riemann–Lebesgue lemma

Rate of decay of Fourier coefficients

- Can show: If f integrable, e.g. continuous, on interval, then

$$\hat{f}_n \rightarrow 0 \quad \text{as } n \rightarrow \infty$$

- Riemann–Lebesgue lemma

- If $f \in C^k$ then $\widehat{f^{(k)}}_n = (in)^k \hat{f}_n \rightarrow 0$

- Hence $\hat{f}_n = o(n^{-k})$

Rate of decay II

- If f is **smooth** (C^∞) then decays faster than any polynomial
- In fact, if analytic in a suitable region then

$$\hat{f}_n = \mathcal{O}(e^{-\alpha n})$$

i.e. coefficients decay **exponentially** fast

- **Spectral convergence**

Representing functions on the computer

- This suggests that we should calculate *enough* terms

Representing functions on the computer

- This suggests that we should calculate *enough* terms
- After a while they should start decaying fast
- Soon reach machine epsilon

Representing functions on the computer

- This suggests that we should calculate *enough* terms
- After a while they should start decaying fast
- Soon reach machine epsilon
- Stop!
- We now have an excellent approximation of our general periodic function to something like machine epsilon!

Summary

- Periodic functions may be expanded in Fourier series
- Expansion in (infinite) linear combination of orthogonal basis functions
- Fourier coefficients decay at a rate determined by smoothness of f