



IMPACT OF YOGIC PRACTICES ON SELECTED BODY COMPOSITION MEASURES AND TRIGLYCERIDES AMONG OBESE MEN

Dr. S. CHIDAMBARA RAJA

Associate Professor, Dept. of Physical Education and Sports Sciences, Annamalai University.

ABSTRACT

The purpose of the study was to find out the effect of yogic practices on selected body composition measures and triglycerides among obese men. To achieve this purpose, 20 obese men, with BMI of 30-40 (W/H^2), were randomly selected as subjects from various places around Chidambaram. The age of the subjects were ranged from 30 to 35 years. The subjects were further classified at random into two equal groups of 10 subjects each, in which, group - I underwent yogic practices for six days (Monday to Saturday) per week for sixteen weeks and group - II acted as control who were not allowed to attend any special training. The selected criterion variables such as percentage of body fat, body mass index and triglycerides were measured before and after the yogic practice period. The selected criterion variables were assessed by using Deurenberg et al formula, Quetelet index and Boehringer Mannheim kit method. The collected data were statistically analysed by using Analysis of Covariance (ANCOVA). From the results of the study it was found that there was a significant reduction in percentage of body fat ($p > .05$) and body mass index ($p > .05$) and a significant increase in high density lipoprotein level ($p > .05$) after the yogic practice when compared with the control group. It was concluded from the result of the study, that yogic practice is a better tool to reduce the percentage of body fat and body mass index and increase the level of high density lipoprotein.

Key Words: Yogic Practices, Body Composition Measures, triglycerides.

INTRODUCTION

Obesity is a medical condition in which excess body fat has accumulated to the extent that it may have an adverse effect on health, leading to reduced life expectancy and/or increased health problems.[1] It is a metabolic disorder which is affecting the people throughout the world and commonly caused by a combination of excessive food energy intake, lack of physical activity, genetic susceptibility, and other psychological problems, although a few cases are caused primarily by genes, endocrine

disorders, medications or psychiatric illness.[2] The negative health (obesity) consequences are less or more insulin resistance, chances of occurring type 2 diabetes, asthma, hyper tension, increase in high total cholesterol, low density lipoproteins, triglycerides and lowering the triglycerides in blood, become sleep apnea, attaining early puberty, etc.[3] Indexes associated with high risk in obese persons often return to normal with appropriate physical activities, dietary habits, and a small weight loss even when body weight and percentage

body fat remain above recommended amounts.[4]

Men suffer a disproportionate burden of disease attribute to overweight and obesity when comparing with men and the men those who are measuring more than 30 percent of the ideal body weight are becoming obese.[5] Men with body weight measuring up to 30 percent more than that of the ideal are known as Obese. The body mass index (BMI), is the most popular and effective way of calculating Female Obesity.[5]

Those children who have BMI of above 95% percentiles are in obese.[6] More children aged 2 to 5 years are obese, as are 17 percent of children aged 6 to 19 according to the Centers of Disease Control and Prevention (CDC).[7] The primary problems for obese children are psychological or emotional.[8] It is also evident that increasing mortality rate during adolescent are due to childhood obesity.[9] A 2008 study has found that children who are obese have carotid arteries which have prematurely aged by as much as thirty years as well as abnormal levels of cholesterol.[10] The obese children were abused and teased by their same age group[11] and also by their family members quite often.[12] [13]

Yoga is a spiritual science for the integrated and holistic development of physical, mental and spiritual aspects of our well being.[14] Yoga is originated in India many thousands of years ago and it is the oldest system of personal development in the world, encompassing body, mind and spirit.[15,16] Yogsana have a deeper significant value in the development of the physical, mental and spiritual personality, whereas pure exercises only have a physical effect on

the muscles and bones.[17] Yoga poses are also designed to tone and exercise the muscles of the body to eliminate excess fat, and make it more flexible and stronger.[18] Yogic practice reduces the obesity and also reduces the risk factors associated with obesity.[19] A study shows there was a significant reduction in total cholesterol and triglycerides after twelve weeks of yoga practices.[20] Various researches suggest that yoga exercise improves the BMI of sedentary human beings including men.[21,22,23,24]

The exact body fat percentage cannot be precisely determined, but multiple methods are used to estimate it.[25] There is no single ideal percentage of body fat for everyone. Levels of body fat are epidemiologically dependent on sex and age.[26] There are many methods examine the percentage of body fat, such like, underwater body weight, skinfold test, bioelectrical impedance analysis, etc. The percentage of body fat will also be estimated with person's body mass index (BMI) by applying Deurenberg *et al* formula.[27] Body mass index (BMI) has recently gained favor as a better measure of adiposity.[28,29]

METHODOLOGY

The purpose of this study was to find out the effect of yogic practices on percentage of body fat, body mass index and triglycerides among obese men. To achieve the purpose of the present study, 20 obese men with the BMI of 95 percentile[30] or above and who were studying in various schools around Chidambaram, Tamilnadu were randomly selected as subjects. The age of the subjects were ranged from 14 to 16 years (mean age = 15.1 ± 0.3 years). All the subjects were residing at their

home, so, the food habits were not same and could not be measured. The selected subjects were divided into two equal groups of ten subjects each. Group - I considered as experimental group who underwent yogic practices for sixteen weeks, six days (Monday to Friday) per week on selected yogic exercises (appendix – I) and the same were taught by yoga teachers from School of Yoga Studies, Annamalai University, Annamalainagar, Chidambaram. Group - II considered as control that did not undergo any training programme or physical activity (either strenuous or recreational) throughout the experimental period. The data were collected on selected criterion variables such as percentage of body fat was

assessed by using Deurenberg *et al*[27,31] formula, body mass index was assessed by Quetelet index[32] and triglycerides was assessed by phosphotungstate/Mg²⁺ method, using the reagent from Boehringer Mannheim Lab, Germany[33] after taking 5 ml of blood from each subject by venous puncture method by the lab technicians, under the supervision of a qualified doctor, before and after the sixteen weeks of yogic practices as pre and post test. The blood samples were collected only after the consulting and getting permission from their parents. Analysis of covariance (ANCOVA) was applied to find out the significant difference if any between the experimental and control groups.

Table – II

Analysis of Covariance on Percentage of Body Fat Body, Body Mass Index and Triglycerides of Yogic Practice Group and Control Group

| Variable Name | Group Name | Yogic Practice Group | Control Group | 'F' Ratio |
|---|------------------------------|----------------------|------------------|-----------|
| Percentage of Body Fat (in Percentage) | Pre-test Mean ± S.D | 28.2521 ± 1.8236 | 29.6861 ± 1.893 | 2.991 |
| | Post-test Mean ± S.D. | 25.8926 ± 1.5406 | 29.8121 ± 2.126 | 18.91* |
| | Adj. Post-test Mean | 25.561 | 29.299 | 32.351* |
| Body Mass Index (kg/m²) | Pre-test Mean ± S.D | 31.8682 ± 1.3317 | 31.1816 ± 1.2628 | 0.1962 |
| | Post-test Mean ± S.D. | 29.3839 ± 0.8623 | 31.2125 ± 1.452 | 19.253* |
| | Adj. Post-test Mean | 27.896 | 31.531 | 29.632* |
| Triglycerides (mg/dl) | Pre-test Mean ± S.D | 134.47 ± 15.226 | 134.67 ± 15.159 | 0.0001 |
| | Post-test Mean ± S.D. | 130.33 ± 14.104 | 134.27 ± 17.694 | 21.844* |
| | Adj. Post-test Mean | 129.89 | 134.175 | 46.832* |

*Significant at 0.05 level of confidence.(The table values required for significance at 0.05 level of confidence for 1 and 18 & 1 and 17 are 4.41 and 4.45 respectively).

RESULTS

The collected data prior to and after the yoga practice on percentage of body fat, body mass index and high density lipoprotein were analyzed by

applying Analysis of Covariance (ANCOVA) are presented in table – II. In all the cases, .05 level of confidence was fixed to test the significance, which was considered as an appropriate.

After applying the Analysis of Covariance, the result of this study shows that there was a significant decrease in percentage of body fat (Exp. Gr. Pre-mean = 28.2521 ± 1.8236 Vs Post-mean = 25.8926 ± 1.5406 & Cont. Gr. Pre-mean = 29.6861 ± 1.893 Vs. Post-test mean = 29.8121 ± 2.126) and body mass index (Exp. Gr. Pre-mean = 31.8682 ± 1.3317 Vs. Post-mean 29.3839 ± 0.8623 & Cont. Gr. Pre-mean = 31.1816 ± 1.2628 Vs. Post-mean 31.2125 ± 1.452) and also there was a significant decrease in triglycerides (Exp. Gr. Pre-mean = 134.47 ± 15.226 Vs. Post-mean 130.33 ± 14.104 & Cont. Gr. Pre-mean = 134.67 ± 15.159 Vs. Post-mean 134.27 ± 17.694) only for yogic practice group.

Further, comparing the adjusted post-test means of the criterion variables (between yogic practice group and control group), such as percentage of body fat (Exp. Gr. = 25.561 Vs. Cont. Gr. = 29.299 & $F = 32.351$, $p < 0.05$), body mass index (Exp. Gr. = 29.896 Vs. Cont. Gr. = 31.531 & $F = 29.632$, $p < 0.05$) the yogic practice group was significantly differ with control group. Triglycerides was also significantly differ for yogic practice group (Exp. Gr. = 129.89 Vs. Cont. Gr. = 134.175 , & $F = 46.832$, $p > 0.05$) with df 1,17.

DISCUSSION

In the present study the subjects were obese children and their BMI was at 95th percentile.

1. The reduction in percentage of body fat and body mass index was significant for yogic practice group when compared with the control group.

2. There was a significant increase in high density lipoprotein cholesterol for yogic practice group when compared with the control group.

CONCLUSIONS

1. The results of the study revealed that there was a significant reduction in percentage of body fat after the yogic practice period. This result is in line with that of the study earlier conducted by **Pal et al**[34] and **Shenbagavalli and Divya**[35] found that there was a significant reduction in percentage of body fat after the yogic practice. **Ruhall, Bhandari and Chakravarti**[36] also found that there was a significant reduction in percentage of body fat after the pranayama practice.

2. The result of the study also shown that there was a significant reduction in body mass index (BMI) after the yogic practice period, when compared with the control group. The findings of **Kumari et al**[37], **Dhananjai et al**[38] and **Chen et al**[39] also found that there was a significant decrease in body mass index after the yogic practice period. **Ankad et al**[40] also found that there was a significant decrease in body mass index after the pranayama practice.

3. The result of the study shown that there was a significant decrease in triglycerides after the yogic practice period, when compared with the control group. The findings of **Telles et al**[41] found that there was a significant

decrease in triglycerides after the yogic practice period.

4. The overall result of the study shown that there was a significant reduction in percentage of body fat, body mass index and triglycerides after the experimental period. In this study, no attempt was taken to control the diet. But, in future, if the effort will be taken to control or modify the diet, the reduction in percentage of body fat, body mass index and triglycerides will be higher.

Reference:

[1] Haslam DW and James WP (2005). "Obesity". *Lancet*, 366:9492, 1197–209.

[2] Retrieved from <http://en.wikipedia.org/wiki/Obesity> on 12/07/2014.

[3] Retrieved from <http://www.obesity.org/resources-for/childhood-overweight.htm> on 15-07-2014.

[4] Aherlthy RP and Black DR (1996). "Healthy Body Weights: an Alternative Perspective". *American Journal of Clinical Nutrition*, 63, 448 – 451.

[5] Retrieved from <http://www.patientsmedical.com/men/obesity/> on 24-07-2014.

[6] Retrieved from http://kidshealth.org/kid/grow/body_stuff/bmi.html on 12-07-2014.

[7] Kimberley Wonderly, "What is a Normal BMI for Men", Retrieved from <http://www.livestrong.com/article/88045-normal-bmi-men/> on 2-06-2014.

[8] Great Britain Parliament House of Commons Health Committee (May 2004). *Obesity - Volume 1 - HCP 23-I, Third Report of session 2003-04. Report, together with formal minutes.*

London, UK: TSO (The Stationery Office). ISBN 978-0-215-01737-6. Retrieved 2007-12-17.

[9] Must A, Jacques PF, Dallal GE, Bajema CJ, Dietz WH (November 1992). "Long-term morbidity and mortality of overweight adolescents. A follow-up of the Harvard Growth Study of 1922 to 1935". *The New England Journal of Medicine*, 327:19, 1350–5.

[10] Facts for Families, "Obesity in Men Teens", Retrieved from http://www.aacap.org/App_Themes/AA_CAP/docs/facts_for_families/79_obesity_in_men_and_teens.pdf on 15-07-2014.

[11] Janssen I, Craig WM, Boyce WF, Pickett W (2004). "Associations between overweight and obesity with bullying behaviors in school-aged men". *Pediatrics*, 113:(5), 1187–94.

[12] Retrieved from <http://web.archive.org/web/20020602165343/http://obesity.org/discrimination/education.shtml> on 12-07-2014.

[13] Gurvinder Kalra, Avinash De Sousa, Sushma Sonavane and Nilesh Shah (January – June 2012). "Psychological Issues in Paediatric Obesity". *Indian Psychiatry Journal*, 21:2, 11-17.

[14] Bhavanani, A.B. (2011). "Application of Yoga Concept in the Health Improvement". In: P.Nikic, ed. Proceedings "Yoga – the Light of Microuniverse" of the International Interdisciplinary Scientific Conference "Yoga in Science – Future and Perspectives", September 23-24, 2010, Belgrade, Serbia. Belgrade: Yoga Federation of Serbia, p. 189-193.

[15] Swami Vishnu Devananda, *The Sivananda Companion to Yoga*, (New York: Fireside Book, Simon and Schuster, 2000), p. 10.

- [16] Retrieved from <http://www.minddisorders.com/Py-Z/Yoga.html> on 24-04-2012.
- [17] Chidambara Raja (November – December, 2012). “Effect of Yogic Practices on Flexibility Cholesterol and Blood Pressure”. *Online International Interdisciplinary Research Journal*, 2:6, 221-225.
- [18] Retrieved from <http://www.abc-of-yoga.com/yoga-and-health/yoga-for-men.asp> on 06-05-2012.
- [19] Dhananjai S, Sadashiv, Kumar Rajjan M.P.S. Negi and Dr. Sunita Tiwari (July 2011). “Effect of yoga practice in Management of Risk Factors with Obesity: A Pilot study”. *Indians Streams Research Journal*, 1:4, 1-7.
- [20] Mathew Ted Ballwin, “Astanga Yoga for Men – A Viable Alternative for Weight Management and Psychological Well Being”, retrieved from <http://www.healthandyoga.com/html/news/therapy/yogakids.aspx> on 12-07-2014.
- [21] T.L. Chen, H.C. Mao, C.H. Lai, C.Y. Li and C.H. Kuo, “The Effect of yoga Exercise Intervention on Health Related Physical Fitness in School – Age Asthmatic Men”. *Hu Li Za Zhi*, 56:2, (April 2009), 42-52.
- [22] Suchetha Kumari N, Damodara Gowda KM, Sukesh N, Madhu LN and Kathyayini (September 2011). “Effect of Yoga Therapy on Body Mass Index and Oxidative Status”. *Nitte University Journal of Health Science*, 1:1-3, 10-14
- [23] Balamurugan D, Ashok Kumar R and Dr. Karikalan I (January to June 2013). “Effect of Varied Yogic Practices on Body Mass Index Component of Obese Engineering College Men Students”. *Asian Journal of Physical Education and Computer Science in Sports*, 8:1, 43-44.
- [24] Yokesh TP and Chandrasekaran K. (2011). “Effect of Yogic Practice on Selected Physical Fitness among Overweighted School Men”. *Recent Research in Science and Technology*, 3:9, 43-45.
- [25] Retrieved from <http://weightloss.about.com/od/glossary/g/percentbodyfat.htm> on 6-8-2012.
- [26] Jackson, AS; Stanforth, PR; Gagnon, J; Rankinen, T; Leon, AS; Rao, DC; Skinner, JS; Bouchard, C; Wilmore, JH (2002). “The Effect of Sex, Age and Race on Estimating Percentage of Body Fat from Body Mass Index: The Heritage Family Study”. *International Journal of Obesity and Related Metabolic Disorder*, 26:6, 789–96.
- [27] Deurenber P, Weststrate JA and Seidell JC (March 1991). “Body Mass Index as a Measure of Body Fatness: Age- and Sex-Specific Prediction Formulas”. *British Journal of Nutrition*, 65:2, 105-14.
- [28] Kraemer H, Berkowitz RI and Hammer LD (1990). “Methodological Difficulties in Studies of Obesity, I: Measurement Issues”. *Ann Behav Med*. 12, 112-118.
- [29] Must A, Dallal GE and Dietz WH (1991). “Reference Data for Obesity: 85th and 95th Percentiles of Body Mass Index (wt/ht²) and Triceps Skinfold Thickness”. *American Journal of Clinical Nutrition*, 53, 839 - 846.
- [30] Retrieved from <http://www.webmd.com/parenting/raising-fit-kids/weight/kids-bmi-for-parents> on 22-06-2014.
- [31] Retrieved from <http://www.iarc.fr/en/publications/pdfs->

online/prev/handbook6/Handbook6-1.pdf on 22-6-2014.

[32] Retrieved from http://en.wikipedia.org/wiki/Body_mass_index on 21-06-2014.

[33] Pierre N.M. Demacker, Marja Hessels, Helga Toenhake-Dijkstra and Henk Baadenhuijsen, (April 1997). "Precipitation Methods for High Density Lipoprotein Cholesterol Measurement Compared, and Final Evaluation Under Routine Operating Conditions of a Method with a Low Sample-to-Reagent Ratio". *Clinical Chemistry*, 43:4, 663-668.

[34] Pal A, Srivastava N, Tiwari S, Verma NS, Narain VS, Agrawal GG, Natu SM and Kumar K (June 2011). "Effect of Yogic Practices on Lipid Profile and Body Fat Composition in Patients of Coronary Artery Disease". *Complementary Therapies in Medicine*, 19:3, 122-7.

[35] Shenbagavalli A and Divya K (2010). "The Effect of Specific Yogic Exercises and Combination of Specific Yogic Exercises with Autogenic Training on Selected Physiological Psychological and Biochemical Variables of College Men Students". *Journal of Exercise Science and Physiotherapy*, 6:2, (2010), 94-101.

[36] Ajay Singh Ruhall, Rakesh Bhandari and Ranjan Chakravarti (2010). "Effect of Kapalabhati on **FORMULAS:**

Calculating the Percentage of Body Fat:

Child Body Fat % = (1.51 x BMI) - (0.70 x Age) - (3.6 x gender) + 1.4

Where as the gender: Male = 1, Female = 0. (Retrieved from <http://www.halls.md/bmi/fat.htm> on 15-08-2012).

Calculating Body Mass Index (BMI):

The weight in kilograms divided by the square of the height in meters, used in the assessment of underweight and obesity. (Retrieved from <http://medical-dictionary.thefreedictionary.com/Quetelet+Index> on 19-05-2012).

Selected Body Composition Variables". *British Journal of Sports Medicine*, 44, i 70.

[37] N. Suchetha Kumari, K.M. Damodara Gowda, N. Sukesh, L.N. Madhu and Kathyayani, "Effect of Yoga Therapy on Body Mass Index and Oxidative Status", *Nitte University Journal of Health Science*, 1:1-3, (September 2011). 10-14.

[38] S. Dhananjai, Sadashiv, Kumar Rajjan, M.P.S. Negi and Sunita Diwari, "Effect of Yoga Practice in the Management of Risk Factors Associated with Obesity: A Pilot Study", *Indian Streams Research Journal*, 1:6, (July 2011).

[39] T.L. Chen, H.C. Mao, C.H. Lai, C.Y. Li and C.H. Kuo, "The Effect of Yoga Exercise Intervention on Health Related Physical Fitness in School-age Asthmatic Men", *Hu Li Za Zhi*, 56:2, (April 2009), 42 – 52.

[40] Roopa B. Ankad, Anita Herur, Shailaja Patil, G.V. Shashikala and Surekharani Chinnagudi, "Effect of Short-term Pranayama and Meditation on Cardiovascular Functions in Healthy Individuals", *Heart Views*, 12:2, (April – June 2011), 58 – 62.

[41] S. Telles, V.K. Naveen, A. Balakrishna and S. Kumar, "Short Term Health Impact of a Yoga and Diet Change Program on Obesity", *Med. Sci. Monit.*, 16:1, (January 2010), 35-40.

APPENDIX - I
TRAINING SCHEDULE FOR YOGIC PRATICE GROUP

| List of Yogasana & Pranayama | Weeks | Duration | Maintaining Duration | Recovery in between Yogasanas | Repetitions | Frequency | Warming up and cooling down |
|------------------------------|---------------|----------|----------------------|-------------------------------|-------------|--|-----------------------------|
| Padmasana | 1- 3 Weeks | 20 min. | 30 seconds | 30 seconds | 2 | Monday Tuesday Wednesday Thursday Friday & Saturday | 5 – 10 Minutes |
| Trikonasana | | | 30 seconds | 30 seconds | | | |
| Dhanurasana | | | 30 seconds | 30 seconds | | | |
| Shashangasana | | | 1 minute | 30 seconds | | | |
| Patchimosthasan | | | 1 minute | 30 seconds | | | |
| Meditation–Omkar. | | | 1 minutes | 30 seconds | | | |
| Pranayama – Nadisuthi | | | 1 minutes | 30 seconds | | | |
| Shavasana | | | 2 minutes | | 1 | | |
| As in previous week | 4 – 6 Weeks | 40 min | 20 minutes | | 2 | Monday Tuesday Wednesday Thursday Friday & Saturday | 5 – 10 Minutes |
| Bhujangasana | | | 1 minute | 30 seconds | | | |
| Shalabasana | | | 1 minute | 30 seconds | | | |
| Utkattasana | | | 1 minute | 30 seconds | | | |
| Gomukasana | | | 1 minute | 30 seconds | | | |
| Meditation – Omkar. | | | 1 minute | 30 seconds | | | |
| Pranayama – Sitali. | | | 1 minute | 30 seconds | | | |
| Shavasana | | | 2 minutes | | 1 | | |
| As in previous week | 7 – 9 Weeks | 66 min | 40 minutes | | 3 | Monday Tuesday Wednesday Thursday Friday & Saturday | 5 – 10 Minutes |
| Sedhupandhasan | | | 1 minutes | 30 seconds | | | |
| Matsyasana | | | 1 minutes | 30 seconds | | | |
| Uttanasana | | | 1 minute | 30 seconds | | | |
| Meditation - Omkar | | | 1 minutes | 30 seconds | | | |
| Pranayama – Bhastrika. | | | 1 minute | 1 minute | | | |
| Shavasana | | | 2 minutes | | 1 | | |
| As in previous week | 10 – 12 Weeks | 86 min | 66 minutes | | 3 | Monday Tuesday Wednesday Thursday Friday & Saturday | 5 – 10 Minutes |
| Paschimottasana | | | 1 minute | 30 seconds | | | |
| Ushatrasana | | | 1 minute | 30 seconds | | | |
| Meditation – Omkar. | | | 1 minutes | 30 seconds | | | |
| Pranayama – ujjayi. | | | 1 minute | 30 seconds | | | |
| Shavasana | | | 2 minutes | | 1 | | |