#### 1D Finite Element Analysis

## Shortcoming of the classical (variational) Ritz method

- Difficult to construct approximation functions
  - Especially in > 1 dimension
  - No systematic way of choosing them

$$u \approx u_h = \sum_{j=1}^m c_j \phi_j \quad ???$$

## Finite Element Method overcomes this shortcoming


• By...

## Finite Element Method overcomes this shortcoming


• By...

Return to day 1

- The domain of the problem is represented by a collection of simple subdomains, called *finite elements*.
  - The collection of finite elements is called the finite element mesh.



- The domain of the problem is represented by a collection of simple subdomains, called *finite elements*.
  - The collection of finite elements is called the finite element mesh.



- The domain of the problem is represented by a collection of simple subdomains, called *finite elements*.
  - The collection of finite elements is called the finite element mesh.
- Over each finite element, the physical process is approximated by functions (polynomials or otherwise) and algebraic equations relating physical quantities at selective points, called *nodes*, are developed.

$$u \approx u_h = \sum_{j=1}^n u_j \psi_j$$

- The domain of the problem is represented by a collection of simple subdomains, called *finite elements*.
  - The collection of finite elements is called the finite element mesh.
- Over each finite element, the physical process is approximated by functions (polynomials or otherwise) and algebraic equations relating physical quantities at selective points, called *nodes*, are developed.
- The element equations are *assembled* using continuity and/or "balance" of physical quantities and solved.

## Determining the interpolation (shape) functions

- The approximate solution should be:
  - Continuous over the element and differentiable (required by weak form).
  - Complete.
  - Should be an interpolant of the nodal variables of interest.

# Determining the interpolation (shape) functions (cont.)

 The interpolation (shape) functions are denoted by:

 $\psi_j$  or  $N_j$ 

 For weak form derivation of element equations the interpolation functions must satisfy the essential boundary conditions (for admissibility).