



Survey

Fine tuning type I interferon responses

Paul J. Hertzog^{a,*}, Bryan R.G. Williams^b^a Centre for Innate Immunity and Infectious Diseases, Monash Institute of Medical Research, Monash University, Clayton, Victoria, Australia^b Centre for Cancer Research, Monash Institute of Medical Research, Monash University, Clayton, Victoria, Australia

ARTICLE INFO

Article history:

Available online 25 May 2013

Keywords:

Interferons
STATs
Signal transduction
Antiviral
Innate Immunity

ABSTRACT

Interferon responses are balanced between protection against pathogens and other disease agents versus toxicity and development of chronic diseases. Optimal outcomes are achieved by regulating the nature, strength and duration of Interferon (IFN) production, IFN-receptor interaction and signalling pathways modulated in a manner appropriate for particular target cells. Modification of cell behaviour is mediated by regulation of positive and negative signalling pathways and by proteins encoded by selected groups of IFN-regulated genes. Understanding how these pathways are regulated and how to measure them by biomarkers or gene signatures will enable us to better understand the role of IFN pathways in the pathogenesis of infectious and inflammatory diseases and cancer. This will lead to improved patient stratification and disease treatment.

© 2013 Elsevier Ltd. All rights reserved.

Contents

1. Introduction	217
2. Description and production of type I IFNs	218
3. Type I IFN signalling is initiated at the receptors.	219
4. Signal transduction pathways activated via type I IFN receptors	219
5. IFN Regulated Gene Induction Facilitated by PLZF.	220
6. Negative regulation of IFN signalling	220
6.1. Receptor level	220
6.2. SOCS proteins	220
6.3. Negative regulation of transcription	221
6.4. Post-transcriptional regulation of signalling	221
7. A global view of the type I IFN response	221
8. Fine tuning the type I IFN response by feedback and feed forward mechanisms.	222
8.1. PKR	222
8.2. PRRs.	223
9. Conclusions and future directions.	223
References	223

1. Introduction

Interferons (IFNs) are a family of cytokines whose name was published in 1957 [1] based on viral interference work that began earlier in the Walter and Eliza Hall Institute in Melbourne [2]. During this period Nagano and Kojima had also described a similar phenomenon [3]. Since then, IFNs have been shown to represent a

family of cytokines that exhibit a broad range of properties beyond just antiviral action. Research in Australia in the last 30 years has made significant contributions to the field ranging from cloning one of the human isoforms (rhu IFN α 4) [4], structure function studies aimed at identifying the role of conserved amino acids [5], clinical studies in a range of diseases with Ian Mackay from WEHI [6–8], cloning and characterisation of receptors [9,10], generation of IFN receptor deficient mice [11] and negative regulation by SOCS proteins [12–14]. In this review we shall discuss how current knowledge informs us of how IFN signals are regulated or fine-tuned, which, after all is the aim of any therapeutic exploitation of

* Corresponding author.

E-mail addresses: Paul.Hertzog@monash.edu (P.J. Hertzog), Bryan.Williams@monash.edu (Bryan R.G. Williams).

these multifunctional cytokines. We will include aspects of our own work consistent with the theme of this volume on cytokine research in Australia.

Interferons are now broadly classified into three major types: Type I IFNs (including α , β , ω , ϵ), Type II (IFN γ) and Type III (IFN λ s, also known as IL28 and 29) [15,16]. They are distinguished by the specificity of their cognate receptors, their primary amino acid sequence homology, genetic locus, and stimulus and cell type of production. This review will focus on the type I IFNs, which have been the focus of our work. In broad terms, type I IFNs have pleiotropic effects on cells – they induce an antiviral state, they can inhibit cell proliferation, modulate cell fate (survival/apoptosis), differentiation and migration. Their effects can be local, at the site of production, or systemic. One major role is the regulation of the development and activity of virtually every effector cell of the innate and adaptive immune responses [17]. Because of the rapid production in response to stimuli, they are considered key effectors of innate immunity that link recruitment and sculpting of the adaptive response [18–20]. The various properties of type I IFNs make them very crucial to host defence and indeed mice that are deficient in IFN responses are susceptible to many diseases including infection (by viruses and some bacteria) and cancers [11,21,22]. However, like many cytokines, the properties of type I IFNs are a double edged sword. Excessive IFN responses are toxic as evident in mice deficient in the negative regulator of IFN signalling, SOCS1 (see below and Linossi et al., 2013). Furthermore, type I IFNs appear to contribute to the pathogenesis of autoimmune diseases such as Systemic Lupus Erythematosus (SLE) [23] and administration of IFNs to patients can result in dose limiting toxicity including side effects of flu-like symptoms, nausea, myalgia, leukopenia and neurological effects [24]. Therefore it is critical to both understanding the type I IFN system and to harnessing its therapeutic potential, that we have an appreciation of how IFN responses are finetuned by a balance of positive and negative regulation of signalling. These issues will be the focus of this review.

2. Description and production of type I IFNs

The type I IFNs are encoded by a family of genes located in a cluster on chromosome 9p in humans and the syntenic region on chromosome 4 in mouse [15]. There are 14 subtypes of IFN α that share 75–100% amino acid identity; a single IFN β subtype with 30% identity to a consensus IFN α sequence; a single IFN ϵ with 30% amino acid identity to consensus IFN α and IFN β ; IFN ω and other subtypes in certain species [15,16]. It is an unusual feature among the cytokine family that 20 or so type I IFNs all signal through the same receptor composed of IFNAR1 and IFNAR2 [25]. It begs the question of why so many subtypes exist. In this regard, it is notable that type I IFNs are rarely produced singly (some features of IFN β and IFN ϵ may be an exception), never all at once and usually in small groups (of IFN α subtypes +/-IFN β). In addition to the divergent sequences of the protein coding regions of these (mostly) single exon genes, the regulatory promoter regions of the genes in the type I IFN locus are also divergent in sequence, resulting in different regulatory elements, suggesting different control of spatio-temporal production.

In order to have a full understanding of the regulation of type I IFN signalling, it is important to consider where and when IFNs are produced. Classically, like other cytokines involved in acute responses, type I IFNs are thought to be only expressed after a stimulus, the usual stimulus being infection [18]. Thus there has been considerable recent excitement about the discovery of the growing families of pattern recognition receptors as factors that sense pathogens and drive the production of type I IFNs, as well as other cytokines and chemokines that are the effectors of the innate inflammatory response [19,26]. IFNs are induced by cell surface

and endosomal TLR signalling pathways: IFN β specifically via TLR4 stimulation by lipopolysaccharide (LPS); mixtures of IFN α subtypes and IFN β by others and usually not by TLR2, which acts in consort with TLRs 1 or 6 [18,19] (although an unusual myeloid cell type in the GI tract has been reported to produce IFN β via an atypical TLR2 pathway [27]). Cytosolic Rig I – Like Helicases (RLHs) drive IFN production after activation by viral nucleic acids via signalling pathways activated by recruitment of these receptor complexes to the mitochondria associated membranes. Other nucleic acid sensors (Aim2, DD/HX proteins, etc.) have been located in the cytosol and nucleus [28,29]. These families of pattern recognition receptors (PRRs) appear to be located at a variety of subcellular locations such that they will sense different parts of a pathogen (cell surface LPS or peptidoglycan or single or double stranded nucleic acids: RNA or DNA) that are exposed to the host cell during the course of infection. PRR recognition initiates a series of signal transduction pathways, the important one for type I IFN production appears to be the activation of one or more of the Interferon Regulatory Factor (IRF) family of transcription factors. IRF 1, 3, 5 and 7 appear to be particularly important in the induction of IFN α gene expression, whereas IRF3 is important in the induction of IFN β gene expression together with NF κ B (which is not involved in IFN α gene expression) [30].

It is generally considered that most cells in the body can produce type I IFNs, but haemopoietic cells are the predominant source and of these, plasmacytoid dendritic cells (pDC) can produce very high levels. The latter observation may be due to the high levels of constitutive IRF7 expression compared to other cells in which the expression of this transcription factor (TF) is usually low and must be primed by an initial burst of IFN β before optimal levels of other type I IFNs are produced [31]. This two-stage mechanism of type I IFN production has been demonstrated in cultured fibroblasts (and GM mice) but the extent it occurs in various cell types has not been ascertained. Indeed in a recent study, we unexpectedly demonstrated that breast cancer cells expressed IRF7, which drove type I IFN production, which, in turn, activated immune surveillance mechanisms that suppressed metastasis [32].

Breast cancer is not the only situation in which type I IFNs are produced in the absence of infection. While the type I IFN system has likely evolved to respond to pathogens, either their cell wall components or nucleic acids that are ‘exposed’ during the course of infection, the same sensing systems can detect DNA from dying cells [33], aberrant nucleic acid metabolism (degradation), which is the underlying cause of Aicardi Goutiere’s Syndrome [34,35] and antibody complexes in autoimmunity – all leading to type I IFN production and contributing to pathology [36].

IFN β is induced by CSF1 during macrophage differentiation where it has an inhibitory effect on the cell [37]. Similarly, IFN β is induced by RANKL during osteoclast generation, again inhibiting proliferation of these cells [38]. Accordingly, *Ifnar1* –/– mice have been reported to be slightly osteoporotic because of excessive numbers of osteoclasts. In both the previous situations, IFN β is produced via a pathway that activates the TF, AP1, which binds to cognate sites in the promoter. This is a distinct mechanism from the IRF3/NF κ B pathway that activates pathogen driven IFN β production. Another situation where type I IFNs are produced in the absence of pathogen stimulation is the recent observation that *Lactobacillus*, a component of commensal flora, stimulated very high production of IFN β [39]. This implies that a “physiological” production of conventional type I IFN may play a role in the “resting” mucosal organs in the maintenance of homeostasis and perhaps in priming innate protection. We recently described an even more specialised mucosal function of a type I IFN in the constitutive production of IFN ϵ in the female reproductive tract. Surprisingly the IFN ϵ promoter contains no binding elements for

IRFs and is therefore not induced by pathogens [40]. Instead it is constitutively expressed, hormone regulated (stimulated by oestrogen and suppressed by progesterone), and its expression levels fluctuate during the oestrus cycle and are switched off during pregnancy at the time of embryo implantation.

In general terms, what we believe this means is that the rules for regulating IFN production are adapted to suit the location and time. Accordingly, depending on the characteristic local influences (e.g. hormones, exposure to environment, microflora, etc.), the production and action of the type I IFN system appears to be adapted to ensure protection is provided but adequately controlled by site-specific mechanisms to ensure toxicity does not ensue.

Therefore, the first level where the type I IFN system is “finetuned” is when and where it is produced. It is important to realise that this is not only in pathological, but also “physiological” circumstances. It is interesting to note that viruses often target critical parts of the host immune response to evade it, survive and establish productive or chronic infection. Thus several viruses target the production of type I IFNs e.g. NS1 and NS3/4 proteases that act at the level of IRF activation – turning down the finetuning to suit their survival [41,42].

3. Type I IFN signalling is initiated at the receptors

All type I IFNs initiate signalling via interaction with the type I IFN components, IFNAR1 and IFNAR2 [15,16,25]. These were cloned in the 1990s as were their murine homologues [9,43–45], which reinforced the view that the murine IFN system is similar in complexity of ligands and specificity of receptors and signalling components and thus represented a good system in which to model human effects. Nineteen years ago, we and others generated mice with null mutations in the *Ifnar1* component of the receptor [11,21]. These mice demonstrated the critical importance of this system in protection from viral infection and in transducing signals from all type I IFNs including the recently described IFN ϵ (indeed this was the result that really defines this cytokine as a type I IFN) [40]. Subsequently there have been over 1,400 papers published using these mice, demonstrating a role for IFN in protection against viral infections, some bacterial infections, cancer and autoimmune diseases. Furthermore, studies with these mice have demonstrated that IFNs contribute to diseases including SOCS1 suppressed multi-organ inflammation and lethality [13], septic shock, bacterial infections (e.g. Chlamydia) [40,46], SLE type autoimmune diseases [47,48], and neurotoxicity associated with nucleic acid-sensing PRRs inducing pathogenic IFN production [33–36]. These mice have also contributed to our understanding of in vivo and cellular type I IFN signal transduction mechanisms including the existence of priming [49].

The IFNAR1 chain of the receptor that was the target for generating the aforementioned knockout mice is considered the “signal transducing chain” for all type I IFNs, whereas the IFNAR2 chain is the high affinity binding chain. However, there have been suggestions that interaction with IFNAR1 could be significant: (1) this receptor was initially cloned by virtue of its ability to bind a particular IFN α subtype and IFN β [43]; and (2) interactions with IFNAR1, in the context of a ternary complex, have been correlated with potency of activity [50]. Even before the receptors were cloned, several studies at Monash and other laboratories, investigated important residues in the ligands that influenced binding. These demonstrated the importance of cysteine residues that “pinned” the 3-dimensional structure of IFNs together [5]. These studies identified conserved hydrophilic residues such as Arg33 and Tyr123 were critical for activity, whereas other conserved residues were not [51]. These data were consistent with the observations that differences in amino acid sequences of different type I IFNs (even different IFN α subtypes) accounted for

differences in the affinity of interaction with cell surface receptors and the potency of the IFN biological effect [52]. While it is well documented that different type I IFN subtypes have different biological potencies, there are no convincing, quantitative experiments to determine that any IFN subtype has a particular activity that others do not. Based on current data, this could only occur if differences in affinity of receptor engagement can result in a different signal.

Experiments mutating residues on the ligand and on cloned receptors were extended over the years to generate an impression of what the binding interfaces on this alpha-helical cytokine were with each receptor component [52,53]. Our understanding of the fine structure of the interaction of type I IFNs with the extracellular domains of IFNAR1 and IFNAR2 were elucidated in 2012 when an X-ray crystallography characterization of the ternary complex was published by the Garcia and Schreiber groups [50]. This had implications for how different IFN subtypes might interact with the two receptor components and transduce signals, and critical interacting residues on all three proteins were predicted.

What remains to be determined is whether any cellular differences in the absolute or relative levels of *Ifnar1* and *Ifnar2* either permanently or at a particular stage of development or response occur, as has been reported for an IFN γ receptor subunit, IFNGR2, in particular T cell subsets [54].

4. Signal transduction pathways activated via type I IFN receptors.

The ternary complex formed by a type I IFN ligand engaging with the extracellular domains of IFNAR1 and IFNAR2 also bring into proximity the intracellular domains of these receptors and their associated signalling adaptors, thus enabling interactions that initiate the signalling cascade. The best characterised signalling proteins that are pre-associated with the receptor chains are Tyk2 (with IFNAR1) and JAK1 (with IFNAR2) [15,55,56]. These are members of the Janus kinase (Just Another Kinase) family that were discovered by Andrew Wilks working at the Melbourne branch of the Ludwig Institute, and have proven to be important in signalling by many cytokines [57,58]. Once activated by receptor dimerisation/multimerisation, JAK kinases phosphorylate each other and tyrosine residues on the receptors, which in turn act as docking sites for Signal Transducers and Activators of Transcription (STAT) proteins, which dock onto phosphotyrosine residues via their SH-2 domains [59]. In addition to the importance of their kinase activation of STATs, JAKs have other activities [60]. For example, the Tyk2 interaction with IFNAR1 stabilises its expression at the plasma membrane [61].

Receptor-docked STATs are phosphorylated by JAK and other kinases, then dissociate from the receptors, dimerise and translocate to the nucleus via interactions with importins, where they bind to DNA and activate transcription [59]. Type I IFN signalling was initially characterised by the formation of a specific complex called Interferon Stimulated Gene Factor 3 (ISGF3) composed of STAT1, STAT2 and IRF9 (originally called ISGF3 γ or p48). STAT1 is certainly critical to type I IFN mediated responses such as antiviral activity and there are many similarities between the phenotype of the *Ifnar1*–/– and *Stat1*–/– mice [62]. However, STAT1 is also crucial to type II IFN signalling and that of other cytokines. In addition to activation of the ISGF3 complex, type I IFNs also activate STAT3 and STAT1 to homo- and hetero-dimerise to form complexes that bind to the GAS element (Gamma Activated Sequence because it was first described as a consequence of IFN γ activation). Again these complexes are not specific to IFN signalling but are also activated by IL6 (Mansell and Jenkins, 2013) and other cytokines.

In order to understand the mechanism of IFN signalling in more detail, we and others have undertaken extensive studies to “map” the domains or residues in the cytoplasmic parts of the receptors to which signalling molecules are bound. Firstly, the JAKs are bound to membrane-proximal regions. The summary of several studies demonstrates that STAT1, 2, 3 bind to the *lfnar2* component [14,56,63–65]. Indeed mutation of tyrosine residues on IFNAR1 resulted in no change to STAT reporter activities indicating that this chain played no direct part in recruiting STAT1, 2, or 3 to the receptor complex [14]. This result is at odds with claims that recruitment of IFNAR1 to other receptors such as IFNGR and the IL6 receptor system recruits STAT1 as a mechanism of cytokine cross-talk [66,67]. Since these studies relied on the use of *lfnar1*–/– mice, it is more likely that cross talk between these receptor systems occurred by a different mechanism; that is to say, STAT1 levels are low in most cells in *lfnar1*–/– mice because of a lack of basal type I IFN priming of many IRGs such as STAT1 (the levels of STAT1 range from 2 to 3 fold below normal levels to 10 fold lower) [49]. Thus JAK/STAT pathway initiation is associated with the IFNAR2 component but requires Tyk2 on IFNAR1.

Other STATs, 4, 5 and 6 are also reportedly activated by type I IFNs, but the receptor chains to which they are associated, or even the upstream signalling pathways that activate them are unknown [59,68,69]. While there is no evidence for IFNAR1 in STAT activation, it may be linked to STAT independent signalling, such as that identified in studies of cells from STAT deficient mice. These studies indicated STAT1-dependent and -independent pathways (for both type I and II IFN signalling). While genes were activated in the absence of STAT1, the nature of the signal transduction pathways were not explored [70]. It should be noted that in addition to factors that transduce activating signals, negative regulation and regulators have also been mapped to the receptor chains (see below).

Supplementing the essential cytokine activated Jak-Stat pathways are alternate signalling pathways that are triggered by IFN receptor engagement to impart optimal transcriptional activation of IRGs and the subsequent biological responses. These include the MAPK (p38 and ERK), NFκB and PI3K/AKT pathways [68,71]. The best characterised of these is the p38 MAP kinase (MAPK) pathway, which is activated coincident with the Jak-Stat pathway [72]. The Erk MAP kinase pathway is also activated by IFN signalling and impacts on ISG mRNA translation via activation of Mnk kinases and eIF4E phosphorylation [73]. Downstream activation of the AKT/mTOR (mammalian target of rapamycin) signalling cascade is also characteristic of IFN receptor engagement but the impact is on mRNA translation of IRGs and production of encoded proteins [74,75]. While the relative contribution of these alternative pathways is likely to be cell and context dependent, many of their functions appear to be essential for IFN-dependent gene transcription, although specific transcript profiles remain to be fully characterised. Indeed a report of type I IFN signalling in T cells indicated a role for T cell receptor signalling molecules [76].

5. IFN Regulated Gene Induction Facilitated by PLZF

The zinc finger and BTB domain containing 16 (*Zbtb16*) gene encoding the protein promyelocytic leukaemia zinc finger (PLZF) is essential for antiviral innate immune responses driven by IFNs [77]. This was initially discovered by studying renal adenocarcinoma cell lines differing in their IFN sensitivity. PLZF stimulated IFN regulated genes that contained putative PLZF binding sites in their promoter regions. Many ISGs including *Oas1g*, *CXCL10*, *Rsad2* (*Viperin*), and *Ifit2* [78] failed to respond to high levels of IFN in *PLZF*–/– mice after viral infection. PLZF also regulates the effector function of NK cells since Granzyme B (*Gzmb*) induction after IFN activation in splenic isolated NK cells from PLZF deficient mice is impaired. A reduction in IFN-

induced CXCL10 expression in NK cell-rich organs is also seen in these mice and likely contributes to their susceptibility to viral infection. The *Zbtb16* gene encoding PLZF is not itself regulated by IFN but instead, PLZF is phosphorylated following IFN signalling. IFN mediates phosphorylation of PLZF at serine residue 76 in vitro via Mitogen-activated protein kinase 8 (JNK) or a kinase in the same pathway. This phosphorylation of PLZF enhances its transcriptional activity and facilitates binding of PLZF to cognate sites in select ISGs. Thus, following IFN stimulation, PLZF acted as a transcriptional activator, a novel finding for this zinc finger protein family member. This activity of PLZF contrasts with its well-described role as a transcriptional repressor [79].

6. Negative regulation of IFN signalling

In order to regulate type I IFN signalling and prevent toxicity, there is negative regulation at virtually every step of the signal transduction process: at the level of ligand production, as mentioned above, at the level of receptor, signalling adaptor, signalling enzymes, TFs, co-regulators of transcription, and post-transcriptional and post-translational modifications, as well as inhibitors of the proteins encoded by IFN regulated genes. The effect of these negative regulators can be to limit the strength or duration of signalling.

6.1. Receptor level

The gene encoding IFNAR2, the primary binding chain, can undergo alternative splicing to generate a truncated transmembrane isoform with no apparent signalling capability; this may act as a decoy receptor, thus inhibiting the interaction of ligand with productive signalling receptors [16,25,44,45]. Both human and murine IFNAR2 genes can be alternatively spliced to form a soluble receptor [9,44]. There has been a report that this may also be produced by protease cleavage by ADAM 17 (TACE) [80]. While it is possible that this soluble receptor might inhibit signalling, as is the case in some cytokine systems, this has not yet been demonstrated for the type I IFN. An alternative possibility is that soluble IFNAR2 might engage in trans-signalling (as defined in case of IL6 soluble receptor), whereby the soluble receptor binds ligand and presents it to the signal transducing chain, in this case, IFNAR1 (gp130 in the case of IL6), resulting in the transduction of particular signals [81]. The *lfnar1* component of the receptor also contains a negative regulatory domain that is the target for ubiquitination and degradation [82].

Proteins associated with the intracellular domains of the receptors can also suppress signalling. These include UBP18/43 associated with IFNAR2 [83] and phosphatases that are associated with IFNAR [84,85]. Uncharacterised activities also exist, such as: reports of a negative regulatory domain in the C terminal region of IFNAR1 [86], and a negative impact of the IFNAR2 chain on GAB2 signalling from IFNAR1 [87].

6.2. SOCS proteins

We have recently mapped the binding of the negative regulator SOCS1 to the IFNAR1 component of the type I IFN receptor [14]. The Suppressors Of Cytokine Signalling (SOCS proteins) were discovered concurrently by three groups including the Hilton laboratory at Walter and Eliza Hall Institute in Melbourne (Linossi et al., 2013). There are 7 members of this family that function to inhibit signalling by a range of cytokines and growth factors (Linossi et al., 2013; Mansell and Jenkins, 2013). SOCS proteins are composed of three key domains: KIR, SH2 and SOCS BOX. These domains mediate interaction with their target proteins: they variously bind to phosphor-tyrosine residues on receptors (e.g. IFNGR and gp130)

or to JAK kinases and target them for degradation by the proteasome. Of these, SOCS1 and SOCS3 are well characterised to inhibit type I IFN signalling. The *in vivo* role of these proteins has been defined in gene-targeted mice. SOCS1 mice die pre-weaning due to multi-organ inflammation [12]. This lethality could be rescued by crossing these mice to IFN γ -deficient mice, which is a vivid demonstration of the dangers of unrestricted cytokine signalling [12]. The *Socs1*^{−/−} lethality at 3 weeks could also be rescued by crossing to *lfnar1*^{−/−} mice demonstrating that unrestricted type I IFN signalling was also pro-inflammatory and potentially lethal [13]. Interestingly, mice were not rescued by crossing to *lfnar2*^{−/−} mice, which demonstrated that SOCS1 probably associated with the *lfnar1* chain of the receptor (and also that there were potentially phenotypic differences between these two receptor deficient mice). We went on, in collaboration with the WEHI scientists, to demonstrate SOCS1 association with *lfnar1* by co-immunoprecipitation [13]; we then demonstrated that SOCS1 did not associate with phosphotyrosine residues in the receptor (as it did in the IFN γ and IL6 receptor systems), but with the Tyk2 kinase which was associated with the juxta-membrane region of the intracellular domain of *lfnar1* [14].

6.3. Negative Regulation of Transcription

Type I IFN signalling is also negatively regulated at the level of transcription factors by interacting proteins, such as Protein Inhibitors of Activated STATs (PIAS) [88], Akt pathway activation of EMSY [89], TRIM pathways acting via retinoic acid receptor (RAR) α [90], and by several members of the IRF family that can act as negative regulators [91]. Type I IFN signalling also up-regulates expression of Sprouty (Spry) proteins 1, 2, and 4. This is Mnk kinase-dependent and results in suppressive effects on the IFN-activated p38 MAP kinase (MAPK), and consequent suppression of transcription of ISGs [92]. The dual roles of some transcriptional regulators like PLZF are interesting, because while it is a positive regulator of type I IFN signalling, it is at other times, a negative regulator of primary signalling [77–79].

6.4. Post-transcriptional regulation of signalling

The most notable negative regulators of signalling that are being investigated at present are the non-coding RNA, especially micro-RNA (miR) [93]. There are about 800 confirmed genes encoding functional miRs in the human and mouse genome and estimates range up to 10,000s [94]. The genes encode short sequences of RNA that bind Argonaute proteins, which direct the duplexes into the RISC complex where transcripts are degraded. MiR targeting of protein coding sequences thus decreases the half-life of transcripts and the encoded proteins [95]. There is considerable characterisation of the role of miRs in the immune response [96] and cancer [97] but little documentation of their role in regulating innate immune responses and the IFN response in particular, where only a few have been reported [98–100]. The targets include SOCS1 by miR-155 involving auto-regulation of this negative feedback loop [101]; and of *lfnar1* by miR-29a involving thymic involution [102]. There is sufficient data on the regulation of miR transcription to suggest that regulation will be similar to that of protein-coding genes. Therefore the pathways discussed here for regulating the expression of protein-coding IRGs are likely to also transcriptionally regulate miRs, providing another level of networked finetuning.

7. A global view of the type I IFN response

The end result of signalling emanating from an IFN ligand interacting with cognate receptors and initiating various signal transduction pathways is the activation of transcription factors

that activate or repress the expression of (sets of) IFN regulated genes (IRGs). It is the proteins encoded by these IRGs that are the effectors of the IFN response. They will elicit protection from viral infection, inhibition of cell proliferation, modulation of immune cell functions, etc. One way to obtain a global view of the IFN response is to perform a microarray analysis of changes in gene expression. The first (using the Affymetrix platform) microarray experiments of an IFN response were performed by the Williams and Silverman laboratories at the Lerner Research Institute, Cleveland Clinic [103,104]. These experiments detected over 1000 genes with altered expression and gene ontology analyses (GO) could identify sets of genes associated with known IFN functions including antiviral activity, cell cycle regulation, survival/apoptosis, immune response, etc. [104]. Ten years later, we decided to collect all of the experiments that had been performed where a cell, mouse or human had been treated with IFN and microarrays performed to measure changes in gene expression (Interferome.org [105]). The tally of IFN regulated genes was then about 2000 (in mouse and human). There was some overlap in genes induced by types I, II and III IFNs, consistent with our knowledge of common signalling pathways [106]. As well, there were a significant number of distinct genes regulated by type I IFNs consistent with our knowledge that these IFNs activate specific pathways.

Having this resource provides a database of the full potential scope of IFN signalling. Analysis of these genes and the pathways leading to their activation is providing a broader view of the IFN response. Analysis of promoter regulatory regions of the IRGs reveals patterns that suggest that sets of genes are co-regulated. We have begun annotating the patterns and pathways and recently have published an updated version of the database with more quantitative analysis of data, thorough annotation of metadata and additional tools such as regulatory regions and tissue expression [107]. This approach has been successfully used to identify subsets of IRGs or pathological significance and the upstream regulatory pathway.

Analysis of transcriptional changes that occur in HIV infected pDC and subsequent Interferome analysis identified that despite inhibition of IFN production, a common feature of viral infection was that a subset of IRGs were induced. Subsequent analysis of the promoters of these genes identified an enrichment of IRF1 sites. These were functionally validated by ChIP experiments. Interestingly, the genes induced by this HIV activated pathway were annotated in GO as anti-apoptotic or pro-survival [108]. Another example of a functional subset of IRGs being activated was our recent study of breast cancer metastases to bone, in which we identified a subset of 540 IRGs that were enriched for IRF7 sites; the gene set and this TF were expressed in the primary tumour cells but were suppressed in metastatic tumour cells [32]. This pathway was demonstrated to activate immune responses to metastasising tumour cells: suppression was necessary to enable metastases to bone, whereas reversal of this process inhibited breast cancer metastases to bone.

One general principle evident from these and other studies is that no situation leads to the induction of all potential IFN regulated genes. Thus, the IFN response is finetuned by the activation of some, but not all possible IFN regulated pathways. It will be important to elucidate why this is if we are to understand and therapeutically manipulate this response. It is the nature, strength, duration, cellular context and activation or repression of these pathways that determine the ultimate response: on the one hand, resolution of infection, repair or healing and return to homeostasis; versus acute disease (e.g. septic shock), chronic disease or aberrant immune response and autoimmunity. We are yet to learn what determines go/no go, strong/weak, short/long signals for each IFN regulated pathway. The selective activation of

pathways likely reflects different IFN subtypes, different cell types containing different levels of signalling factors and potentially expressing different levels of receptor components on the cell surface, as well as the amount and duration of IFN exposure. This different ability of different cells to respond was elegantly illustrated in a recent report of experiments examining a population of cells that responded to a viral stimulus resulting in the production of type I IFNs. It was shown that only a low proportion of cells produced IFN. This was because when the levels of signalling components required for this pathway were measured, most cells were deficient in one or more factors and therefore did not respond. Only cells that contained relatively high levels of all components of the signalling pathway produced IFN [109]. This very important observation is likely to explain why basal IFN is produced as a primer of certain IRGs that are required for an optimal response to a danger signal.

8. Fine tuning the type I IFN response by feedback and feed forward mechanisms

We have presented several scenarios that demonstrate that type I IFN may be produced in the absence of an overt pathological stimulus. This included myeloid and osteoclast cell development, gastrointestinal tract microbiota, reproductive tract epithelium and breast cancer and normal epithelial cells. In the latter situation we have not identified an extracellular IFN stimulus, but the IFN regulator, IRF7, is expressed, as are 540 of its potential regulated

genes; the impact on the immune response in this case is profound. This “physiological” production is likely to be important in priming the IFN system to ensure the levels of critical signalling factors are maintained at a sufficient level to ensure a robust response. Consistent with this thesis, we and others [110] have demonstrated that gene expression in “resting” cells from *Ifnar1* null mice is different from wild type. The expression levels of 100s of (IFN regulated) genes show lower expression, presumably because of the inability of these cells to respond to endogenous IFN production. Among these genes are several, such as STAT1, that are involved in the IFN response. However it must also be remembered that this auto-regulation by IFN of its own response, can be inhibitory as reviewed above for the IFN induced inhibitors, SOCS1 and miRs. Nevertheless, the constitutive IFN does prime positively for its own response because cells with higher levels of signalling components will signal better, while others with low levels will not [109]. Some of the more interesting IRGs are sensors themselves, such as PKR and PRRs (Fig. 1).

8.1. PKR

Protein Kinase R (PKR) is an interferon regulated gene product that is constitutively expressed in all differentiated cells as an inactive protein monomer. It is an effector molecule of the innate immune system that responds to cellular stress, most typically exemplified by viral infection. Viral RNA replication intermediates, or protein activators induced by different stressors, trigger

Fine Tuning Type I Interferon Responses




Fig. 1. Fine tuning of type I IFN responses. The figure shows a diagrammatic representation of major factors involved in the production of type I IFNs α and β by upstream pattern recognition receptors such as TLRs, RLHs, STING and DDX which sense pathogens and other stimuli in different subcellular compartments. They activate IF production by phosphorylation of IRF family members by the kinases Tbk or IKK ϵ . IFN acts via its receptors to activate STAT pathways or alternative pathways such as PLZF which in turn contribute to the activation (or repression) of IFN regulated genes (IRGs). Regulation of PRRs constitutes positive feedback loop for its own production (reverse arrows). IFN primes its own signalling by regulating factors such as STAT1. Moreover IFN induces negative regulators (in red) such as SOCS1 and SOCS3, which act with others such as UBP18/45 and PIAS to dampen the response.

dimerisation and autophosphorylation to generate the active enzyme [111–114]. Activated PKR inhibits protein translation through phosphorylation of eukaryotic initiation factor 2 α (eIF2 α). PKR has been shown to modulate different cell-signalling pathways including those controlled by the c-Jun Terminal Kinase (JNK) and nuclear factor- κ B (NF κ B) [115,116]. The precise mechanism by which PKR effects JNK and NF κ B has not been established but, in the case of NF κ B, appears to be independent of kinase activity [117]. In this case PKR is likely playing a scaffolding role interacting with the adaptor TRAF6 [118]. Recently a role has been proposed for PKR control of the actin cytoskeleton [119]. PKR associates with the key actin regulatory protein Gelsolin (GSN) and regulates its activity. Ablation of PKR affects actin dynamics and GSN-dependent processes that include cell motility, membrane ruffling, endocytosis, and phagocytosis. This identifies a mechanism where the innate immune effector protein PKR controls the actin cytoskeleton by keeping GSN in an inactive conformation. This constitutes a novel means by which the innate immune protein PKR functions in response to pathogens. PKR bolsters innate immunity by limiting pathogen entry into fibroblasts, while increasing macrophage-mediated clearance of infected cells. While the association between PKR and GSN is disrupted by activation of PKR, the active enzyme triggers cell signalling, as described above, and phosphorylates eIF2 α limiting viral protein production.

8.2. PRRs

Many PRRs (TLRs, STING, RLHs, Aim2, DNA sensors) are also strongly IFN regulated. Indeed in some cases, this property was an element of their discovery. Some PRR signalling adaptors are IRGs as are some members of the IRF family of TFs, especially IRF7. The constitutive production of type I IFN (or the constitutive activation of certain pathways) thus primes the system responsible for its production to ensure optimal levels are produced in response to a danger signal such as infection. IFN also thus primes the levels of components necessary for its own production.

9. Conclusions and future directions

We have reviewed herein the state-of-the art knowledge on fine tuning type I IFN responses. The balance between a protective response, without harmful effects and a return to homeostasis is offset against the risk of infection (acute and dangerous such as sepsis or chronic infectious disease with sequelae such as cancer in the case of hepatitis B/C), toxicity and autoimmune disease, and is achieved by tightly controlled positive and negative signals. Control occurs at every level from the machinery for stimulating IFN production to its interaction with cognate receptors and transducing signals in the responding cell. We have more yet to learn about the nature and control of IFN signalling, particularly at the global level of induction and suppression of 1000s of genes in signature sets which encode important effector proteins. Much of this new information will come from the application of next generation sequencing technologies and of studies of non-coding RNAs and their role in regulation and as potential biomarkers of disease or responses. Knowledge of fine tuning of the production and response to cytokines such as type I IFNs will be a key to translating this knowledge into clinical application in diagnosing disease, therapeutic use or blocking of their action or tailoring treatments to more responsive patient groups.

References

- [1] Isaacs A, Lindenmann J. Virus interference. I. The interferon. *Proceedings of the Royal Society of London Series B Biological Sciences* 1957;147:258–67.
- [2] Isaacs A, Edney M. Interference between inactive and active influenza viruses in the chick embryo. *The Australian Journal of Experimental Biology and Medical Science* 1950;28:635–45.
- [3] Nagano Y, Kojima Y. Pouvoir immunisant du virus vaccinal inactivé par des rayons ultraviolets. *Comptes Rendus des Séances de la Société de Biologie et de ses Filiales* 1954;148:1700–2.
- [4] Linnane AW, Beilharz MW, McMullen GL, Macreadie IG, Murphy M, Nisbet IT, et al. Nucleotide sequence and expression in *E. coli* of a human interferon-alpha gene selected from a genomic library using synthetic oligonucleotides. *Biochemistry International* 1984;8:725–32.
- [5] Nisbet IT, Beilharz MW, Hertzog PJ, Tymms MJ, Linnane AW. Single amino acid substitutions at conserved residues of human interferon-alpha can effect antiviral specific activity. *Biochemistry International* 1985;11:301–9.
- [6] Hertzog PJ, Emery P, Cheetham BF, Mackay IR, Linnane AW. Interferons in rheumatoid arthritis: alterations in production and response related to disease activity. *Clinical Immunology and Immunopathology* 1988;48:192–201.
- [7] Hertzog PJ, Wright A, Harris G, Linnane AW, Mackay IR. Intermittent interferonemia and interferon responses in multiple sclerosis. *Clinical Immunology and Immunopathology* 1991;58:18–32.
- [8] Johns TG, Mackay IR, Callister KA, Hertzog PJ, Devenish RJ, Linnane AW. Antiproliferative potencies of interferons on melanoma cell lines and xenografts: higher efficacy of interferon beta. *Journal of the National Cancer Institute* 1992;84:1185–90.
- [9] Owczarek CM, Hwang SY, Holland KA, Gulluyan LM, Tavaría M, Weaver B, et al. Cloning and characterization of soluble and transmembrane isoforms of a novel component of the murine type I interferon receptor, IFNAR 2. *The Journal of Biological Chemistry* 1997;272:23865–70.
- [10] Hertzog PJ, Hwang SY, Holland KA, Tymms MJ, Iannello R, Kola I. A gene on human chromosome 21 located in the region 21q22.2 to 21q22.3 encodes a factor necessary for signal transduction and antiviral response to type I interferons. *The Journal of Biological Chemistry* 1994;269:14088–93.
- [11] Hwang SY, Hertzog PJ, Holland KA, Sumarsono SH, Tymms MJ, Hamilton JA, et al. A null mutation in the gene encoding a type I interferon receptor component eliminates antiproliferative and antiviral responses to interferons alpha and beta and alters macrophage responses. *Proceedings of the National Academy of Sciences of the United States of America* 1995;92:11284–88.
- [12] Alexander WS, Starr R, Fenner JE, Scott CL, Handman E, Sprigg NS, et al. SOCS1 is a critical inhibitor of interferon gamma signaling and prevents the potentially fatal neonatal actions of this cytokine. *Cell* 1999;98:597–608.
- [13] Fenner JE, Starr R, Cornish AL, Zhang JG, Metcalf D, Schreiber RD, et al. Suppressor of cytokine signaling 1 regulates the immune response to infection by a unique inhibition of type I interferon activity. *Nature Immunology* 2006;7:33–9.
- [14] Piganis RA, De Weerd NA, Gould JA, Schindler CW, Mansell A, Nicholson SE, et al. Suppressor of cytokine signaling (SOCS) 1 inhibits type I interferon (IFN) signaling via the interferon alpha receptor (IFNAR1)-associated tyrosine kinase Tyk2. *The Journal of Biological Chemistry* 2011;286:33811–18.
- [15] Pestka S, Krause CD, Walter MR. Interferons, interferon-like cytokines, and their receptors. *Immunological Reviews* 2004;202:8–32.
- [16] de Weerd NA, Samarajiwa SA, Hertzog PJ. Type I, interferon receptors: biochemistry and biological functions. *The Journal of Biological Chemistry* 2007;282:20053–57.
- [17] Hervas-Stubbs S, Perez-Gracia JL, Rouzaut A, Sanmamed MF, Le Bon A, Melero I. Direct effects of type I interferons on cells of the immune system. *Clinical Cancer Research An Official Journal of the American Association for Cancer Research* 2011;17:2619–27.
- [18] Hertzog PJ, O'Neill LA, Hamilton JA. The interferon in TLR signaling: more than just antiviral. *Trends in Immunology* 2003;24:534–9.
- [19] Noppert SJ, Fitzgerald KA, Hertzog PJ. The role of type I interferons in TLR responses. *Immunology and Cell Biology* 2007;85:446–57.
- [20] Gonzalez-Navajas JM, Lee J, David M, Raz E. Immunomodulatory functions of type I interferons. *Nature Reviews Immunology* 2012;12:125–35.
- [21] Muller U, Steinhoff U, Reis LF, Hemmi S, Pavlovic J, Zinkernagel RM, et al. Functional role of type I and type II interferons in antiviral defense. *Science* 1994;264:1918–21.
- [22] Swann JB, Hayakawa Y, Zerafa N, Sheehan KC, Scott B, Schreiber RD, et al. Type I, IFN contributes to NK cell homeostasis, activation, and antitumor function. *Journal of Immunology* 2007;178:7540–9.
- [23] Crow MK. Type I, interferon in systemic lupus erythematosus. *Current Topics in Microbiology and Immunology* 2007;316:359–86.
- [24] Jonasch E, Haluska FG. Interferon in oncological practice: review of interferon biology, clinical applications, and toxicities. *The Oncologist* 2001;6:34–55.
- [25] de Weerd NA, Nguyen T. The interferons and their receptors – distribution and regulation. *Immunology and Cell Biology* 2012;90:483–91.
- [26] Kawai T, Akira S. The role of pattern-recognition receptors in innate immunity: update on Toll-like receptors. *Nature Immunology* 2010;11:373–84.
- [27] Barbalat R, Lau L, Locksley RM, Barton GM. Toll-like receptor 2 on inflammatory monocytes induces type I interferon in response to viral but not bacterial ligands. *Nature Immunology* 2009;10:1200–7.
- [28] Keating SE, Baran M, Bowie AG. Cytosolic DNA sensors regulating type I interferon induction. *Trends in Immunology* 2011;32:574–81.
- [29] Cavlart T, Ablasser A, Hornung V. Induction of type I IFNs by intracellular DNA-sensing pathways. *Immunology and Cell Biology* 2012;90:474–82.

- [30] Tamura T, Yanai H, Savitsky D, Taniguchi T. The IRF, family transcription factors in immunity and oncogenesis. *Annual Review of Immunology* 2008; 26:535–84.
- [31] Prakash A, Smith E, Lee CK, Levy DE. Tissue-specific positive feedback requirements for production of type I interferon following virus infection. *The Journal of Biological Chemistry* 2005;280:18651–57.
- [32] Bidwell BN, Slaney CY, Withana NP, Forster S, Cao Y, Loi S, et al. Silencing of *Irf7* pathways in breast cancer cells promotes bone metastasis through immune escape. *Nature Medicine* 2012.
- [33] Okabe Y, Kawane K, Akira S, Taniguchi T, Nagata S. Toll-like receptor-independent gene induction program activated by mammalian DNA escaped from apoptotic DNA degradation. *The Journal of Experimental Medicine* 2005; 202:1333–9.
- [34] Crow YJ, Hayward BE, Parmar R, Robins P, Leitch A, Ali M, et al. Mutations in the gene encoding the 3'–5' DNA exonuclease TREX1 cause Aicardi-Goutieres syndrome at the AGS1 locus. *Nature Genetics* 2006;38:917–20.
- [35] Crow YJ, Leitch A, Hayward BE, Garner A, Parmar R, Griffith E, et al. Mutations in genes encoding ribonuclease H2 subunits cause Aicardi-Goutieres syndrome and mimic congenital viral brain infection. *Nature Genetics* 2006; 38:910–6.
- [36] Barbalat R, Ewald SE, Mouchess ML, Barton GM. Nucleic acid recognition by the innate immune system. *Annual Review of Immunology* 2011;29:185–214.
- [37] Hamilton JA, Whitty GA, Kola I, Hertzog PJ. Endogenous IFN- α beta suppresses colony-stimulating factor (CSF)-1-stimulated macrophage DNA synthesis and mediates inhibitory effects of lipopolysaccharide and TNF- α . *Journal of Immunology* 1996;156:2553–7.
- [38] Takayanagi H, Kim S, Taniguchi T. Signaling crosstalk between RANKL and interferons in osteoclast differentiation. *Arthritis Research* 2002;4(Suppl 3):S227–32.
- [39] Weiss G, Maaetoft-Udsen K, Stifter SA, Hertzog P, Goriely S, Thomsen AR, et al. MyD88 drives the IFN- β response to *Lactobacillus acidophilus* in dendritic cells through a mechanism involving IRF1, IRF3, and IRF7. *Journal of Immunology* 2012;189:2860–8.
- [40] Fung KY, Mangan NE, Cumming H, Horvat JC, Mayall JR, Stifter SA, et al. Interferon- ϵ protects the female reproductive tract from viral and bacterial infection. *Science* 2013;339:1088–92.
- [41] Garcia-Sastre A. Induction and evasion of type I interferon responses by influenza viruses. *Virus Research* 2011;162:12–8.
- [42] Taylor KE, Mossman KL. Recent advances in understanding viral evasion of type I interferon. *Immunology* 2013;138:190–7.
- [43] Uze G, Lutfalla G, Gresser I. Genetic transfer of a functional human interferon α receptor into mouse cells: cloning and expression of its cDNA. *Cell* 1990;60:225–34.
- [44] Novick D, Cohen B, Rubinstein M. The human interferon α /beta receptor: characterization and molecular cloning. *Cell* 1994;77:391–400.
- [45] Lutfalla G, Holland SJ, Cinato E, Monneron D, Reboul J, Rogers NC, et al. Mutant USA cells are complemented by an interferon- α beta receptor subunit generated by alternative processing of a new member of a cytokine receptor gene cluster. *The EMBO Journal* 1995;14:5100–8.
- [46] Monroe KM, McWhirter SM, Vance RE. Induction of type I interferons by bacteria. *Cellular Microbiology* 2010;12:881–90.
- [47] Nacionales DC, Kelly-Scumpia KM, Lee PY, Weinstein JS, Lyons R, Sobel E, et al. Deficiency of the type I interferon receptor protects mice from experimental lupus. *Arthritis and Rheumatism* 2007;56:3770–83.
- [48] Thibault DL, Graham KL, Lee LY, Balboni I, Hertzog PJ, Utz PJ. Type I, interferon receptor controls B-cell expression of nucleic acid-sensing Toll-like receptors and autoantibody production in a murine model of lupus. *Arthritis Research & Therapy* 2009;11:R112.
- [49] Gough DJ, Messina NL, Hii L, Gould JA, Sabapathy K, Robertson AP, et al. Functional crosstalk between type I and II interferon through the regulated expression of STAT1. *PLoS Biology* 2010;8:e1000361.
- [50] Thomas C, Moraga I, Levin D, Krutzik PO, Podoplelova Y, Trejo A, et al. Structural linkage between ligand discrimination and receptor activation by type I interferons. *Cell* 2011;146:621–32.
- [51] Tymms MJ, McInnes B, Waine GJ, Cheetham BF, Linnane AW. Functional significance of amino acid residues within conserved hydrophilic regions in human interferons- α . *Antiviral Research* 1989;12:37–47.
- [52] Jaks E, Gavutis M, Uze G, Martal J, Piehler J. Differential receptor subunit affinities of type I interferons govern differential signal activation. *Journal of Molecular Biology* 2007;366:525–39.
- [53] Cajean-Feroldi C, Nosal F, Nardeux PC, Gallet X, Guymarho J, Baychelier F, et al. Identification of residues of the IFNAR1 chain of the type I human interferon receptor critical for ligand binding and biological activity. *Biochemistry* 2004;43:12498–512.
- [54] Bach EA, Szabo SJ, Dighe AS, Ashkenazi A, Aguet M, Murphy KM, et al. Ligand-induced autoregulation of IFN- γ receptor beta chain expression in T helper cell subsets. *Science* 1995;270:1215–8.
- [55] Yan H, Krishnan K, Lim JT, Contillo LG, Krolewski JJ. Molecular characterization of an alpha interferon receptor 1 subunit (IFNAR1) domain required for TYK2 binding and signal transduction. *Molecular and Cellular Biology* 1996;16:2074–82.
- [56] Domanski P, Fish E, Nadeau OW, Witte M, Platanias LC, Yan H, et al. A region of the beta subunit of the interferon alpha receptor different from box 1 interacts with Jak1 and is sufficient to activate the Jak-Stat pathway and induce an antiviral state. *The Journal of Biological Chemistry* 1997;272:26388–93.
- [57] Wilks AF. Two putative protein-tyrosine kinases identified by application of the polymerase chain reaction. *Proceedings of the National Academy of Sciences of the United States of America* 1989;86:1603–7.
- [58] Wilks AF, Harpur AG, Kurban RR, Ralph SJ, Zurcher G, Ziemiński A. Two novel protein-tyrosine kinases, each with a second phosphotransferase-related catalytic domain, define a new class of protein kinase. *Molecular and Cellular Biology* 1991;11:2057–65.
- [59] Stark GR, Darnell Jr JE. The JAK-STAT pathway at twenty. *Immunity* 2012; 36:503–14.
- [60] O'Shea JJ, Plenge R. JAK and STAT signaling molecules in immunoregulation and immune-mediated disease. *Immunity* 2012;36:542–50.
- [61] Ragimbeau J, Dondi E, Alcover A, Eid P, Uze G, Pellegrini S. The tyrosine kinase Tyk2 controls IFNAR1 cell surface expression. *The EMBO Journal* 2003; 22:537–47.
- [62] Durbin JE, Fernandez-Sesma A, Lee CK, Rao TD, Frey AB, Moran TM, et al. Type I, IFN modulates innate and specific antiviral immunity. *Journal of Immunology* 2000;164:4220–8.
- [63] Li X, Leung S, Kerr IM, Stark GR. Functional subdomains of STAT2 required for preassociation with the alpha interferon receptor and for signaling. *Molecular and Cellular Biology* 1997;17:2048–56.
- [64] Yang CH, Shi W, Basu L, Murti A, Constantinescu SN, Blatt L, et al. Direct association of STAT3 with the IFNAR-1 chain of the human type I interferon receptor. *The Journal of Biological Chemistry* 1996;271:8057–61.
- [65] Zhao W, Lee C, Piganis R, Plumlee C, de Weerd N, Hertzog PJ, et al. A conserved IFN- α receptor tyrosine motif directs the biological response to type I IFNs. *Journal of Immunology* 2008;180:5483–9.
- [66] Takaoka A, Mitani Y, Suemori H, Sato M, Yokochi T, Noguchi S, et al. Cross talk between interferon- γ and - α /beta signaling components in caveolar membrane domains. *Science* 2000;288:2357–60.
- [67] Mitani Y, Takaoka A, Kim SH, Kato Y, Yokochi T, Tanaka N, et al. Cross talk of the interferon- α /beta signalling complex with gp130 for effective interleukin-6 signalling. *Genes to Cells Devoted to Molecular & Cellular Mechanisms* 2001;6:631–40.
- [68] Platanias LC. Mechanisms of type-I- and type-II-interferon-mediated signalling. *Nature Reviews Immunology* 2005;5:375–86.
- [69] van Boxel-Dezaire AH, Rani MR, Stark GR. Complex modulation of cell type-specific signaling in response to type I interferons. *Immunity* 2006;25:361–72.
- [70] Ramana CV, Gil MP, Han Y, Ransohoff RM, Schreiber RD, Stark GR. Stat1-independent regulation of gene expression in response to IFN- γ . *Proceedings of the National Academy of Sciences of the United States of America* 2001;98:6674–9.
- [71] Yang CH, Murti A, Pfeffer SR, Kim JG, Donner DB, Pfeffer LM. Interferon α /beta promotes cell survival by activating nuclear factor kappa B through phosphatidylinositol 3-kinase and Akt. *The Journal of Biological Chemistry* 2001;276:13756–61.
- [72] Goh KC, Haque SJ, Williams BR. p38 MAP kinase is required for STAT1 serine phosphorylation and transcriptional activation induced by interferons. *The EMBO Journal* 1999;18:5601–8.
- [73] Joshi S, Kaur S, Redig AJ, Goldsborough K, David K, Ueda T, et al. Type I interferon (IFN)-dependent activation of Mnk1 and its role in the generation of growth inhibitory responses. *Proceedings of the National Academy of Sciences of the United States of America* 2009;106:12097–102.
- [74] Kaur S, Sassano A, Majchrzak-Kita B, Baker DP, Su B, Fish EN, et al. Regulatory effects of mTORC2 complexes in type I IFN signaling and in the generation of IFN responses. *Proceedings of the National Academy of Sciences of the United States of America* 2012;109:7723–8.
- [75] Kaur S, Sassano A, Dolniak B, Joshi S, Majchrzak-Kita B, Baker DP, et al. Role of the Akt pathway in mRNA translation of interferon-stimulated genes. *Proceedings of the National Academy of Sciences of the United States of America* 2008;105:4808–13.
- [76] Petricoin 3rd EF, Ito S, Williams BL, Audet S, Stancato LF, Gamero A, et al. Antiproliferative action of interferon- α requires components of T-cell-receptor signalling. *Nature* 1997;390:629–32.
- [77] Ozato K. PLZF outreach: a finger in interferon's pie. *Immunity* 2009;30:757–8.
- [78] Sadler AJ, Williams BRG. Interferon-inducible antiviral effectors. *Nature Reviews Immunology* 2008;8:559–68.
- [79] Xu D, Holko M, Sadler AJ, Scott B, Higashiyama S, Berkofsky-Fessler W, et al. Promyelocytic leukemia zinc finger protein regulates interferon-mediated innate immunity. *Immunity* 2009;30:802–16.
- [80] Pioli PD, Saleh AM, El Fiky A, Nastiuk KL, Krolewski JJ. Sequential proteolytic processing of an interferon- α receptor subunit by TNF- α converting enzyme and presenilins. *Journal of Interferon & Cytokine Research The Official Journal of the International Society for Interferon and Cytokine Research* 2012;32:312–25.
- [81] Hardy MP, Owczarek CM, Trajanovska S, Liu X, Kola I, Hertzog PJ. The soluble murine type I interferon receptor *Ifnar-2* is present in serum, is independently regulated, and has both agonistic and antagonistic properties. *Blood* 2001;97:473–82.
- [82] Kumar KG, Krolewski JJ, Fuchs SY. Phosphorylation and specific ubiquitin acceptor sites are required for ubiquitination and degradation of the IFNAR1 subunit of type I interferon receptor. *The Journal of Biological Chemistry* 2004;279:46614–20.
- [83] Malakhova OA, Kim KI, Luo JK, Zou W, Kumar KG, Fuchs SY, et al. UBP43 is a novel regulator of interferon signaling independent of its ISG15 isopeptidase activity. *The EMBO Journal* 2006;25:2358–67.

- [84] Min W, Pober JS, Johnson DR. Interferon induction of TAP1: the phosphatase SHP-1 regulates crossover between the IFN- α /beta and the IFN- γ signal-transduction pathways. *Circulation Research* 1998;83:815–23.
- [85] Coccia EM, Uze G, Pellegrini S. Negative regulation of type I interferon signaling: facts and mechanisms. *Cellular and Molecular Biology (Noisy-le-Grand France)* 2006;52:77–87.
- [86] Gibbs VC, Takahashi M, Aguet M, Chuntharapai A. A negative regulatory region in the intracellular domain of the human interferon- α receptor. *The Journal of Biological Chemistry* 1996;271:28710–16.
- [87] Baychelier F, Nardeux PC, Cajean-Feroldi C, Ermonval M, Guymarho J, Tovey MG, et al. Involvement of the Gab2 scaffolding adapter in type I interferon signalling. *Cellular Signalling* 2007;19:2080–7.
- [88] Shuai K, Liu B. Regulation of gene-activation pathways by PIAS proteins in the immune system. *Nature Reviews Immunology* 2005;5:593–605.
- [89] Ezell SA, Polyarchou C, Hatziaepoulou M, Guo A, Sanidas I, Bihani T, et al. The protein kinase Akt1 regulates the interferon response through phosphorylation of the transcriptional repressor EMSY. *Proceedings of the National Academy of Sciences of the United States of America* 2012;109:E613–21.
- [90] Tisserand J, Khetchoumian K, Thibault C, Dembele D, Chambon P, Lossos R. Tripartite motif 24 (Trim24/Tif1 α) tumor suppressor protein is a novel negative regulator of interferon (IFN)/signal transducers and activators of transcription (STAT) signaling pathway acting through retinoic acid receptor α (RAR α) inhibition. *The Journal of Biological Chemistry* 2011;286:33369–79.
- [91] Yoshida K, Yamamoto K, Kohno T, Hironaka N, Yasui K, Kojima C, et al. Active repression of IFN regulatory factor-1-mediated transactivation by IFN regulatory factor-4. *International Immunology* 2005;17:1463–71.
- [92] Sharma B, Joshi S, Sassano A, Majchrzak B, Kaur S, Aggarwal P, et al. Sprouty proteins are negative regulators of interferon (IFN) signaling and IFN-inducible biological responses. *The Journal of Biological Chemistry* 2012;287:42352–60.
- [93] Winter J, Jung S, Keller S, Gregory RI, Diederichs S. Many roads to maturity: microRNA biogenesis pathways and their regulation. *Nature Cell Biology* 2009;11:228–34.
- [94] Kozomara A, Griffiths-Jones S. miRBase: integrating microRNA annotation and deep-sequencing data. *Nucleic Acids Research* 2011;39:D152–7.
- [95] Pillai RS, Bhattacharyya SN, Artus CG, Zoller T, Cougot N, Basyuk E, et al. Inhibition of translational initiation by Let-7 MicroRNA in human cells. *Science* 2005;309:1573–6.
- [96] Gantier MP, Sadler AJ, Williams BR. Fine-tuning of the innate immune response by microRNAs. *Immunology and Cell Biology* 2007;85:458–62.
- [97] Zhang B, Pan X, Cobb GP, Anderson TA. microRNAs as oncogenes and tumor suppressors. *Developmental Biology* 2007;302:1–12.
- [98] Pedersen IM, Cheng G, Wieland S, Volinia S, Croce CM, Chisari FV, et al. Interferon modulation of cellular microRNAs as an antiviral mechanism. *Nature* 2007;449:919–22.
- [99] Bruni R, Marcantonio C, Tritarelli E, Tataseo P, Stellacci E, Costantino A, et al. An integrated approach identifies IFN-regulated microRNAs and targeted mRNAs modulated by different HCV replicon clones. *BMC Genomics* 2011;12:485.
- [100] McCoy CE. The role of miRNAs in cytokine signaling. *Frontiers in Bioscience* 2011;16:2161–71.
- [101] Wang P, Hou J, Lin L, Wang C, Liu X, Li D, et al. Inducible microRNA-155 feedback promotes type I IFN signaling in antiviral innate immunity by targeting suppressor of cytokine signaling 1. *Journal of Immunology* 2010;185:6226–33.
- [102] Papadopoulos AS, Dooley J, Linterman MA, Pierson W, Ucar O, Kyewski B, et al. The thymic epithelial microRNA network elevates the threshold for infection-associated thymic involution via miR-29a mediated suppression of the IFN- α receptor. *Nature Immunology* 2012;13:181–7.
- [103] Der SD, Zhou A, Williams BR, Silverman RH. Identification of genes differentially regulated by interferon α , β , or γ using oligonucleotide arrays. *Proceedings of the National Academy of Sciences of the United States of America* 1998;95:15623–28.
- [104] de Veer MJ, Holko M, Frevel M, Walker E, Der S, Paranjape JM, et al. Functional classification of interferon-stimulated genes identified using microarrays. *Journal of Leukocyte Biology* 2001;69:912–20.
- [105] Samarajiwa SA, Forster S, Auchettl K, Hertzog PJ. INTERFEROME, the database of interferon regulated genes. *Nucleic Acids Research* 2009;37:D852–7.
- [106] Hertzog P, Forster S, Samarajiwa S. Systems biology of interferon responses. *Journal of Interferon & Cytokine Research The Official Journal of the International Society for Interferon and Cytokine Research* 2011;31:5–11.
- [107] Rusinova I, Forster S, Yu S, Kannan A, Masse M, Cumming H, et al. Interferome v2.0: an updated database of annotated interferon-regulated genes. *Nucleic Acids Research* 2013;41:D1040–46.
- [108] Harman AN, Lai J, Turville S, Samarajiwa S, Gray L, Marsden V, et al. HIV infection of dendritic cells subverts the IFN induction pathway via IRF-1 and inhibits type 1 IFN production. *Blood* 2011;118:298–308.
- [109] Zhao M, Zhang J, Phatnani H, Scheu S, Maniatis T. Stochastic expression of the interferon- β gene. *PLoS Biology* 2012;10:e1001249.
- [110] Thomas KE, Galligan CL, Newman RD, Fish EN, Vogel SN. Contribution of interferon- β to the murine macrophage response to the toll-like receptor 4 agonist, lipopolysaccharide. *The Journal of Biological Chemistry* 2006;281:31119–30.
- [111] Carpick BW, Graziano V, Schneider D, Maitra RK, Lee X, Williams BRG. Characterization of the solution complex between the interferon-induced, double-stranded RNA-activated protein kinase and HIV-1 trans-activating region RNA. *Journal of Biological Chemistry* 1997;272:9510–6.
- [112] Ito T, Yang M, May WS. RAX, a cellular activator for double-stranded RNA-dependent protein kinase during stress signaling. *Journal of Biological Chemistry* 1999;274:15427–32.
- [113] Patel CV, Handy I, Goldsmith T, Patel RC. PACT, a stress-modulated cellular activator of interferon-induced double-stranded RNA-activated protein kinase, PKR. *Journal of Biological Chemistry* 2000;275:37993–98.
- [114] Patel RC, Sen GC. PACT, a protein activator of the interferon-induced protein kinase, PKR. *The EMBO Journal* 1998;17:4379–90.
- [115] Goh KC, deVeer MJ, Williams BR. The protein kinase PKR is required for p38 MAPK activation and the innate immune response to bacterial endotoxin. *The EMBO Journal* 2000;19:4292–7.
- [116] Kumar A, Yang YL, Flati V, Der S, Kadereit S, Deb A, et al. Deficient cytokine signaling in mouse embryo fibroblasts with a targeted deletion in the PKR gene: role of IRF-1 and NF- κ B. *The EMBO Journal* 1997;16:406–16.
- [117] Bonnet MC, Weil R, Dam E, Hovanessian AG, Meurs EF. PKR stimulates NF- κ B irrespective of its kinase function by interacting with the I κ B kinase complex. *Molecular and Cellular Biology* 2000;20:4532–42.
- [118] Jiang Z, Zamanian-Daryoush M, Nie H, Silva AM, Williams BR, Li X. Poly(I:C)-induced Toll-like receptor 3 (TLR3)-mediated activation of NF κ B and MAP kinase is through an interleukin-1 receptor-associated kinase (IRAK)-independent pathway employing the signaling components TLR3-TRAF6-TAK1-TAB2-PKR. *The Journal of Biological Chemistry* 2003;278:16713–19.
- [119] Irving AT, Wang D, Vasilevski O, Latchoumanin O, Kozar N, Clayton AH, et al. Regulation of actin dynamics by protein kinase R control of gelsolin enforces basal innate immune defense. *Immunity* 2012;36:795–806.



Prof. Paul J. Hertzog is Director of the Centre for Innate Immunity and Infectious Diseases at Monash Institute of Medical Research. His research interests are in understanding the molecular control of signalling in the innate immune response. His focus is in interferon signalling where current projects address structure-function of receptors, negative regulation of signalling, characterisation of interferon ϵ function in reproductive tract immunity and disease. He has a particular interest in signal transduction and gene regulation in innate immunity and is using systems biology to understand the control of immune response in infectious and inflammatory disease and cancer. He is supported by a National Health and Medical Research Council Senior Principal Research Fellowship and the Victorian Government's Operational Infrastructure Support Program.



Prof. Bryan R.G. Williams is the Director of the Monash Institute of Medical Research, Melbourne, Australia. Following postdoctoral training at the National Institute for Medical Research, Mill Hill, London, he held faculty positions at the University of Toronto and the Hospital for Sick Children, Toronto, Canada. He was Chairman of the Department of Cancer Biology at the Lerner Research Institute, Cleveland Clinic Foundation, USA. He has a distinguished research career spanning more than 30 years in the areas of innate immunity and cancer biology, and has published over 300 peer-reviewed journal articles and reviews. He is internationally recognised for his work in innate immunity, and for his studies on protein kinase R, an important innate immune signalling molecule. He has also gained prominence in the field of RNA interference. Professor Williams received the 1990 Milstein Award from the International Society for Interferon Research, in recognition of his contributions to advancing interferon research for the treatment of human diseases. He is a Fellow of the American Academy of Microbiology and Fellow of the Australian Academy of Science. He is supported by the Australian Research Council, the Australian National Health and Medical Research Council and the Victorian Government's Operational Infrastructure Support Program.