Table 5 Health state utility values included in HTA reports

Region	HTA report (year)	Type of economic evaluation	Patient population	Treatment	Source of utility values	Utility values reported
Australia	PBAC (2018) [73]	CUA	R/R PMBCL	Pembrolizumab Standard of care chemotherapies including ICE, DHAP, or GDP	Keynote—170 trial	Progression-free state: 0.80 Progressed disease: 0.62
Australia	PBAC (2019) [74]	CUA	Patients with R/R DLBCL who have failed prior therapies and are either ineligible for SCT (due to comorbidities, age, or a failed salvage regimen), or who have had disease relapse after HSCT	Pola-BR BR	JULIET trial/tisagenlecleucel NICE report[21]	Progression-free off treatment: 0.83 Progressive disease: 0.71
Canada	CADTH (2019) [75]	CUA	Adult patients with R/R LBCL after ≥ 2 lines of systemic therapy including DLBCL NOS, HGBCL, and DLBCL arising from FL	Tisagenlecleucel	JULIET trial (8 March 2017 data cutoff)	Disutility values: Short-term AEs associated with treatments (except for CRS): − 0.15 CRS-related or non–CRS-related ICU stay: − 0.83
				Salvage chemotherapy (rituximab plus GDP)	SCHOLAR-1 [76]	Disutility values: Short-term AEs associated with treatments: − 0.15
England and Wales	NICE (2020) [77]	CUA	Patients with R/R DLBCL who are ineligible for HSCT	Pola-BR BR	NICE (2019)	Progression-free survival: 0.72 Progressive disease: 0.65
England and Wales	NICE (2019) [78]	CUA	Adult patients with R/R DLBCL after ≥ 2 lines of systemic therapy	Tisagenlecleucel Salvage chemotherapy (GDP)	JULIET trial	Progression-free state: 0.83 Progressed disease: 0.71 AE disutility: 0.15
England and Wales	NICE (2019) [79]	CUA	Adult patients with R/R DLBCL and PMBCL after ≥ 2 lines of systemic therapy	Axi-cel Salvage chemotherapy excluding pixantrone	ZUMA-1 trial	By follow-up: Screening: 0.739 Week 4: 0.675 Month 3: 0.756 Month 6: 0.758 By response: Complete response: 0.743 Partial response: 0.788 Stable disease: 0.636 Progression-free disease: 0.722 Progressed disease: 0.647
England and Wales	NICE (2014) [80]	CUA	Adults with aggressive de novo or transformed NHL relapsed after ≥ 2 chemotherapy regimens, per ITT population of PIX301	Pixantrone Physician's choice	Published literature for patients receiving second- and subsequent-line treatment for renal cell carcinoma	Manufacturer's submission: Preprogression: 0.81 Postprogression: 0.60 Revised model: Preprogression: 0.76 Postprogression: 0.68

AE adverse event; *axi-cel* axicabtagene ciloleucel; *BR* bendamustine and rituximab; *CADTH* Canadian Agency for Drugs and Technologies in Health; *CRS* cytokine release syndrome; *CUA* cost-utility analysis; *DHAP* dexamethasone, cytarabine, and cisplatin; *DLBCL* diffuse large B cell lymphoma; *FL* follicular lymphoma; *GDP* gemcitabine, dexamethasone, and cisplatin; *HGBCL* high-grade B cell lymphoma; *HSCT* hematopoietic stem cell transplantation; *HTA* health technology assessment; *ICE* ifosfamide, carboplatin, and etoposide; *ICU* intensive care unit; *ITT* intent to treat; *LBCL* large B cell lymphoma; *NHL* non-Hodgkin lymphoma; *NICE* National Institute for Health and Care Excellence; *NOS* not otherwise specified; *PBAC* Pharmaceutical Benefits Advisory Committee; *PMBCL* primary mediastinal large B cell lymphoma; *pola-BR* polatuzumab vedotin, bendamustine, and rituximab; *R/R* relapsed or refractory