

Macromolecules are polymers, built from monomers

Large carbohydrates, proteins, and nucleic acids, also known as **macromolecules** for their huge size, are chain-like molecules called polymers (from the Greek *polys*, many, and *meros*, part). A **polymer** is a long molecule consisting of many similar or identical building blocks linked by covalent bonds, much as a train consists of a chain of boxcars. The repeating units that serve as the building blocks of a polymer are smaller molecules called **monomers** (from the Greek *monos*, single). In addition to forming polymers, some monomers have functions of their own.

The Synthesis and Breakdown of Polymers


Although each class of polymer is made up of a different type of monomer, the chemical mechanisms by which cells make polymers (polymerization) and break them down are similar for all classes of large biological molecules. In cells, these processes are facilitated by **enzymes**, specialized macromolecules (usually proteins) that speed up chemical reactions. The reaction that connects a monomer to another monomer or a polymer is a **condensation reaction**, a reaction in which two molecules are covalently bonded to each other with the loss of a small molecule. If a water molecule is lost, it is known as a **dehydration reaction**. For example, carbohydrate and protein polymers are synthesized by dehydration reactions. Each reactant contributes part of the water molecule that is released during the reaction: One provides a hydroxyl group (—OH), while the other provides a hydrogen (—H) (**Figure 5.2a**). This reaction is repeated as monomers are added to the chain one by one, lengthening the polymer.

Polymers are disassembled to monomers by **hydrolysis**, a process that is essentially the reverse of the dehydration reaction (**Figure 5.2b**). Hydrolysis means water breakage (from the Greek *hydro*, water, and *lysis*, break). The bond between monomers is broken by the addition of a water molecule, with a hydrogen from water attaching to one monomer and the hydroxyl group attaching to the other. An example of hydrolysis within our bodies is the process of digestion. The bulk of the organic material in our food is in the form of polymers that are much too large to enter our cells. Within the digestive tract, various enzymes attack the polymers, speeding up hydrolysis. Released monomers are then absorbed into the bloodstream for distribution to all body cells. Those cells can then use dehydration reactions to assemble the monomers into new, different polymers that can perform specific functions required by the cell. (Dehydration reactions and hydrolysis can also be involved in the formation and breakdown of molecules that are not polymers, such as some lipids.)

The Diversity of Polymers

A cell has thousands of different macromolecules; the collection varies from one type of cell to another. The inherited

▼ **Figure 5.2** The synthesis and breakdown of carbohydrate and protein polymers.



➔ Mastering Biology Animation: Making and Breaking Polymers

differences between close relatives, such as human siblings, reflect small variations in polymers, particularly DNA and proteins. Molecular differences between unrelated individuals are more extensive, and those between species greater still. The diversity of macromolecules in the living world is vast, and the possible variety is effectively limitless.

What is the basis for such diversity in life's polymers? These molecules are constructed from only 40 to 50 common monomers and some others that occur rarely. Building a huge variety of polymers from such a limited number of monomers is analogous to constructing hundreds of thousands of words from only 26 letters of the alphabet. The key is arrangement—the particular linear sequence that the units follow. However, this analogy falls far short of describing the great diversity of macromolecules because most biological polymers have many more monomers than the number of letters in even the longest word. Proteins, for example, are built from 20 kinds of amino acids arranged in chains that are typically hundreds of amino acids long. The molecular logic of life is simple but elegant: Small molecules common to all organisms act as building blocks that are ordered into unique macromolecules.

Despite this immense diversity, molecular structure and function can still be grouped roughly by class. Let's examine each of the four major classes of large biological molecules. For each class, the large molecules have emergent properties not found in their individual components.