

List homomorphisms

Troels Henriksen (athas@sigkill.dk)

DIKU
University of Copenhagen

29-11-2021

Preliminaries

Material

- *The Third Homomorphism Theorem* (Gibbons; 1995)
- *Construction of List Homomorphisms by Tupling and Fusion* (Hu, Iwasaki, Takeichi; 1996)

Approach

- **This material is dense.**
- Do not get dissuaded!
 - ▶ Ask about notation that is unclear.
- Proofs in the papers, not the lecture.
- We focus on intuition about working algebraically with (parallel!) programs.
- My notation is slightly different than in the papers.
 - ▶ I write *hom* for homomorphisms (and leave out neutral elements).
 - ▶ I write *map f* instead of f^* .

List homomorphisms

- Definitions and properties

- Homomorphism theorems

- Applying the third homomorphism theorem

Deriving efficient almost homomorphisms

- Almost homomorphisms

- Deriving initial almost homomorphisms by tupling

- Fusion theorems for homomorphisms

- Fusion of *mss*

Conclusions

List homomorphisms

Definition: list homomorphism

A function h is called a list homomorphism if there exists a binary operator \odot such that for all finite lists x, y

$$h (x \mathbin{++} y) = h x \odot h y$$

where $\mathbin{++}$ means list concatenation.

- Intuitively, homomorphisms are functions where we can split the input arbitrarily, solve the subproblems, and combine the partial results.
- Each “split” can be solved in parallel.

Examples of list homomorphisms

Summation

$$\textit{sum} [] = 0$$

$$\textit{sum} [x] = x$$

$$\textit{sum} (x \mathbin{++} y) = \textit{sum} x + \textit{sum} y$$

Length

$$\textit{len} [] = 0$$

$$\textit{len} [x] = 1$$

$$\textit{len} (x \mathbin{++} y) = \textit{len} x + \textit{len} y$$

Observations

$$h (x \mathbin{+} y) = h x \odot h y$$

- \odot is necessarily associative because $\mathbin{+}$ is associative.
- $h []$ is necessarily a neutral element for \odot .

For concision, we write

$$\text{hom } (\odot) f$$

for a list homomorphism h , where

$$f x = h [x]$$

$$0_{\odot} = h []$$

Examples

$$\begin{aligned} \text{sum} &= \text{hom } (+) \text{ id} \\ \text{length} &= \text{hom } (+) (\text{const } 1) \end{aligned}$$

Reductions

Reductions are homomorphisms

A function of the form

$$\text{hom } (\odot) \text{ id}$$

is called a *reduction* with operator \odot .

- Equivalent to what we would write as `reduce \odot 0 \odot` in Futhark.
- For concision we will abbreviate as follows:

$$\text{red } \odot = \text{hom } (\odot) \text{ id}$$

- Given \odot , the neutral element 0_{\odot} is always unique (think about why).

Maps are homomorphisms

A function of the form

$$\text{hom } (+) ([\cdot] \circ f)$$

is called a *map* with function f .

- $[\cdot]$ is the function $\lambda x \rightarrow [x]$ that turns x into $[x]$.
- We'll write this as *map* f .
- Really is just what you expect, no surprises here.

Leftwards and rightwards functions

A function on lists h is \oplus -leftwards if and only if for all elements a and lists y ,

$$h ([a] \mathbin{\dot{+}} y) = a \oplus h y$$

- \oplus need not be associative.
- **Intuitively, what does this say about h ?**

Leftwards and rightwards functions

A function on lists h is \oplus -leftwards if and only if for all elements a and lists y ,

$$h ([a] \mathbin{++} y) = a \oplus h y$$

- \oplus need not be associative.
- **Intuitively, what does this say about h ?**
- Suppose

$$h [] = e$$

then

$$h [a, b, c] = a \oplus (b \oplus (c \oplus e))$$

- A \oplus -leftwards function is a right-to-left *fold*, denoted by *foldr* $(\oplus) e$.
- Symmetrically, for an \otimes -rightwards function h ,

$$h (y \mathbin{++} [a]) = h y \otimes a$$

which we denote by *foldl* $(\otimes) e$.

List homomorphisms

Definitions and properties

Homomorphism theorems

Applying the third homomorphism theorem

Deriving efficient almost homomorphisms

Almost homomorphisms

Deriving initial almost homomorphisms by tupling

Fusion theorems for homomorphisms

Fusion of *mss*

Conclusions

First Homomorphism Theorem

Every monomorphism can be written as the composition of a reduction and a map.

$$\text{hom } (\odot) f = \text{hom } (\odot) \text{id} \circ \text{map } f$$

Conversely, every such composition is a homomorphism.

- **This is why *map – reduce* compositions are so crucial.**
- Sometimes it is easiest to invent parallel algorithms by thinking “homomorphism-style” in terms of recursively splitting and combining subresults.
 - ▶ Can turn this into data-parallel map-reduce code mechanically.

Second Homomorphism Theorem

If \odot is associative, then

$$\begin{aligned} \text{hom } (\odot) f &= \text{foldr } (\oplus) 0_{\odot} \quad \text{where } x \oplus y = f \ x \odot y \\ &= \text{foldl } (\otimes) 0_{\otimes} \quad \text{where } x \otimes y = x \odot f \ y \end{aligned}$$

That is, *every homomorphism is both a leftwards and a rightwards function.*

- **Intuition:** as a homomorphism allows us to split “however we want”, we can also split consistently with a left (or right) bias.
- **What about the other direction?**

Third Homomorphism Theorem

If h is leftwards and rightwards, then h is a homomorphism.

- **Why this is useful:** If we can come up with left-folds and right-folds that both solve the problem, then a homomorphism must also exist.
- **Put another way:** If a problem can be solved with both a *foldr* and a *foldl*, then a parallel algorithm for the problem also exists.

Third Homomorphism Theorem

If h is leftwards and rightwards, then h is a homomorphism.

- **Why this is useful:** If we can come up with left-folds and right-folds that both solve the problem, then a homomorphism must also exist.
- **Put another way:** If a problem can be solved with both a *foldr* and a *foldl*, then a parallel algorithm for the problem also exists.
- **Unfortunately:**
 - ▶ Not all homomorphisms are usefully parallel.
 - ▶ The proof does not tell us how to *construct* a practical homomorphism from the left and right folds.

Proof sketch (full details in Gibbons' paper)

The statement to prove

If h is leftwards and rightwards, then h is a homomorphism.

Steps:

1. Prove that

$$h\ v = h\ x \wedge h\ w = h\ y \Rightarrow h\ (v \uplus w) = h\ (x \uplus y)$$

► Rough steps:

$$\begin{aligned} h\ (v \uplus w) &= \dots \\ &= h\ (v \uplus y) && \text{(By treating } h \text{ as leftwards function.)} \\ &= \dots \\ &= h\ (x \uplus y) && \text{(By treating } h \text{ as rightwards function.)} \end{aligned}$$

2. Prove that this means h is a homomorphism.

The function h is a homomorphism if and only if the implication

$$h\ v = h\ x \wedge h\ w = h\ y \Rightarrow h\ (v \# w) = h\ (x \# y)$$

holds for all lists v, w, x, y .

“Only if” part

If h is a homomorphism then for all a, b there is a \oplus such that,

$$h\ (a \# b) = h\ a \oplus h\ b$$

and then

$$h\ (v \# w) = h\ v \oplus h\ w \quad (\text{Def. of homomorphism})$$

$$= h\ x \oplus h\ y \quad (\text{LHS of implication})$$

$$= h\ (x \# y) \quad (\text{Def. of homomorphism})$$

The function h is a homomorphism if and only if the implication

$$h\ v = h\ x \wedge h\ w = h\ y \Rightarrow h\ (v \uplus w) = h\ (x \uplus y)$$

holds for all lists v, w, x, y .

“If” part

- Choose g such that $h \circ g \circ h = h$ and define

$$a \odot b = h\ (g\ a \uplus g\ b)$$

- (We'll show that g exists in a moment.)
- We have $h\ x = h\ (g\ (h\ x))$ and $h\ y = h\ (g\ (h\ y))$.
- With $v = g\ (h\ x)$ and $w = g\ (h\ y)$ then

$$\begin{aligned} h\ (x \uplus y) &= h\ (g\ (h\ x)) \uplus g\ (h\ y) \\ &= h\ x \odot h\ y \end{aligned}$$

Existence of g

Given h , we must construct g such that

$$h \circ g \circ h = h$$

- Can we be sure such a g exists for any h ?
 - ▶ **Yes:** To compute $g x$, enumerate domain of h and produce first y such that $h y = x$.
 - ▶ **Theoretically sound, but not really practical.**
 - ▶ When we apply the Third Homomorphism Theorem we better come up with a more reasonable g than this.

List homomorphisms

Definitions and properties

Homomorphism theorems

Applying the third homomorphism theorem

Deriving efficient almost homomorphisms

Almost homomorphisms

Deriving initial almost homomorphisms by tupling

Fusion theorems for homomorphisms

Fusion of *mss*

Conclusions

Consider sorting

We can define

$$\text{sort} = \text{foldr ins } []$$

where

$$\begin{aligned} \text{ins } a [] &= [a] \\ \text{ins } a (b : x) &= \begin{cases} a : b : x & \text{if } a \leq b \\ b : \text{ins } a x & \text{otherwise} \end{cases} \end{aligned}$$

That is, $\text{ins } a l$ inserts a at the appropriate location in the sorted list l .

Consider sorting

We can define

$$\textit{sort} = \textit{foldr ins []}$$

where

$$\begin{aligned}\textit{ins } a [] &= [a] \\ \textit{ins } a (b : x) &= \begin{cases} a : b : x & \text{if } a \leq b \\ b : \textit{ins } a x & \text{otherwise} \end{cases}\end{aligned}$$

That is, $\textit{ins } a l$ inserts a at the appropriate location in the sorted list l .

$$\textit{sort} = \textit{foldl ins' []}$$

where

$$\textit{ins'} x a = \textit{ins } a x$$

As \textit{sort} is both leftwards and rightwards, there is a homomorphic sorting algorithm.

Homomorphic sorting

By the Third Homomorphism Theorem, we get a homomorphism $hom \odot [\cdot]$ where

$$u \odot v = sort (unsort\ u \uplus unsort\ v)$$

for some function *unsort* such that

$$sort \circ unsort \circ sort = sort$$

Can you think of such a function?

Homomorphic sorting

By the Third Homomorphism Theorem, we get a homomorphism $hom \odot [\cdot]$ where

$$u \odot v = sort (unsort\ u \uplus unsort\ v)$$

for some function *unsort* such that

$$sort \circ unsort \circ sort = sort$$

Can you think of such a function?

$$unsort = id$$

This gives

$$u \odot v = sort (u \uplus v)$$

Is this a good sorting algorithm?

Homomorphic sorting

By the Third Homomorphism Theorem, we get a homomorphism $hom \odot [\cdot]$ where

$$u \odot v = sort (unsort\ u \uplus unsort\ v)$$

for some function *unsort* such that

$$sort \circ unsort \circ sort = sort$$

Can you think of such a function?

$$unsort = id$$

This gives

$$u \odot v = sort (u \uplus v)$$

Is this a good sorting algorithm?

Homomorphic sorting in words

“To sort the concatenation of two sorted lists u and v , concatenate u and v and then sort the result using some other sorting algorithm.”

Improving the homomorphic sort

- In the paper, Gibbons goes on to mechanically derive a more efficient \odot , based on the property that u will be sorted.

$$\begin{aligned}u \odot v &= \text{sort } (u \uplus v) \\&= \text{foldl } \text{ins}' \ [] \ (u \uplus v) \\&= \text{foldl } \text{ins}' \ (\text{foldl } \text{ins}' \ [] \ u) v \\&= \text{foldl } \text{ins}' \ (\text{sort } u) v \\&= \text{foldl } \text{ins}' \ u \ v\end{aligned}$$

- Now $u \odot v$ merges two sorted lists in $O(n)$ time.
- By also assuming v is sorted, Gibbons eventually derives mergesort.
 - ▶ Read the paper!
- **Not parallel**, but we could in principle have derived a parallel \odot instead.

List homomorphisms

Definitions and properties

Homomorphism theorems

Applying the third homomorphism theorem

Deriving efficient almost homomorphisms

Almost homomorphisms

Deriving initial almost homomorphisms by tupling

Fusion theorems for homomorphisms

Fusion of *mss*

Conclusions

Almost homomorphisms

Some functions are not homomorphisms.

- Knowing the means of arrays x and y does not let us compute the mean of $x \uplus y$.

But they can be turned into homomorphisms if we compute a little extra information.

- Knowing the means x_{mean}, y_{mean} and sizes x_n, y_n of x, y lets us compute the mean and size of $x \uplus y$.

This is called an *almost homomorphism*.

mean as an almost homomorphism

$$\begin{aligned} \text{mean } [] &= (0, 0) \\ \text{mean } [x] &= (x, 1) \\ \text{mean } (x \mathbin{++} y) &= \left(\frac{x_{\text{mean}} \cdot x_n + y_{\text{mean}} \cdot y_n}{x_n + y_n}, x_n + y_n \right) \end{aligned}$$

where

$$(x_{\text{mean}}, x_n) = \text{mean } x$$

$$(y_{\text{mean}}, y_n) = \text{mean } y$$

To compute the mean we then project out a component

$$\pi_1 \circ \text{mean}$$

List homomorphisms

Definitions and properties

Homomorphism theorems

Applying the third homomorphism theorem

Deriving efficient almost homomorphisms

Almost homomorphisms

Deriving initial almost homomorphisms by tupling

Fusion theorems for homomorphisms

Fusion of *mss*

Conclusions

The maximum segment sum problem

Problem statement

Given a list A , find the largest sum of a subsequence $A[i : j]$.

Examples

$$mss [0, 3, -2] = 3$$

$$mss [4, -1, 0] = 4$$

$$mss [0, 3, -2, 4, -1, 0] = 5$$

Not a homomorphism

- Knowing $mss\ x$ and $mss\ y$ is not enough to give us $mss(x \# y)$.
- E.g. $x = [0, 3, -2], y = [4, -1, 0]$ as above.

But is an almost homomorphism

- With human creativity we can come up with an associative operator on 4 – *tuples* that lets us solve MSS as an almost homomorphism.
- But can we also derive it systematically?

Specification of mss

$$mss = max^s \circ (map^s \text{ sum}) \circ segs$$

Note that max^s and map^s are set operations.

$$mss = \text{max}^s \circ (\text{map}^s \text{ sum}) \circ \text{segs}$$

$$\text{segs } [] = \{\}$$

$$\text{segs } [x] = \{[x]\}$$

$$\text{segs } (xs \mathrel{++} ys) = \text{segs } xs \cup \text{segs } ys \cup (\text{tails } xs \mathcal{X}_{++} \text{inits } ys)$$

$$\text{inits } [] = []$$

$$\text{inits } [x] = [[x]]$$

$$\text{inits } (xs \mathrel{++} ys) = \text{inits } xs \mathrel{++} \text{map } (xs \mathrel{++}) (\text{inits } ys)$$

$$\text{tails } [] = []$$

$$\text{tails } [x] = [[x]]$$

$$\text{tails } (xs \mathrel{++} ys) = \text{map } (+ys) (\text{tails } xs) \mathrel{++} \text{tails } ys$$

$$[x_1, \dots, x_n] \mathcal{X}_{\oplus} [y_1, \dots, y_m] = \{x_1 \oplus y_1, \dots, x_1 \oplus y_m, \dots, x_n \oplus y_1, \dots, x_n \oplus y_m\}$$

$$(f \triangle g) x = (f x, g x)$$

So $f \triangle g$ is a function that applies both f and g to its argument.

$$a(\oplus \triangle \otimes)b = (a \oplus b, a \otimes b)$$

Example

$$(+1) \triangle (\cdot 2) \triangle (-3) = \lambda x \rightarrow (x + 1, x \cdot 2, x - 3)$$

$$(+) \triangle (\cdot) \triangle (-) = \lambda(x, y) \rightarrow (x + y, x \cdot y, x - y)$$

$$(f \triangle g) x = (f x, g x)$$

So $f \triangle g$ is a function that applies both f and g to its argument.

$$a(\oplus \triangle \otimes)b = (a \oplus b, a \otimes b)$$

Example

$$(+1) \triangle (\cdot 2) \triangle (-3) = \lambda x \rightarrow (x + 1, x \cdot 2, x - 3)$$

$$(\+) \triangle (\cdot) \triangle (-) = \lambda(x, y) \rightarrow (x + y, x \cdot y, x - y)$$

$$\Delta_1^n f_i = f_1 \triangle \cdots \triangle f_n$$

So Δ_1^n tuples the n functions f_i :

$$(\Delta_1^n f_i) x = (f_1 x, \cdots, f_n x)$$

Tupling of homomorphisms

Let h_1, \dots, h_n be mutually defined as follows:

$$\begin{aligned}h_j [] &= 0_{\oplus_j} \\h_j [x] &= f_j x \\h_j (xs \uplus ys) &= ((\Delta_1^n h_i) xs) \oplus_j ((\Delta_i^n h_i) ys)\end{aligned}$$

Then

$$\Delta_1^n h_i = \text{hom } (\Delta_1^n \oplus_i) (\Delta_1^n f_i) (0_{\oplus_1}, \dots, 0_{\oplus_n})$$

Implication: tupling mutually recursive functions that *traverse the same list in a specific way* will produce a list homomorphism.

Notation reminder

$$(\Delta_1^n f_i) x = (f_1 x, \dots, f_n x)$$

Writing *segs* in the required form

We have

$$\begin{aligned} \text{segs } [] &= \{\} \\ \text{segs } [x] &= \{[x]\} \\ \text{segs } (xs \mathbin{++} ys) &= \underline{\text{segs } xs} \cup \underline{\text{segs } ys} \cup (\underline{\text{tails } xs} \mathcal{X}_{++} \underline{\text{inits } ys}) \end{aligned}$$

and want n mutually recursive functions

$$\begin{aligned} h_j [] &= 0_{\oplus_j} \\ h_j [x] &= f_j x \\ h_j (xs \mathbin{++} ys) &= ((\Delta_1^n h_i) xs) \oplus_j ((\Delta_i^n h_i) ys) \end{aligned}$$

- *segs* must be tupled with *tails* because *segs* and *tails* traverse the same list *xs*.
- *segs* must be tupled with *inits* because *segs* and *inits* traverse the same list *xs*.

Considering *inits* and *tails*

$$\begin{aligned} \text{inits } [] &= [] \\ \text{inits } [x] &= [[x]] \\ \text{inits } (xs \mathbin{++} ys) &= \underline{\text{inits } xs} \mathbin{++} \text{map } (\underline{xs} \mathbin{++}) (\text{inits } ys) \end{aligned}$$

$$\begin{aligned} \text{tails } [] &= [] \\ \text{tails } [x] &= [[x]] \\ \text{tails } (xs \mathbin{++} ys) &= \text{map } (\mathbin{++} \underline{ys}) (\text{tails } xs) \mathbin{++} \underline{\text{tails } ys} \end{aligned}$$

We need to tuple with *id*:

$$\begin{aligned} \text{id } [] &= [] \\ \text{id } [x] &= [x] \\ \text{id } (xs \mathbin{++} ys) &= \text{id } xs \mathbin{++} \text{id } ys \end{aligned}$$

- To summarise, our tupled function will be

$$\text{segs } \triangle \text{ inits } \triangle \text{ tails } \triangle \text{ id}$$

Tupling $segs \triangle inits \triangle tails \triangle id$

Still need to phrase this in the form

$$\begin{aligned}h_j [] &= 0_{\oplus_j} \\h_j [x] &= f_j x \\h_j (xs \# ys) &= ((\Delta_1^n h_i) xs) \oplus_j ((\Delta_i^n h_i) ys)\end{aligned}$$

i.e. derive f_1, \oplus_1 for $segs$, f_2, \oplus_2 for $inits$, f_3, \oplus_3 for $tails$, f_4, \oplus_4 for id .

- Actually straightforward: pick corresponding recursive calls from the tuples.

Deriving f_1 and \oplus_1

Original

$$\begin{aligned} \text{segs } [] &= \{\} \\ \text{segs } [x] &= \{[x]\} \\ \text{segs } (xs \text{ ++ } ys) &= \underline{\text{segs } xs} \cup \underline{\text{segs } ys} \cup (\underline{\text{tails } xs} \mathcal{X}_{\text{++}} \underline{\text{inits } ys}) \end{aligned}$$

Goal

$$\begin{aligned} h_1 [] &= 0_{\oplus_1} \\ h_1 [x] &= f_1 x \\ h_1 (xs \text{ ++ } ys) &= ((\Delta_1^n h_i) xs) \oplus_1 ((\Delta_i^n h_i) ys) \end{aligned}$$

Derivation

$$\begin{aligned} f_1 x &= \{[x]\} \\ (s_1, i_1, t_1, d_1) \oplus_1 (s_2, i_2, t_2, d_2) &= s_1 \cup s_2 \cup (t_1 \mathcal{X}_{\text{++}} i_2) \end{aligned}$$

All derivations

Skeleton as a reminder

$$\begin{aligned}h_1 [] &= 0_{\oplus_1} \\h_1 [x] &= f_1 x \\h_1 (xs \# ys) &= ((\Delta_1^n h_i) xs) \oplus_1 ((\Delta_i^n h_i) ys)\end{aligned}$$

Derivation

$$\begin{aligned}f_1 x &= \{[x]\} \\(s_1, i_1, t_1, d_1) \oplus_1 (s_2, i_2, t_2, d_2) &= s_1 \cup s_2 \cup (t_1 \mathcal{X}_{\#} i_2) \\f_2 x &= [[x]] \\(s_1, i_1, t_1, d_1) \oplus_2 (s_2, i_2, t_2, d_2) &= i_1 \# \text{map } (d_1 \#) i_2 \\f_3 x &= [[x]] \\(s_1, i_1, t_1, d_1) \oplus_2 (s_2, i_2, t_2, d_2) &= \text{map } (d_2 \#) t_1 \# t_2 \\f_4 x &= [x] \\(s_1, i_1, t_1, d_1) \oplus_2 (s_2, i_2, t_2, d_2) &= d_1 \# d_2\end{aligned}$$

The almost homomorphism

$$\text{segs} = \pi_1 \circ \text{hom } (\Delta_1^n \oplus_i) (\Delta_1^n f_i)$$

List homomorphisms

Definitions and properties

Homomorphism theorems

Applying the third homomorphism theorem

Deriving efficient almost homomorphisms

Almost homomorphisms

Deriving initial almost homomorphisms by tupling

Fusion theorems for homomorphisms

Fusion of *mss*

Conclusions

Fusion

Reminder of where we are

$$mss = max^s \circ (map^s sum) \circ segs$$

where

$$segs = \pi_1 \circ hom (\Delta_1^4 \oplus i) (\Delta_1^4 f_i)$$

Fusion

Merging adjacent operations to avoid intermediate structures or to allow further simplification.

Example of fusion

$$map f \circ map g = map (f \circ g)$$

Can we fuse $segs$ into $(map^s sum)$?

Fusion theorem for homomorphisms

Let h and $\text{hom}(\oplus) f$ be given. If there exists \otimes such that

$$\forall x, y. h(x \oplus y) = h x \otimes h y$$

then

$$h \circ \text{hom}(\oplus) f = \text{hom}(\otimes) (h \circ f)$$

Example of applying fusion theorem

- Suppose $h = (2 \times)$, $(\oplus) = (\uparrow)$, $f = \text{abs}$.

(\uparrow is the “max” operator)

- Must find \otimes such that

$$\forall x, y. 2(x \uparrow y) = 2x \otimes 2y$$

Example of applying fusion theorem

- Suppose $h = (2\times), (\oplus) = (\uparrow), f = \text{abs}$. (\uparrow is the “max” operator)
- Must find \otimes such that

$$\forall x, y. 2(x \uparrow y) = 2x \otimes 2y$$

We fiddle a bit and find

$$x \otimes y = x \uparrow y$$

- Which means we can fuse as

$$\begin{aligned}(2\times) \circ \text{hom } (\uparrow) \text{ abs} &= \text{hom } (\uparrow) ((2\times) \circ \text{abs}) \\ &= \text{red } (\uparrow) \circ \text{map } ((2\times) \circ \text{abs})\end{aligned}$$

Sadly this is not yet enough

Reminder:

$$mss = \max^s \circ (\text{map}^s \text{ sum}) \circ \text{segs}$$

where

$$\text{segs} = \pi_1 \circ \text{hom} (\Delta_1^4 \oplus i) (\Delta_1^4 f_i)$$

- We want to fuse *segs* into $(\text{map}^s \text{ sum})$
- Since *segs* is not a homomorphism but an *almost homomorphism*, we cannot apply the fusion theorem.

Fortunately:

- There is also a fusion theorem for almost homomorphisms.
- Idea is to create a function H that “contains” h and “respects” π_1 .

Fusion theorem for almost homomorphisms

Let h and $hom (\Delta_1^n \oplus_i) (\Delta_1^n f_i)$ be given. If there exists \otimes_i ($i = 1, \dots, n$) and a function $H = h_1 \times \dots \times h_n$ where $h_1 = h$ such that for all i ,

$$\forall x, y. h_i(x \oplus_i y) = H x \otimes_i H y$$

then

$$h \circ \pi_1 \circ hom (\Delta_1^n \oplus_i) (\Delta_1^n f_i) = \pi_1 \circ hom (\Delta_1^n \otimes_i) (\Delta_1^n (h_i \circ f_i))$$

Notation note: crossing functions, sometimes called *maps*

$$(h_1 \times \dots \times h_n) (x_1, \dots, x_n) = (h_1 x_1, \dots, h_n x_n)$$

Example of applying almost fusion theorem

Suppose we want to fuse

$$\log \circ \pi_1 \circ \text{hom} (\Delta_1^2 \oplus_i) (\Delta_1^2 f_i)$$

where

$$\begin{aligned} f_1 x &= x & (x_{\text{mean}}, x_n) \oplus_1 (y_{\text{mean}}, y_n) &= \frac{x_{\text{mean}} \times x_n + y_{\text{mean}} \times y_n}{x_n + y_n} \\ f_2 x &= 1 & (x_{\text{mean}}, x_n) \oplus_2 (y_{\text{mean}}, y_n) &= x_n + y_n \end{aligned}$$

We must define

$$\begin{aligned} h_1 &= \log \\ h_2 &=? \\ \otimes_1 &=? \\ \otimes_2 &=? \end{aligned}$$

such that

$$\begin{aligned} h_1 (x \oplus_1 y) &= (h_1 \times h_2) x \otimes_1 (h_1 \times h_2) y \\ h_2 (x \oplus_2 y) &= (h_1 \times h_2) x \otimes_2 (h_1 \times h_2) y \end{aligned}$$

Finding h_2, \otimes_2

With

$$\begin{aligned}(x_{mean}, x_n) \otimes_2 (y_{mean}, y_n) &= x_n + y_n \\ h_2 x &= x\end{aligned}$$

then

$$\begin{aligned}h_2 (x \oplus_2 y) &= x_n + y_n \\ &= (\log x_{mean}, h_2 x_n) \otimes_2 (\log y_{mean}, h_2 y_n) \\ &= (\log \times id) x \otimes_2 (\log \times id) y \\ &= H x \otimes_2 H y\end{aligned}$$

With

$$\begin{aligned}(x_{mean}, x_n) \otimes_1 (y_{mean}, y_n) &= \log \left(\frac{\exp x_{mean} \times x_n + \exp y_{mean} \times y_n}{x_n + y_n} \right) \\ h_1 x &= \log x \\ h_2 x &= x\end{aligned}$$

then

$$\begin{aligned}h_1 (x \oplus_1 y) &= h_1 ((x_{mean}, x_n) \oplus_1 (y_{mean}, y_n)) \\ &= \log \left(\frac{x_{mean} \times x_n + y_{mean} \times y_n}{x_n + y_n} \right) \\ &= (\log x_{mean}, x_n) \otimes_1 (\log y_{mean}, y_n) \\ &= (\log \times id) x \otimes_1 (\log \times id) y \\ &= H x \otimes_1 H y\end{aligned}$$

Applying almost fusion

So we can fuse

$$\log \circ \pi_1 \circ \text{hom} (\Delta_1^2 \oplus_i) (\Delta_1^2 f_i)$$

to

$$\pi_1 \circ \text{hom} (\Delta_1^2 \otimes_i) (\Delta_1^2 (h_i \circ f_i))$$

where

$$h_1 x = \log x$$

$$h_2 x = x$$

$$(x_{\text{mean}}, x_n) \otimes_1 (y_{\text{mean}}, y_n) = \log \left(\frac{\exp x_{\text{mean}} \times x_n + \exp y_{\text{mean}} \times y_n}{x_n + y_n} \right)$$

$$(x_{\text{mean}}, x_n) \otimes_2 (y_{\text{mean}}, y_n) = x_n + y_n$$

$$mss = max^s \circ (map^s \text{ sum}) \circ \pi_1 \circ hom (\Delta_1^4 \oplus_i) (\Delta_1^4 f_i)$$

We need to find $h_1, h_2, h_3, h_4, \otimes_1, \otimes_2, \otimes_3, \otimes_4$ that solve these equations ($j = 1, \dots, 4$):

$$\begin{aligned} h_j ((s_1, i_1, t_1, d_1) \oplus_j (s_2, i_2, t_2, d_2)) = \\ (h_j s_1, h_2 i_1, h_3 t_1, h_4 d_1) \otimes_j (h_j s_2, h_2 i_2, h_3 t_2, h_4 d_2) \end{aligned}$$

Procedure: calculate LHS of equation j by promoting h_j into result and determine h_j, \otimes_j by matching with its RHS.

Example for $j = 1$

Equation

$$\begin{aligned} \text{map}^s \text{sum} ((s_1, i_1, t_1, d_1) \oplus_1 (s_2, i_2, t_2, d_2)) = \\ (\text{map}^s \text{sum } s_1, h_2 \ i_1, h_3 \ t_1, h_4 \ d_1) \otimes_1 (\text{map}^s \text{sum } s_2, h_2 \ i_2, h_3 \ t_2, h_4 \ d_2) \end{aligned}$$

Calculation

$$\begin{aligned} & \text{map}^s \text{sum} ((s_1, i_1, t_1, d_1) \oplus_1 (s_2, i_2, t_2, d_2)) \\ &= (\text{map}^s \text{sum}) (s_1 \cup s_2 \cup (t_1 \ \mathcal{X}_+ \ i_2)) \\ &= \text{map}^s \text{sum } s_1 \cup \text{map}^s \text{sum } s_2 \cup \text{map}^s \text{sum}(t_1 \ \mathcal{X}_+ \ i_2) \\ &= \text{map}^s \text{sum } s_1 \cup \text{map}^s \text{sum } s_2 \cup (t_1 \ \mathcal{X}_{\text{map}^s \text{sum} \circ +} \ i_2) \\ &= \text{map}^s \text{sum } s_1 \cup \text{map}^s \text{sum } s_2 \cup (\text{map sum } t_1 \ \mathcal{X}_+ \ \text{map sum } i_2) \end{aligned}$$

Matching last expression with RHS of the equation gives

$$\begin{aligned} h_2 = h_3 = \text{map sum} \\ (s_1, i_1, t_1, d_1) \otimes_1 (s_2, i_2, t_2, d_2) = s_1 \cup s_2 \cup (t_1 \ \mathcal{X}_+ \ i_2) \end{aligned}$$

Example for $j = 2$

Equation

$$\begin{aligned} \text{map sum } ((s_1, i_1, t_1, d_1) \oplus_2 (s_2, i_2, t_2, d_2)) = \\ (\text{map sum } s_1, h_2 \ i_1, h_3 \ t_1, h_4 \ d_1) \otimes_2 (\text{map sum } s_2, h_2 \ i_2, h_3 \ t_2, h_4 \ d_2) \end{aligned}$$

Calculation

$$\begin{aligned} & \text{map sum } ((s_1, i_1, t_1, d_1) \oplus_2 (s_2, i_2, t_2, d_2)) \\ &= \text{map sum } (i_1 \# \text{map } (d_1 \#) \ i_2) \\ &= \text{map sum } i_1 \# \text{map sum } (\text{map } (d_1 \#) \ i_2) \\ &= \text{map sum } i_1 \# \text{map } (\text{sum} \circ (d_1 \#)) \ i_2 \\ &= \text{map sum } i_1 \# \text{map } (\text{sum } d_1 \#) \ (\text{map sum } i_2) \end{aligned}$$

Matching last expression with RHS of the equation gives

$$\begin{aligned} h_4 &= \text{sum} \\ (s_1, i_1, t_1, d_1) \otimes_2 (s_2, i_2, t_2, d_2) &= i_1 \# \text{map } (d_1 \#) \ i_2 \end{aligned}$$

If we continue

$$h_1 = \text{map}^s \text{ sum}$$

$$h_2 = \text{map sum}$$

$$h_3 = \text{map sum}$$

$$h_4 = \text{sum}$$

$$(s_1, i_1, t_1, d_1) \otimes_1 (s_2, i_2, t_2, d_2) = s_1 \cup s_2 \cup (t_1 \mathcal{X}_+ i_2)$$

$$(s_1, i_1, t_1, d_1) \otimes_2 (s_2, i_2, t_2, d_2) = i_1 \uplus \text{map } (d_1 +) i_2$$

$$(s_1, i_1, t_1, d_1) \otimes_3 (s_2, i_2, t_2, d_2) = \text{map } (+d_2) t_1 \uplus t_2$$

$$(s_1, i_1, t_1, d_1) \otimes_4 (s_2, i_2, t_2, d_2) = d_1 + d_2$$

The f parts

To apply the almost homomorphism fusion theorem we also need fused f functions:

$$f_1 x = \{[x]\}$$

$$f_2 x = [[x]]$$

$$f_3 x = [[x]]$$

$$f_4 x = [x]$$

$$f'_1 x = ((map^s sum) \circ f_1) x = \{x\}$$

$$f'_2 x = ((map sum) \circ f_2) x = [x]$$

$$f'_3 x = ((map sum) \circ f_3) x = [x]$$

$$f'_4 x = (sum \circ f_4) x = [x]$$

And finally we obtain:

$$map^s sum \circ \pi_1 \circ hom (\Delta_1^4 \oplus_i) (\Delta_1^4 f_i) = \pi_1 \circ hom (\Delta_1^4 \otimes_i) (\Delta_1^4 f'_i)$$

Finishing up

$$mss = max^s \circ \pi_1 \circ hom (\Delta_1^4 \otimes_i) (\Delta_1^4 f'_i)$$

We can fuse max^s with the remaining almost homomorphism with

$$H = max^s \times max \times max \times id$$

and get the final almost homomorphism

$$mss = \pi_1 \circ hom (\Delta_1^4 \otimes'_i) id$$

where

$$(s_1, i_1, t_1, d_1) \otimes'_1 (s_2, i_2, t_2, d_2) = s_1 \uparrow s_1 \uparrow (t_1 + i_2)$$

$$(s_1, i_1, t_1, d_1) \otimes'_2 (s_2, i_2, t_2, d_2) = i_1 \uparrow (d_1 + i_2)$$

$$(s_1, i_1, t_1, d_1) \otimes'_3 (s_2, i_2, t_2, d_2) = (t_1 + d_2) \uparrow t_2$$

$$(s_1, i_1, t_1, d_1) \otimes'_4 (s_2, i_2, t_2, d_2) = d_1 + d_2$$

What has been accomplished?

We went from

$$mss = max^s \circ (map^s sum) \circ segs$$

to

$$mss = \pi_1 \circ hom (\Delta_1^4 \otimes'_i) id$$

- Specification was an inefficient $O(n^2)$ algorithm involving computationally difficult objects (sets).
- Final derivation has only (tuples of) scalars as intermediate values.
 - ▶ You could easily translate the final bit into Futhark or any other parallel language.

What do we make of all this?

- Easy to get lost in the details of formalisms.
- These theorems and techniques **do not give us algorithms for free**.
- They **help us structure our search**.
- Usually easier to find small functions that satisfy specific properties than to come up with algorithms from scratch using sheer creativity.
- (But maybe some of this could be mechanised? Maybe a fun (research) project!)