

See discussions, stats, and author profiles for this publication at: <https://www.researchgate.net/publication/360897960>

# Extraction of variations in agricultural practices over rice fields using unsupervised learning

Poster · May 2022

DOI: 10.13140/RG.2.2.26986.18880

CITATIONS

0

READS

22

4 authors:



**Thomas Di Martino**

CentraleSupélec

7 PUBLICATIONS 7 CITATIONS

[SEE PROFILE](#)



**Régis Guinvarc'h**

CentraleSupélec

122 PUBLICATIONS 455 CITATIONS

[SEE PROFILE](#)



**Elise Colin**

The French Aerospace Lab ONERA

92 PUBLICATIONS 849 CITATIONS

[SEE PROFILE](#)



**Laetitia Thirion-Lefevre**

CentraleSupélec

110 PUBLICATIONS 440 CITATIONS

[SEE PROFILE](#)

Some of the authors of this publication are also working on these related projects:



MEDUSA [View project](#)



Materials orientation and polarimetric signatures [View project](#)



# Extraction of variations in agricultural practices over rice fields using unsupervised learning

Thomas Di Martino<sup>1,2</sup>, Régis Guinvarc'h<sup>2</sup>, Elise Colin<sup>1</sup>, Laetitia Thirion-Lefevre<sup>2</sup>

1 ONERA, Traitement de l'information et systèmes, Université Paris-Saclay, 91123 Palaiseau, France

2 SONRA, CentraleSupélec, Université Paris-Saclay, 91190 Gif-sur-Yvette, France

A3.04 Agriculture - Methods and Algorithms, Science, Applications and Policy

living planet  
symposium | BONN  
23-27 May  
2022


## INTRODUCTION

Existing variations of agricultural practices within a single crop type:

- Various sowing, harvesting dates [1].
- Optional flooding of rice fields, for bird resting areas [2], seeding period...

**Convolutional Autoencoders** [3] can model and **extract variations within a single crop type** and **group them**, using different semantic criteria, **without supervision**.

Average Sentinel-1 backscatter temporal profile of rice fields over the studied area





## USED DATA

- Location of rice crops: Sector BXII, near Sevilla, Spain
- Details: 3500 ha of crops, growing in 2017.
- Sentinel-1 Data: 61 acquisitions (Jan 2017 to Dec 2017), dual-pol (VV, VH), preprocessing in [4]





Rice fields, in a Guadalquivir Marshes landscape, Andalusia, Spain (Source: [5])


Study Area



## AUTOENCODERS AND APPLICATION



Spatial visualization of groups of crops in the embedding space



Optional Flooding Period

Approximate Seeding Period

End of the year flooding Period

## CONCLUSION

The application of an autoencoder to SAR time series of rice crops allows to:


- Generate a 3D **embedding image** that **highlights groups** of crops.
- **Visualize** these **groups** of crops by mapping the embedding space to the **RGB** color space.
- Find out **which period** of a field contributes to **making the groups** of crops using the Grad-CAM methodology.

## REFERENCES

- [1] Alexandre Bouvet, Thuy Le Toan and Lam-Dao Nguyen, "Monitoring of the Rice Cropping System in the Mekong Delta Using ENVISAT/ASAR Dual Polarization Data," IEEE Transactions on Geoscience and Remote Sensing, vol. 47, no. 2, pp. 517-526, 2009.
- [2] Claire A. Pernollet, Anis Guelmami, Andy J. Green, Antoni Curcó Masip, Bosco Dies, Giuseppe Bogliani, Franco Tesio, Anne Brogi, Michel Gauthier-Clerc, Matthieu Guillemain, "A comparison of wintering duck numbers among European rice production areas with contrasting flooding regimes," Biological Conservation, vol. 186, pp. 214-224, 2015.
- [3] Thomas Di Martino, Régis Guinvarc'h, Laetitia Thirion-Lefevre and Elise Colin, "Beets or Cotton? Blind Extraction of Fine Agricultural Classes Using a Convolutional Autoencoder Applied to Temporal SAR Signatures," IEEE Transactions on Geoscience and Remote Sensing, vol. 60, pp. 1-18, 2022.
- [4] Alejandro Mestre-Quereda, Juan M. Lopez-Sanchez, Fernando Vicente-Guijalba, Alexander W. Jacob, and Marcus E. Engdahl, "Time-series of sentinel-1 interferometric coherence and backscatter for crop-type mapping," IEEE Journal of Selected Topics in Applied Earth Observations and Remote Sensing, vol. 13, pp. 4070-4084, 2020.
- [5] [https://es.wikipedia.org/wiki/Marismas\\_del\\_Guadalquivir](https://es.wikipedia.org/wiki/Marismas_del_Guadalquivir)
- [6] Ramprasaath R. Selvaraju, Abhishek Das, Ramakrishna Vedantam, Devi Parikh and Dhruv Batra, "Grad-cam: Visual explanations from deep networks via gradient-based localization," IEEE International Conference on Computer Vision, pp. 618-626, 2017.


## RESULTS AND VISUALISATION

Red channel: 1st embedding dimension




Similarities between crops in shades of red imply similarities in harvest strategies in key periods (i.e. Till, Seeding, Harvest, End of the year flooding)

Extraction of date importance for 1st embedding generation using the Grad-CAM method [6]




Periods of interest for 1st embedding:  
• All, but start of the year optional flooding

Green channel: 2nd embedding dimension




Extraction of date importance for 2nd embedding generation using the Grad-CAM method [6]




Periods of interest for 2nd embedding:  
• End of Field preparation  
• Seeding  
• Harvest  
• End of the year optional flooding

Blue channel: 3rd embedding dimension



Extraction of date importance for 3rd embedding generation using the Grad-CAM method [6]



Periods of interest for 3rd embedding:  
• Start of the year optional flooding  
• Growth