




# Unravelling the Impact of Halogen Interactions in Medicinal Chemistry

Christoph P. Sager



SCS Spring School on Digital Chemistry – 19 April 2023

# Agenda





- Halogen bonds and their interactions
- Exercise #2.1
- Application of halogen bonds in our projects
- Sulfur-oxygen interactions
- Fluorine multipolar interactions
- Exercise #2.2
- Summary



# Halogen bonds

# Halogen bonding – the $\sigma$ -hole

Scholfield et al. / Shinada et al. / Cramer et al.





- Group VII has 5 electrons in the  $p$ -atomic orbitals
  - Single valence electron ( $p_z$  orbital) forms the covalent  $\sigma$ -bond
  - The opposite side of the  $p_z$  orbital is depopulated  $\rightarrow$   $\sigma$ -hole
  - Remaining 4 electrons ( $p_{xy}$  orbitals) form the negative belt

VII

9	F	Fluorine	Halogen
17	Cl	Chlorine	Halogen
35	Br	Bromine	Halogen
53	I	Iodine	Halogen
85	At	Astatine	Halogen
117	Ts	Tennessine	Halogen

# Halogen bonding – the $\sigma$ -hole



Scholfield et al. / Shinada et al. / Cramer et al.





- Group VII has 5 electrons in the  $p$ -atomic orbitals
  - Single valence electron ( $p_z$  orbital) forms the covalent  $\sigma$ -bond
  - The opposite side of the  $p_z$  orbital is depopulated  $\rightarrow$   $\sigma$ -hole
  - Remaining 4 electrons ( $p_{xy}$  orbitals) form the negative belt
- Polarizability/Size:  $F < Cl < Br < I \sim 4-8 \text{ kJ mol}^{-1}$ 
  - Electron withdrawing partner:  $16 \text{ kJ mol}^{-1}$
- Classical hydrogen bonds:  $18 - 21 \text{ kJ mol}^{-1}$
- Desolvation cost of a hydroxyl group:  $11 - 24 \text{ kJ mol}^{-1}$

# Halogen bonding - geometries

Scholfield et al. / Bissantz et al.




- Distance:  $\leq \Sigma r_{vdW}$  Å,
  - $\leq 3.24$  Å,  $\leq 3.37$  Å,  $\leq 3.5$  Å
- Linearity ( $\Theta_1$ ) =  $160^\circ$ – $165^\circ$
- Planarity ( $\Theta_2$ ) =  $120^\circ$




# X-bond interactions

Shinada et al.




Tyr228 in Factor Xa



- 2pr3
- 3q3k
- 4y71
- 2boh



Cl



Br



Dataset

Non-redundant



# Exercise #2.1

## Cathepsin L & MAP ERK kinase



**Odd-Groups** (1, 3, 5, ...)

Open: Cathepsin\_L\_example.pdb

**Even-Groups** (2, 4, 6, ...)

Open: MAP\_ERK\_kinase\_example.pdb

Goal: Add/Change only 1 atom to improve affinity


Time: 10 min

Presentation: Two groups (1 odd & 1 even group), 2 min each



# Application of halogen bonds in our projects

# Crystallography enables virtual screening

How to find a halogen bond, step by step




- Crystal Structures: 120+ (avg. 1.4 Å)
- Starting point: 4500 nM
- Docking: 14'000
- Synthesized: 17 (220 – 4000 nM)
- Follow-up: 27 (90 – 9000 nM)
- 2nd follow-up: 17 (80 – 350 nM)





## Knowledge transfer to other series




The image displays a collection of 20 chemical structures and 5 question marks, arranged in a grid. The structures are various substituted benzene rings with different functional groups and substituents. The substituents are color-coded: Br (brown), Cl (green), F (light green), N (blue), O (orange), and CN (dark blue). The question marks are red.

The structures are as follows:

- Row 1: 1-bromo-2-chlorobenzene, 1-fluoro-2-chlorobenzene, 1-chloro-3-methylbenzene, 1-fluoro-2-chlorobenzene, 1-chloro-3-methylbenzene, 1,1-difluoro-2-chlorobenzene, 1-chloro-3-chlorobenzene, and a red question mark.
- Row 2: 1-chloro-2-methylbenzene, 1-chloro-2-methylbenzene, a red question mark, 1-methoxy-2-chlorobenzene, a red question mark, 1,2-dichlorobenzene, a red question mark, 1-chloro-2-methylbenzene, and 1-bromo-2-chlorobenzene.
- Row 3: 1-chloro-2-methylbenzene, a red question mark, 1-cyano-2-chlorobenzene, 1-bromo-2-chlorobenzene, 1-cyano-2-chlorobenzene, 1,2-dichlorobenzene, 1-chloro-2-methylbenzene, and 1-methoxy-2-chlorobenzene.

# Increasing the $\sigma$ -hole effect

Electrostatic surface potentials




	Phenyl	Meta-F	Meta-Cl	Meta-Br	Meta-meta-Cl	Meta-meta-Br	Meta-Cl meta-cyano	Meta-Br meta-cyano	Meta-Cl para-CF <sub>3</sub>
ESP [kcal/mol]	20.3	5.6	9.7	12.9	13.2	16.5	17.3	20.6	12.4
IC <sub>50</sub> [nM]	67 (n = 4)	49 (n = 1)	16 (n = 6)	22 (n = 3)	13 (n = 16)	12 (n = 4)	15 (n = 7)	11 (n = 4)	147 (n = 1)

Jaguar  
Theory: M06-2X  
Basis-Set: CC-PVDZ++



# Increasing the $\sigma$ -hole effect

Affinity vs. ESP



Jaguar



Theory: M06-2X

Basis-Set: CC-PVDZ++



# Crystal structure analysis

Gabriele Conti




72 crystal structures analyzed in term of:

- $d_{X \cdots O}$
- $\Theta_1$
- $\Theta_2$
- $\sum r_{vdW}$
- $\leq 3.24 \text{ \AA}, \leq 3.37 \text{ \AA}, \leq 3.5 \text{ \AA}$
- $\Delta d (d_{X \cdots O} - \sum r_{vdW})$

# Crystal structure analysis

$$\Delta d (d(X \cdots O) - \sum r_{vdW})$$




Optimal values:

- $\Delta d (d(X \cdots O) - \sum r_{vdW}) < 0$
- The shorter the X – O distances, the higher the affinities

# Crystal structure analysis

Fluorine  $\Theta_1$




Optimal values:

- $\Delta d (d \text{ X} \cdots \text{O} - \sum r_{\text{vdW}}) < 0$
- The shorter the X – O distances, the higher the affinities
- Literature:  $\Theta_1 \approx 160 - 165^\circ$
- No fluorine halogen bond



# Crystal structure analysis

Bromine  $\Theta_1$




Optimal values:

- $\Delta d (d \text{ X} \cdots \text{O} - \sum r_{\text{vdW}}) < 0$
- The shorter the X – O distances, the higher the affinities
- Literature:  $\Theta_1 \approx 160 - 165^\circ$
- Para substituent lower affinity

# Crystal structure analysis

Chlorine  $\Theta_1$




Optimal values:

- $\Delta d (d \text{ X} \cdots \text{O} - \sum r_{\text{vdW}}) < 0$
- The shorter the X – O distances, the higher the affinities
- Literature:  $\Theta_1 \approx 160 - 165^\circ$
- Para substituent lower affinity

# Crystal structure analysis

## Summary $\Theta_1$




Optimal values:

- $\Delta d (d_{X-O} - \sum r_{vdW}) < 0$
- The shorter the X – O distances, the higher the affinities
- Literature:  $\Theta_1 \approx 160 - 165^\circ$
- Compounds with  $\Theta_1$  values close to  $156 - 164^\circ$  show the highest activity
- Para substituent influences affinity:  $CF_3 < CH_3 < Cl < F < H$
- Lipophilic groups allowed



# Crystal structure analysis

## Summary $\Theta_1$




Optimal values:

- $\Delta d (d \text{ X} \cdots \text{O} - \sum r_{\text{vdW}}) < 0$
- The shorter the X – O distances, the higher the affinities
- Literature:  $\Theta_1 \approx 160 - 165^\circ$
- Compounds with  $\Theta_1$  values close to  $156 - 164^\circ$  show the highest activity
- Para substituent influences affinity:  $\text{CF}_3 < \text{CH}_3 < \text{Cl} < \text{F} < \text{H}$
- Lipophilic groups allowed


# Crystal structure analysis

## Summary $\Theta_2$



Optimal values:


- Literature:  $\Theta_2 \approx 120^\circ$
- Our observed range: 160 – 175°
- $\Theta_2$  less impact on affinity



# Summary

## Halogen Bonds in Medicinal Chemistry


- Crystal structure support is a great asset for challenging projects
- In depths (geometric) analysis of high-resolution structures combined with QM calculations lead to more specific SAR understanding
- Proper understanding of the interactions involved in a system is essential for making informed decisions and optimizing outcomes.



# Sulfur-oxygen interactions


# Sulfur-oxygen interaction – $\sigma$ -hole part II

Zhang et al.






# Sulfur-oxygen interaction at Idorsia

Benzothiazoles with a  $\sigma$ -hole



	Benzothiazole	Benzothiazole-Me
IC <sub>50</sub> [nM]	20 (n = 5)	31 (n = 3)









# Fluorine multipolar interactions

# Fluorine in Pharmaceuticals: Looking Beyond Intuition



Diederich group



**Prozac**



**Lipitor**




**Ciprobay**


- Fluorine substituents are widely used in drugs due to the development of safe and selective fluorinating agents.
- Fluorine has significant effects on the physical and pharmacokinetic properties of a drug compound (ADME).
- The inductive effects of fluorine are relatively well understood and can enhance bioavailability by reducing the basicity of neighboring amines.

# C-F...H-N interactions

Diederich group









- The C-F unit is **not a good** acceptor for hydrogen bonds because it has a low proton affinity and is weakly polarizable.
- A study of **788** C-F containing structures found that in only **11** of them, the C-F part of the molecule points towards the H-N bond when the F...N distance is below **3.1 Å** and the angles  $\alpha_1$  and  $\alpha_2$  meet certain criteria ( $\alpha_1 \geq 150^\circ$  and  $90^\circ \leq \alpha_2 \leq 150^\circ$ )
- Thrombin inhibitor comparison: The F-containing inhibitor is shown to be **5 times more potent** than the H-containing inhibitor.



# C–F...C=O interactions

Diederich group



- Distance:  $2.8 \text{ \AA} < d_1 < 3.8 \text{ \AA}$ 
  - $C_{\text{vdW}}: 1.7 \text{ \AA}$
  - $F_{\text{vdW}}: 1.47 \text{ \AA}$
- Linearity ( $\alpha_1$ ) =  $180^\circ$ 
  - Weaker dependence at longer distance
- Planarity ( $\alpha_2$ ) =  $90^\circ \pm 10\text{-}40^\circ$ 
  - Distance dependent
- $\Delta\Delta G$  from  $C_{\text{sp}^2}\text{-H}$  to  $C_{\text{sp}^2}\text{-F}$ 
  - $-1.0 \pm 0.2 \text{ kJ mol}^{-1}$

# Exercise #2.2

## Galectin-3

## All Groups

Open: Galectin-3\_example.pdb

**Goal:** Implement what you have learned

- Halogen bonds
- Sulfur-oxygen interaction
- Fluorine multipolar interactions


**Time:** 15 min

**Presentation:** Min. two groups



# Summary

## Interactions in Medicinal Chemistry



- Proper understanding of the interactions involved in a system is essential for making informed decisions and optimizing outcomes.
  - Perfectly placed halogens and thiazoles enhance protein-ligand affinity and may serve as core interactions for lead discovery projects.
  - Fluorine residues can enhance physicochemical properties to improve ADME properties and strengthen protein-ligand binding interactions.
- Academic research plays a critical role in advancing our understanding of complex systems and developing new solutions.
- Continuous learning and improvement is essential for staying up-to-date and making meaningful contributions to scientific research and innovation.

# Acknowledgement

Modeling	Christoph Sager, Gabriele Conti, Benedetta Girardi, Elisa Liberatore
Crystallography	Alain Chambovey, Aengus Mac Sweeney, Manon Müller, Micha Wicky
Biology	John Gatfield, Celia Müller
Chemistry	Christine Berthion, Martin Bolli, Edouard Ehret, Michael Erhardt, Sven Glutz Corinna Grisostomi, Claire Hinder, Daniela Kruesi, Claus Mueller, Marc Masmünster, Jean-Marie Pilven, Luboš Remeň, Gunther Schmidt, Jürgen Seifert, Christina Stamm, Marco Tschanz, Rene Vogelsanger, Raphael Vogt, Cornelia Zumbrunn



Curious  
to learn more?



# References

## Constantly expanding our knowledge

- General

- A systematic analysis of atomic protein–ligand interactions in the PDB, *Med. Chem. Commun.*, 2017, 8, 1970
- Conformational constraint in protein ligand design and the inconsistency of binding entropy, *Biopolymers*, 2008, 8, 653-667
- Aromatic interactions in model systems, *Curr. Op. Chem. Biol.*, 2002, 6, 736-741
- A Medicinal Chemist's Guide to Molecular Interactions, *J. Med. Chem.* 2010, 53, 5061–5084
- Hydroxyl Groups in Synthetic and Natural-Product-Derived Therapeutics: A Perspective on a Common Functional Group, *J. Med. Chem.* 2019, 62, 20, 8915–8930
- Small Molecule Conformational Preferences Derived from Crystal Structure Data. A Medicinal Chemistry Focused Analysis, *J. Chem. Inf. Model.* 2008, 48, 1-24

- Protein Structures

- [Electron density maps - Proteopedia, life in 3D](#)
- [PDBe | Protein Data Bank in Europe \(ebi.ac.uk\)](#)
- [Coot website](#)
- Features and development of Coot, *Acta Crystallographica Section D* 66, 2010, 486-501
- [PyMol website](#)

- C-X...O

- Halogens in Protein–Ligand Binding Mechanism: A Structural Perspective, *J. Med. Chem.* 2019, 62, 21, 9341-9356
- Halogen bonding (X-bonding): A biological perspective, *Protein Science*. 2013, 22:139-152
- Halogen Bonds: Benchmarks and Theoretical Analysis, *J. Chem. Theory Comput.* 2013, 9, 1918-1931
- Halogen Bonding Interactions: Revised Benchmarks and a New Assessment of Exchange vs. Dispersion, *J. Chem. Theory Comput.* 2018, 14, 1, 180–190

- C-F...O

- The many faces of fluorine: Some noncovalent interactions of fluorine compounds, *J. Chem. Thermodynamics* 156 (2021) 106382

- S...O

- Intermolecular Sulfur...Oxygen Interactions: Theoretical and Statistical Investigations, *J. Chem. Inf. Model.* 2015, 55, 10, 2138–2153

