



Trends in Survival based on Treatment Modality in Lung Cancer Patients: A Population-Based Study

Journal:	<i>Cancer Investigation</i>
Manuscript ID	LCNV-2019-OR-0022
Manuscript Type:	Original Article
Date Submitted by the Author:	21-Jan-2019
Complete List of Authors:	Akhtar-Danesh, Noori; McMaster University, School of Nursing and Department of Health Research Methods, Evidence, and Impact Akhtar-Danesh, Gileh-Gol; McMaster University, Department of Surgery Seow, Hsien; McMaster University, Department of Oncology; McMaster University, Department of Health Research Methods, Evidence, and Impact Shakeel, Saad; University of Toronto, School of Medicine Finley, Christian; McMaster University, Division of Thoracic Surgery, Department of Surgery
Keywords:	Non-small cell lung cancer, Survival analysis, Treatment modality, Trends in survival

SCHOLARONE™
Manuscripts

**Trends in Survival based on Treatment Modality in Lung Cancer Patients:
A Population-Based Study**

Noori Akhtar-Danesh^{1,2}, PhD

Gileh-Gol Akhtar-Danesh³, MD

Hsien Yeang Seow^{2,4}, PhD

Saad Shakeel⁵, MPH

Christian Finley⁶, MD, FRCS(C)

¹School of Nursing, McMaster University

²Department of Health Research Methods, Evidence, and Impact, McMaster University

³Department of Surgery, McMaster University

⁴Department of Oncology, McMaster University

⁵School of Medicine, University of Toronto

⁶Division of Thoracic Surgery, Department of Surgery, McMaster University

Conflict of Interest

The authors have no conflicts of interest to declare.

Corresponding Author:

Noori Akhtar-Danesh,
School of Nursing,
McMaster University,
1280 Main St. West, Room 3N28B
Hamilton, Ontario, L8S 4K1
Canada
daneshn@mcmaster.ca
Tel: 905-525-9140 Ext. 22297
Fax: 905-521-8834

ABSTRACT

Objectives: We compared the trends in survival based on treatment modality among lung cancer patients in the province of Ontario, Canada, from 2007 to 2015.

Methods: We compared the trends in survival based on treatment modality.

Results: Among 56,417 identified patients, the mean age at diagnosis was 70.1 years (SD=10.7). Treatment modalities varied significantly over time ($p < 0.001$). Overall, 23.0% of patients received surgical treatments. We observed more than 20% increase in five-year survival rates for all surgical groups over time.

Conclusions: Patients undergoing sublobar/lobar resections had the best survival.

Key words: Non-small cell lung cancer; Survival analysis; Treatment modality; Trends in survival

Introduction

Lung cancer remains the leading cause of cancer-related mortality worldwide (1). In 2017, lung cancer represented 17% of all new cancer diagnoses in Canada. It was also the leading cause of cancer death, with its mortality surpassing the combined mortalities of colorectal, breast, and prostate cancers (2). Similar figures have been reported in the United States, where lung cancer represented nearly 14% of all new cancer diagnoses, and 27% of all cancer-related deaths (3). The high mortality rate is partly attributable to the initial relatively asymptomatic nature of lung cancer patients. By the time symptoms manifest, the malignancy is often advanced and curative options are no longer available (4). Furthermore, there is still no screening test that can be implemented at a population-level, although provincial and national medical associations have proposed screening programs for high-risk patients (5).

For patients in whom the disease is identified early, there are several treatment modalities available. In those who are medically fit and able to tolerate an operation, surgical resection is the treatment of choice. Patients unfit for surgery can undergo a combination of radiation and chemotherapy (6). The decision-making process surrounding treatment modalities for patients with lung cancer is quite complex, and confounded by the existing co-morbidities, as the surgery places a tremendous physiologic demand on the cardiovascular and respiratory systems. In patients with early disease identification, the five-year survival can be as high as 73% compared to a five-year survival of 2-13% in patients with metastatic disease (7). Nevertheless, there is a paucity of literature comparing the associated effects of various treatment modalities for lung cancer on overall survival.

In this study, we investigated different treatment modalities, their potential covariates and effects on the survival of Canadian patients in the province of Ontario. To our best knowledge,

1
2
3 this is the first population-based study in Canada to investigate trends in survival based on
4
5 treatment modality among lung cancer patients. We explored the survival differences between
6
7 varying treatment modalities as the main independent variable. The analysis was adjusted for
8
9 demographics and tumor characteristics such as stage and histology of tumor at the time of
10
11 diagnosis. This study was limited to patients with non-small cell lung cancer, as the natural
12
13 history and prognosis of small cell lung cancer is drastically different from the non-small cell
14
15 population.
16
17
18
19

20 **Materials and methods**

21 ***Data Source***

22
23
24 The main dataset used in this analysis was the Ontario Cancer Registry (OCR) dataset,
25
26 which includes diagnosis date, stage, and tumor characteristics for each cancer patient. The OCR
27
28 includes all diagnosed cases of invasive neoplasia, except basal cell and squamous cell skin
29
30 cancers, in Ontario (8). For this analysis we merged the OCR with several population-based or
31
32 administrative datasets held at the Institute for Clinical Evaluative Sciences (ICES). These
33
34 datasets include: Cancer Activity Level Reporting (chemotherapy and radiation), Discharge
35
36 Abstracts Database (surgery, radiation, and chemotherapy), National Ambulatory Care Reporting
37
38 System (chemotherapy and radiation), Ontario Health Insurance Plan Claims (chemotherapy and
39
40 radiation), Same Day Surgery datasets (chemotherapy and radiation), and Registered Person
41
42 Database (age, sex, date of death). These datasets were merged to include all Ontario residents
43
44 diagnosed with primary invasive non-small cell lung cancer (NSCLC) with known disease stage
45
46 from 2007 to 2015, with follow-up to the end of 2016.
47
48
49
50
51

52
53 Using the International Statistical Classification of Diseases and Related Health
54
55 Problems, Tenth Revision (ICD-10), we identified patients with NSCLC with the codes of C34.0
56
57
58
59
60

(malignant neoplasm of main bronchus), C34.1 (malignant neoplasm upper lobe, bronchus or lung), C34.2 (malignant neoplasm of middle lobe, bronchus or lung), C34.3 (malignant neoplasm of lower lobe, bronchus or lung), C34.8 (overlapping malignant lesion of bronchus or lung), and C34.9 (malignant neoplasm of bronchus or lung unspecified) in the OCR dataset. Then, using the International Classification of Diseases for Oncology (ICD-O-2 or ICD-O-3), we excluded all small cell carcinoma, carcinoid tumors, salivary gland tumors, mesenchymal tumors, lymphoproliferative tumors, and miscellaneous tumors (9). Patients were also excluded if the diagnosis was solely based on death certificate or autopsy as the survival time could not be identified for these patients. The analysis included patients who were 18 years and older at the time of diagnosis. Age at diagnosis was then classified as < 60, 60-69, 70-79, and \geq 80 years. This analysis includes only patients with known disease stage, as staging and age are important factors that are taken into consideration when deciding on treatment modalities (6, 10).

Treatment modalities were investigated based on the Canadian Classification of Health Information (CCI) (11) and Canadian Classification of Diagnostic, Therapeutic, and Surgical Procedures (CCP) (12). Treatment modalities were classified as: 1) *no-treatment* for patients who received no treatment, or we could not find any record of treatment, 2) *radiation only* (RT) for patients that received only radiation therapy, 3) *chemotherapy only* (CT) for patients who received only chemotherapy, and 4) *chemoradiation* for patients who received both radiation and chemotherapy but no surgical resection. The surgical groups include: 5) *sublobar resection or lobectomy only* (Sub/Lob only) with no record of radiation or chemotherapy for this group, 6) *sublobar resection or lobectomy* with radiation or chemotherapy (Sub/Lob + RT/CT), 7) *pneumonectomy* with no record of radiation or chemotherapy (*pneumonectomy only*), and finally, 8) *pneumonectomy* with radiation or chemotherapy (*pneumonectomy* + RT/CT). To determine

1
2
3 treatment classifications, a preliminary analysis was conducted to examine the Kaplan-Meier
4 survival curves for all surgical and non-surgical groups. The Kaplan-Meier curves were adjusted
5
6 for patient's age, stage of disease, and histology of tumor. Then, based on the similarity of the
7
8 Kaplan-Meier curves (Figure A.1) sublobar-only and lobectomy-only resection groups were
9
10 combined into one group (Sub/Lob only). Also, sublobar and lobectomy surgical groups with
11
12 radiation or chemotherapy were combined into one group (Sub/Lob + RT/CT). For each patient,
13
14 a treatment modality was accounted for and included in the analysis only if it was received up to
15
16 one year after diagnosis.
17
18
19
20
21

22 Staging data has been made available in Ontario since 2007. Cancer Care Ontario uses a
23
24 best stage grouping approach where stage is assigned to each case based on pathologic TNM
25
26 when available, and clinical TNM otherwise (13). It collects stage data using the staging criteria
27
28 of the American Joint Committee on Cancer or the Collaborative Stage initiative. If a case has
29
30 more than one valid stage value, the "best stage" is derived based on a pre-specified algorithm;
31
32 Otherwise in case of availability, the stage group provided by the treating regional cancer centre
33
34 is used (14). This analysis only includes patients with known cancer stage.
35
36
37
38

39 Tumor histology was classified as: 1) carcinoma-NOS, 2) squamous cell carcinoma
40
41 (SCC), 3) adenocarcinoma-NOS (AC), or 4) other.
42
43

44 **Statistical Analysis**

45
46 First, we conducted a logistic regression analysis to quantify the effects of demographics
47
48 (sex and age groups) and tumor characteristics (stage and histology) on receiving any surgical
49
50 resection. Treatment modalities were regrouped as *surgery* and *no-surgery* (reference group).
51
52 The surgery group included all surgical resection groups with or without any adjuvant
53
54 chemotherapy or radiation. The no-surgery group included all the non-surgical groups.
55
56
57
58
59
60

Second, the survival rate was estimated for patients diagnosed with NSCLC from 2007 and 2015 in Ontario with follow-up until the end of 2016. We used a flexible parametric (Royston-Parmar) model (15, 16) to estimate the trends in survival rate based on treatment modality, adjusted for sex, age, year of diagnosis, stage of tumor, and histology of tumor. This method uses a piecewise approach in modeling, which is more capable of mimicking the actual trends in mortality (hazard rate) and survival patterns compared to other traditional methods (17). We fitted the model by incorporating age, sex, year of diagnosis, histology of tumour, stage of disease, treatment modality, and the interaction term between each two variables into a multiple statistical model using a forward approach. The likelihood ratio test was used to compare different models, and to reach to a final model. All variables included in the final model were statistically significant ($p \leq 0.05$). Then, based on the final model, trends in one-, two-, and five-year survival rates for each treatment modality were estimated while adjusting for the other variables.

All the analyses were conducted using Stata/MP 15.1 (Stata Corporation, College Station, TX). The flexible parametric model was fitted using the freely available **stpm2** program (15) written for Stata package. The local ethics board at McMaster University, Hamilton, Canada, reviewed the research proposal and provided ethical approval for this retrospective study.

Results

Overall, 56,417 patients from the OCR dataset with invasive NSCLC were identified, of which 52.3% (n = 29,499) were male. The mean age at diagnosis was 70.1 years (SD=10.7); 70.5 (SD=10.3) years for males and 69.7 (SD=11.0) years for females. Approximately, 49.1% of patients were diagnosed with stage IV disease (50.4% of males and 47.7% of females). Adenocarcinoma was the most common subtype of NSCLC, accounting for 36.0% of cases.

Table 1 summarizes the distribution of sex, age group, histology of tumor, stage of disease, and year of diagnosis based on treatment modality. Overall, 23.0% of patients received surgical treatments, 26.3% radiation only, 7.2% chemotherapy only, and 21.0% chemoradiation; 22.5% did not receive any active treatment. We found a significant difference in the treatment modalities between males and females. About 21.0% of males underwent operative intervention, while 25.3% of females received surgical treatment ($p < 0.001$). The differences in treatment modalities were even more pronounced based on age group. The proportion of patients who underwent surgical resection was 26.1%, 28.3%, 24.7%, and 10.2% for age groups <60 , 60-69, 70-79, and ≥ 80 years, respectively. Older patients were more likely to receive no treatment or radiation only, with the proportion of no-treatment or radiation only varying from 30.2% to 78.0% from the youngest to oldest age group (Table 1). However, the largest discrepancy in treatment modality was based on disease stage. While 62.8% of patients with stage I disease received surgical treatment, only 2.1% of patients with stage IV disease underwent operative intervention. Conversely, 62.0% of patients with stage IV disease received no treatment or radiation only, compared to 32.1% of patients with stage I disease. The proportion of patients receiving chemotherapy only varied from 1.35% to 11.0% for patients with stage I to stage IV disease. The proportion of patients who received chemoradiation varied from 3.8% in stage I disease to 33.8% among patients with stage III disease.

Changes in treatment modalities

As can be seen in Table 1, the changes in the treatment modalities were quite complex. Statistical analysis based on a Chi-square test indicated that the distribution of treatment modalities was not the same over the nine years of the study period ($p < 0.001$). Two obvious trends that can be found are: 1) the proportion of radiation only increased from 22.4% in 2007 to

28.7% in 2015 (p-for-trend < 0.001); and 2) the proportion of pneumonectomy, with or without radiation or chemotherapy, decreased from 2.4% in 2007 to 1.0% in 2015 (p-for-trend < 0.001). However, there has not been a steady increase in the resection rate over time. In general, the resection rate decreased from 24.8% in 2007 to 20.9% in 2010 and then increased to 25.7% in 2015. However, the linear trend indicated a significant increase from 2007 to 2015 (p-for-trend < 0.001).

Predictors of surgical resection

Table 2 presents the predictors of surgical resection as described earlier using a multiple logistic regression. It predicts the chance of surgical resection based on sex, age group, year of diagnosis, histology and stage of tumor. The results are presented as odds ratio (OR) with 95% confidence interval (95% CI). Female patients were 10% more likely to receive surgical resection compared to male patients (OR 1.10; 95% CI: 1.05- 1.17). The odds ratios for surgical resection for patients 70-79 years and 80 and over were 0.48 (95% CI: 0.44- 0.52) and 0.12 (95% CI: 0.11- 0.14), respectively, compared to patients < 60 years. The chance of surgical resection for patients with SCC and AC tumors was 3.26 (95% CI: 2.97- 3.58) and 3.40 (95% CI: 3.12- 3.73) times higher compared to patients with carcinoma NOS. Most importantly, those with a stage IV disease had merely a 1% chance of receiving a surgical resection compared to those with a stage I disease.

Trends in survival rate based on treatment modality

The final model from fitting a flexible parametric model included sex, age group, treatment modality, stage of disease, histology of tumor, and year of diagnosis. The following interaction terms were also included in the model: age group and stage, age group and histology, treatment modality and year of diagnosis, treatment modality and stage, and stage and year of

diagnosis. In addition, age group, stage of tumor, treatment modality, histology of tumor, and year of diagnosis were included as time dependent variables. Then, based on the final model, trends in one-, two-, and five-year survival rates from 2007 to 2015 were estimated for each treatment modality. Results of fully the adjusted survival rates are shown in Figure 1. In general, we observed increasing trends in survival for all treatment modalities from 2007 to 2015, although it was more pronounced for the surgical resection groups. We observed an approximately 10% increase in two-year survival and an increase of more than 20% in five-year survival rates for all surgical groups.

We observed an increase of 10-15% in one-, two-, and five-year survival rate based on age group and sex from 2007 to 2015. Results of the fully adjusted survival model for five-year survival are shown in Figure 2. The trends were similar for all age groups, although a higher survival was evident for women across all age groups. Figure 3 depicts the trends in one-, two-, and five-year survival rates based on stage of disease. Although all one-, two-, and five-year survival rates improved for patients with stages I-III disease, they were less pronounced for patients with stage IV disease.

Overall five-year survival rate

The model-based average five-year survival rates based on histology, stage, and treatment modality are shown in Table 3. In general, the highest five-year survival rates were observed for the *Sub/Lob only* surgical resection group, followed by the *Sub/Lob + RT/CT* group. In comparison, the lowest survival rates were observed for the no-treatment group, the RT only group, and the CT only group. Chemoradiation seems to be more effective among patients with adenocarcinoma than carcinoma NOS and squamous cell carcinoma.

Discussion

In this large retrospective population-based study, we found that treatment modality was dependent on a patient’s age, sex, and histology and stage of disease among lung cancer patients in Ontario. The overall survival of patients with lung cancer increased over the 9-year study period. The differences in survival rates highly varied based on treatment modality, with surgical resection offering the greatest survival advantage.

Surgery is the mainstay of treatment for non-small cell lung cancer, and offers the best chance of cure or longer survival (18). However, due to the physiologic burden associated with lung resection, not all patients are appropriate candidates for surgery (18). Furthermore, given the often advanced stage of disease at the time of diagnosis, curative resection may not be feasible (6). In our study, only 23.0% of patients received surgery. This is similar to a large study from the United States in which 29.1% of patients were treated surgically (19). Women were significantly more likely to undergo an operation, as were the youngest group of patients. In contrast, elderly patients were more likely to receive no treatment or receive radiation only. These findings are consistent with the existing body of literature. Women with lung cancer tend to be younger at the time of diagnosis, and are less likely to report significant comorbidities, which may account for differences in resection rates (20, 21). However, our analysis indicates a higher chance of surgical resection for women after adjusting for age, stage, and histology of tumor. Previous studies have also reported lower perioperative morbidity and mortality in women undergoing lung resections (21). In contrast, elderly patients with NSCLC are more likely to be offered radiation only, or best supportive care, despite existing evidence regarding the safety of surgery in selected elderly patients (22-24).

The survival of patients with NSCLC increased over the study period. The difference in survival was most pronounced for the younger age groups, patients with stages I to III disease, and those undergoing operative intervention. As outlined above, numerous studies have shown the safety of surgery in carefully selected elderly patients (22-24). The improvement in survival of patients over the age of 80 from 2007 to 2015 (Figure 2) suggests that these patients are actually receiving treatment with curative intent, as opposed to best supportive care. This hypothesis is supported by further analysis that more elderly patients were treated with radiation-only towards the latter portion of the study period (see Table A.1). The observed differences in survival based on disease stage are not surprising, and in keeping with the poor prognosis of advanced stages of lung cancer (6). This study highlights the differences in survival based on treatment modality, while controlling for disease stage, information that is invaluable to patients and clinicians. The improved survival of patients with stages I to III from 2007 to 2015 is likely attributable to the improved medical care of these patients, including advances in radiation and peri-operative care. These results are similar to a recent American study, which examined the survival of lung cancer patients within the Surveillance, Epidemiology, and End Results (SEER) database (25). Another large study from Singapore also reported improvement in the survival of lung cancer patients over time, and attributed this to the increasing incidence of never-smokers in patients with NSCLC (26). Never-smokers represent an increasing proportion of patients with non-small cell lung cancer (27); however, the smoking status of patients was outside the scope of this study. Since the introduction of screening recommendations by the Canadian Medical Association in 2016, it can be hypothesized that the survival of lung cancer in Canada may continue to improve due to early detection. In contrast to the patients with local or regional

disease, the overall survival of patients with stage IV lung cancer continues to be poor, despite advances in targeted therapies (28).

Patients undergoing surgical resection via a sublobar/lobar resection had the best survival rates. Patients undergoing a pneumonectomy, with or without radiation or chemotherapy, had improved survival compared to those undergoing no treatment, radiation only, chemotherapy only, or chemoradiation but poorer survival than the lobar/sublobar resection patients. This is likely attributable to the increased morbidity associated with a pneumonectomy (29). As per guidelines put forth by the National Comprehensive Cancer Network (NCCN), radiation therapy currently plays a role in all stages of NSCLC (30). For patients with early disease, radiation therapy is offered to patients medically unfit for surgery, or those who decline surgery. In patients with advanced or metastatic disease, radiation therapy can help achieve palliation and symptom control. In comparison, chemotherapy plays a role in the multimodal treatment of locally advanced disease, in adjuvant therapy of resected disease, or as palliative therapy for metastatic disease (30). However, the platinum-based chemotherapy regimens, used as part of the NCCN guidelines, are often poorly tolerated and have significant toxic side effects (31). Examining specific chemotherapy regimens was beyond the scope of this study, and a stage-based analysis of the utility of chemotherapy is necessary to determine which patients should be offered systemic therapies.

This study has several limitations. First, incomplete submission of cancer staging data for all patients diagnosed with NSCLC in Ontario limited our ability to analyze the entire cohort between 2007 and 2015. Second, besides sex, age, stage, histology of tumor, various other factors could be involved in treatment decisions (e.g. comorbidity and patient preference) that could not be captured using administrative datasets. Third, this study is susceptible to coding and

1
2
3 recording error, a limitation inherent to most studies conducted with administrative datasets.

4
5 Finally, this study was conducted within a single-payer healthcare system in Ontario and its
6
7 findings may not be generalizable to private healthcare systems. The treatment effects of
8
9 radiation, chemotherapy, and surgery in patients with lung cancer can be hypothesized to differ
10
11 in multi-payer systems, and the results of this study should therefore be taken in context of a
12
13 universal healthcare system.
14
15

16
17 The strengths of this study include the use of a high-quality population-based dataset and
18
19 long follow-up over an extended period. The ability to report one- and two-, and five-year
20
21 survival rates provides more insight into the mortality and survival of lung cancer patients. Also,
22
23 the use of a flexible parametric model for analysis allows for a more flexible and powerful
24
25 approach in mimicking the actual trends in mortality and survival patterns (17, 32).
26
27

28 29 30 **Conclusions**

31
32 In summary, the present study found that the survival of lung cancer patients in Ontario
33
34 has improved from 2007 to 2015. We observed more than 20% increase in five-year survival
35
36 rates for all surgical groups. This study also indicates that radiation therapy offers significant
37
38 long-term survival advantage over no treatment in elderly patients with lung cancer. Thus,
39
40 radiation therapy plays an important role in both the treatment of patients with unresectable
41
42 disease, and the palliation of patients with lung cancer.
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

Acknowledgement

This study was supported by the Institute for Clinical Evaluative Sciences (ICES), which is funded by an annual grant from the Ontario Ministry of Health and Long-Term Care (MOHLTC). The opinions, results and conclusions reported in this paper are those of the authors and are independent from the funding sources. No endorsement by ICES or the Ontario MOHLTC is intended or should be inferred. Parts of this material are based on data and/or information compiled and provided by CIHI. However, the analyses, conclusions, opinions and statements expressed in the material are those of the authors, and not necessarily those of CIHI. Also, parts of this material are based on data and information provided by Cancer Care Ontario (CCO). The opinions, results, view, and conclusions reported in this paper are those of the authors and do not necessarily reflect those of CCO. No endorsement by CCO is intended or should be inferred. We are indebted to ICES staff at McMaster, specifically, to Richard Perez and Joshua Cerasuolo, for their invaluable help and support.

Disclosure statement

The authors report no conflicts of interest. The authors alone are responsible for the content and writing of the article.

Review Only

References

1. International Agency for Research on Cancer (IARC). World Cancer Report 2014. Lyon, France: IARC; 2014 2014.
2. Canadian Cancer Society's Advisory Committee on Cancer Statistics. Canadian Cancer Statistics 2017. Toronto, ON: Canadian Cancer Society; 2017.
3. Siegel RL, Miller KD, Jemal A. Cancer statistics, 2015. *CA Cancer J Clin*. 2015;65(1):5-29.
4. Walters S, Maringe C, Coleman MP, Peake MD, Butler J, Young N, et al. Lung cancer survival and stage at diagnosis in Australia, Canada, Denmark, Norway, Sweden and the UK: a population-based study, 2004-2007. *Thorax*. 2013;68(6):551-64.
5. Lewin G, Morissette K, Dickinson J, Bell N, Bacchus M, Singh H, et al. Recommendations on screening for lung cancer. *CMAJ*. 2016;188(6):425-32.
6. Molina JR, Yang P, Cassivi SD, Schild SE, Adjei AA. Non-small cell lung cancer: epidemiology, risk factors, treatment, and survivorship. *Mayo Clin Proc*. 2008;83(5):584-94.
7. Canadian Cancer Society. Survival statistics for non-small cell lung cancer: Canadian Cancer Society; 2018 [updated 2018. Available from: <http://www.cancer.ca/en/cancer-information/cancer-type/lung/prognosis-and-survival/non-small-cell-lung-cancer-survival-statistics/?region=on>.
8. Cancer Care Ontario. Ontario Cancer Registry [
9. Parkin M, Samet J, Tyczynski JE, Shields P, Boffetta P, Caporaso N. Lung cancer epidemiology and etiology. In: Travis WD, Brambilla E, Muller-Hermelink HK, Harris CC, editors. *World Health Organization Classification of Tumours: Pathology & Genetics, Tumours of the Lung, Pleura, Thymus, and Heart*. Geneva, Switzerland: WHO Press; 2004.
10. Qin J, Li G, Zhou J. Characteristics of elderly patients with COPD and newly diagnosed lung cancer, and factors associated with treatment decision. *Int J Chron Obstruct Pulmon Dis*. 2016;11:1515-20.
11. Canadian Institute for Health Information. Canadian Classification of Health Intervention Ottawa, Ontario: Canadian Institute for Health Information; 2015 [Available from: https://www.cihi.ca/en/cc_i_volume_three_2015_en.pdf.
12. Statistics Canada. Canadian Classification of Diagnostic, Therapeutic, and Surgical Procedures. Ottawa, Ontario: Statistics Canada; 1986.
13. Booth CM, Li G, Zhang-Salomons J, Mackillop WJ. The impact of socioeconomic status on stage of cancer at diagnosis and survival: a population-based study in Ontario, Canada. *Cancer*. 2010;116(17):4160-7.
14. Ginsburg OM, Fischer HD, Shah BR, Lipscombe L, Fu L, Anderson GM, et al. A population-based study of ethnicity and breast cancer stage at diagnosis in Ontario. *Curr Oncol*. 2015;22(2):97-104.
15. Lambert PC, Royston P. Further development of flexible parametric models for survival analysis. *Stata Journal*. 2009;9(2):265-90.
16. Royston P, Parmar MK. Flexible parametric proportional-hazards and proportional-odds models for censored survival data, with application to prognostic modelling and estimation of treatment effects. *Stat Med*. 2002;21(15):2175-97.

17. Rutherford MJ, Dickman PW, Lambert PC. Comparison of methods for calculating relative survival in population-based studies. *Cancer Epidemiol.* 2012;36(1):16-21.

18. Ettinger DS, Wood DE, Aisner DL, Akerley W, Bauman J, Chirieac LR, et al. Non-Small Cell Lung Cancer, Version 5.2017, NCCN Clinical Practice Guidelines in Oncology. *J Natl Compr Canc Netw.* 2017;15(4):504-35.

19. Little AG, Rusch VW, Bonner JA, Gaspar LE, Green MR, Webb WR, et al. Patterns of surgical care of lung cancer patients. *Ann Thorac Surg.* 2005;80(6):2051-6; discussion 6.

20. Harichand-Herdt S, Ramalingam SS. Gender-associated differences in lung cancer: clinical characteristics and treatment outcomes in women. *Semin Oncol.* 2009;36(6):572-80.

21. Tong BC, Kosinski AS, Burfeind WR, Jr., Onaitis MW, Berry MF, Harpole DH, Jr., et al. Sex differences in early outcomes after lung cancer resection: analysis of the Society of Thoracic Surgeons General Thoracic Database. *J Thorac Cardiovasc Surg.* 2014;148(1):13-8.

22. Senbaklavaci O. Lobar lung resection in elderly patients with non-small cell lung carcinoma: impact of chronic obstructive pulmonary disease on surgical outcome. *Int Surg.* 2014;99(4):319-24.

23. Senbaklavaci O, Taspinar H, Hartert M, Ergun S, Keranen S, Vahl CF. Lobar lung resection in elderly patients with non-small cell lung carcinoma: impact of cardiac comorbidity on surgical outcome. *Swiss Med Wkly.* 2012;142:w13742.

24. Takamochi K, Oh S, Matsuoka J, Suzuki K. Risk factors for morbidity after pulmonary resection for lung cancer in younger and elderly patients. *Interact Cardiovasc Thorac Surg.* 2011;12(5):739-43.

25. Xia W, Yu X, Mao Q, Xia W, Wang A, Dong G, et al. Improvement of survival for non-small cell lung cancer over time. *Onco Targets Ther.* 2017;10:4295-303.

26. Toh CK, Ong WS, Lim WT, Tan DS, Ng QS, Kanavarar R, et al. A Decade of Never-smokers Among Lung Cancer Patients-Increasing Trend and Improved Survival. *Clin Lung Cancer.* 2018.

27. Cufari ME, Proli C, De Sousa P, Raubenheimer H, Al Sahaf M, Chavan H, et al. Increasing frequency of non-smoking lung cancer: Presentation of patients with early disease to a tertiary institution in the UK. *Eur J Cancer.* 2017;84:55-9.

28. Bustamante Alvarez JG, Gonzalez-Cao M, Karachaliou N, Santarpia M, Viteri S, Teixido C, et al. Advances in immunotherapy for treatment of lung cancer. *Cancer Biol Med.* 2015;12(3):209-22.

29. Alexiou C, Beggs D, Onyeaka P, Kotidis K, Ghosh S, Beggs L, et al. Pneumonectomy for stage I (T1N0 and T2N0) nonsmall cell lung cancer has potent, adverse impact on survival. *Annals of Thoracic Surgery.* 2003;76(4):1023-8.

30. Ettinger DS, Aisner DL, Wood DE, Akerley W, Bauman J, Chang JY, et al. NCCN Guidelines Insights: Non-Small Cell Lung Cancer, Version 5.2018. *J Natl Compr Canc Netw.* 2018;16(7):807-21.

31. Hartmann JT, Lipp HP. Toxicity of platinum compounds. *Expert Opin Pharmacother.* 2003;4(6):889-901.

32. Dickman PW, Lambert PC, Coviello E, Rutherford MJ. Estimating net survival in population-based cancer studies. *Int J Cancer.* 2013;133(2):519-21.

Table 1. Distribution of lung cancer patients based on demographics, tumor stage, year of diagnosis and treatment

	No treatment	Radiation only	Chemo. only	Chemorad.	Sub/Lob only	Sub/Lob + RT/CT	Pneum. only	Pneum. + RT/CT	Total
Sex									
Male	6,795 (23.0)	7,995 (27.1)	2,190 (7.4)	6,342 (21.5)	3,292 (11.2)	2,338 (7.9)	163 (0.6)	384 (1.3)	29,499
Female	5,899 (21.9)	6,822 (25.3)	1,858 (6.9)	5,529 (20.5)	3,971 (14.8)	2,474 (9.2)	109 (0.4)	256 (1.0)	26,918
Age group									
< 60 years	1,034 (10.9)	1,837 (19.3)	796 (8.4)	3,355 (35.3)	1,116 (11.7)	1,110 (11.7)	48 (0.5)	207 (2.2)	9,503
60 – 69 years	2,395 (14.6)	3,621 (22.1)	1,285 (7.9)	4,428 (27.1)	2,427 (14.8)	1,869 (11.4)	87 (0.5)	248 (1.5)	16,360
70 – 79 years	4,230 (22.2)	5,419 (28.5)	1,369 (7.2)	3,326 (17.5)	2,838 (14.9)	1,581 (8.3)	113 (0.6)	168 (0.9)	19,044
≥ 80 years	5,035 (43.7)	3,940 (34.2)	598 (5.2)	762 (6.6)	882 (7.7)	252 (2.2)	24 (0.2)	17 (0.2)	11,510
Age: Mean (SD)	75.4 (10.3)	72.2 (10.2)	68.5 (10.5)	65.1 (9.9)	69.1 (9.1)	66.1 (9.0)	68.5 (8.9)	63.5 (9.4)	70.1 (10.7)
Histology									
Carc. ¹	2,536 (20.8)	4,462 (36.5)	882 (7.2)	3,366 (27.5)	387 (3.2)	465 (3.8)	34 (0.3)	92 (0.8)	12,224
SCC ²	1,611 (13.9)	3,464 (29.9)	508 (4.4)	2,646 (22.8)	1,736 (15.0)	1,182 (10.2)	134 (1.2)	302 (2.6)	11,583
AC ³	3,292 (16.2)	5,436 (26.8)	1,992 (9.8)	5,119 (25.2)	2,603 (12.8)	1,687 (8.3)	51 (0.3)	140 (0.7)	20,320
Other	5,255 (42.8)	1,455 (11.8)	666 (5.4)	740 (6.0)	2,537 (20.6)	1,478 (12.0)	53 (0.4)	106 (0.9)	12,290
Stage									
I	1,172 (9.8)	2,652 (22.3)	161 (1.4)	447 (3.8)	5,918 (49.7)	1,451 (12.2)	71 (0.6)	38 (0.3)	11,910
II	512 (11.1)	763 (16.6)	82 (1.8)	410 (8.9)	893 (19.4)	1,609 (35.0)	113 (2.5)	217 (4.7)	4,599
III	2,385 (19.6)	2,838 (23.3)	754 (6.2)	4,128 (33.8)	318 (2.6)	1,364 (11.2)	74 (0.6)	337 (2.8)	12,198
IV	8,625 (31.1)	8,564 (30.9)	3,051 (11.0)	6,886 (24.9)	134 (0.5)	388 (1.4)	14 (0.1)	48 (0.2)	27,710

Table 1. (Continued)

	No treatment	Radiation only	Chemo. only	Chemorad.	Sub/Lob only	Sub/Lob + RT/CT	Pneum. only	Pneum. + RT/CT	Total
Year									
2007	986 (19.5)	1,133 (22.4)	456 (9.0)	1,230 (24.3)	660 (13.0)	474 (9.4)	39 (0.8)	85 (1.7)	5,063
2008	1,129 (20.9)	1,369 (25.3)	438 (8.1)	1,213 (22.4)	690 (12.7)	454 (8.4)	44 (0.8)	77 (1.4)	5,414
2009	1,357 (23.4)	1,474 (25.4)	433 (7.5)	1,239 (21.3)	687 (11.8)	515 (8.9)	30 (0.5)	72 (1.2)	5,807
2010	1,668 (25.2)	1,723 (26.0)	465 (7.0)	1,391 (21.0)	794 (12.0)	485 (7.3)	28 (0.4)	78 (1.2)	6,632
2011	1,557 (24.2)	1,798 (27.9)	399 (6.2)	1,335 (20.7)	741 (11.5)	520 (8.1)	27 (0.4)	67 (1.0)	6,444
2012	1,763 (26.2)	1,684 (25.0)	396 (5.9)	1,314 (19.5)	857 (12.7)	601 (8.9)	32 (0.5)	78 (1.2)	6,725
2013	1,762 (25.1)	1,873 (26.7)	468 (6.7)	1,365 (19.4)	876 (12.5)	583 (8.3)	29 (0.4)	66 (0.9)	7,022
2014	1,433 (20.7)	1,930 (27.9)	498 (7.2)	1,410 (20.3)	997 (14.8)	569 (8.2)	26 (0.4)	68 (1.0)	6,931
2015	1,039 (16.3)	1,833 (28.7)	495 (7.8)	1,374 (21.5)	961 (15.1)	611 (9.6)	17 (0.3)	49 (0.8)	6,379
Total	12,694 (22.5)	14,817 (26.3)	4,048 (7.2)	11,871 (21.0)	7,263 (12.9)	4,812 (8.5)	272 (0.5)	640 (1.1)	56,417

1- Carcinoma, NOS 2- Squamous cell carcinoma 3- Adenocarcinoma

Table 2. Predictors of surgical resection among patients diagnosed with non-small cell lung cancer

Variable	OR (95% CI)
Sex	
Male	Referent
Female	1.10 (1.05, 1.17) ⁺⁺
Age group	
< 60 years	Referent
60 – 69 years	0.81 (0.75, 0.88) ⁺⁺
70 – 79 years	0.48 (0.44, 0.52) ⁺⁺
≥ 80 years	0.12 (0.11, 0.14) ⁺⁺
Year of diagnosis	0.95 (0.94, 0.96) ⁺⁺
Histology	
Carcinoma, NOS	Referent
SCC ¹	3.26 (2.97, 3.58) ⁺⁺
AC ²	3.40 (3.12, 3.73) ⁺⁺
Other	5.88 (5.35, 6.47) ⁺⁺
Stage	
I	Referent
II	0.97 (0.90, 1.05)
III	0.11 (0.10, 0.11) ⁺⁺
IV	0.01 (0.01, 0.01) ⁺⁺

+ p < 0.05, ++ p < 0.001, 1- Squamous cell carcinoma, 2- Adenocarcinoma

Table 3. Five-year survival rate (95% CI) based on histology, stage, and treatment modality

Histology	Stage	No treatment	RT only	CT only	Chemorad.	Sub/Lob only	Sub/Lob + RT/CT	Pneum. only	Pneum. + RT/CT
Carc. ¹	I	.17 (.16, .19)	.35 (.34, .37)	.07 (.05, .10)	.19 (.16, .21)	.75 (.74, .76)	.58 (.56, .60)	.55 (.43, .68)	.44 (.27, .61)
	II	.09 (.07, .10)	.16 (.15, .17)	.05 (.02, .08)	.18 (.16, .20)	.53 (.50, .57)	.57 (.56, .59)	.29 (.24, .34)	.54 (.50, .58)
	III	.04 (.04, .04)	.07 (.07, .07)	.04 (.03, .05)	.16 (.15, .16)	.40 (.34, .46)	.43 (.42, .44)	.18 (.11, .25)	.35 (.32, .37)
	IV	.02 (.02, .02)	.01 (.01, .01)	.04 (.04, .04)	.03 (.03, .04)	.25 (.20, .30)	.19 (.18, .21)	**	.16 (.12, .19)
SCC ²	I	.19 (.18, .21)	.38 (.37, .39)	.09 (.06, .12)	.17 (.15, .19)	.73 (.73, .74)	.55 (.54, .56)	.50 (.46, .55)	.32 (.24, .40)
	II	.08 (.08, .09)	.16 (.15, .17)	.06 (.04, .09)	.19 (.18, .21)	.53 (.51, .54)	.54 (.53, .54)	.31 (.27, .34)	.51 (.49, .52)
	III	.05 (.04, .05)	.07 (.07, .07)	.05 (.04, .05)	.16 (.15, .16)	.36 (.34, .39)	.40 (.40, .41)	.23 (.20, .26)	.35 (.33, .36)
	IV	.02 (.02, .02)	.01 (.01, .01)	.04 (.04, .05)	.03 (.03, .03)	.28 (.24, .32)	.15 (.14, .16)	**	.13 (.11, .15)
AC ³	I	.27 (.25, .28)	.44 (.43, .45)	.17 (.14, .20)	.28 (.26, .30)	.79 (.79, .79)	.63 (.62, .64)	.57 (.51, .63)	.53 (.45, .62)
	II	.13 (.12, .15)	.24 (.22, .25)	.10 (.07, .14)	.26 (.24, .28)	.59 (.58, .61)	.60 (.60, .61)	.36 (.29, .44)	.59 (.56, .63)
	III	.07 (.07, .07)	.11 (.11, .12)	.07 (.06, .07)	.23 (.23, .24)	.43 (.41, .46)	.49 (.48, .50)	.27 (.19, .36)	.41 (.39, .44)
	IV	.04 (.04, .04)	.02 (.02, .02)	.08 (.07, .08)	.06 (.06, .06)	.33 (.31, .35)	.21 (.20, .23)	.08 (.00, .17)	.22 (.20, .26)
Other	I	.22 (.21, .23)	.45 (.44, .47)	.11 (.07, .14)	.24 (.20, .29)	.81 (.80, .81)	.64 (.63, .65)	.62 (.54, .69)	.50 (.38, .62)
	II	.10 (.09, .11)	.20 (.17, .22)	.08 (.05, .10)	.27 (.22, .32)	.59 (.57, .60)	.62 (.61, .63)	.42 (.35, .48)	.64 (.60, .68)
	III	.05 (.05, .05)	.10 (.09, .11)	.05 (.04, .06)	.23 (.21, .24)	.46 (.44, .49)	.51 (.50, .52)	.28 (.23, .34)	.43 (.41, .46)
	IV	.03 (.03, .03)	.02 (.02, .02)	.06 (.06, .06)	.06 (.06, .07)	.31 (.27, .34)	.24 (.23, .25)	.10 (.03, .16)	.23 (.20, .26)

1- Carcinoma, NOS, 2-Squamous cell carcinoma, 3- Adenocarcinoma

Table A.1 Trends of radiation in elderly patients (80+) over the study period (p-for-trend < 0.001)

Year of diagnosis	No treatment	Radiation only	Chemotherapy only	Chemoradiation	Surgery groups	Total
2007	333 (38.9)	278 (32.5)	68 (7.9)	64 (7.5)	113 (13.2)	856
2008	426 (43.3)	314 (31.9)	60 (6.1)	74 (7.5)	110 (11.2)	984
2009	480 (45.3)	363 (34.3)	58 (5.5)	60 (5.7)	99 (9.3)	1,060
2010	629 (48.3)	405 (31.1)	53 (4.1)	86 (6.6)	129 (9.9)	1,302
2011	622 (45.8)	475 (35.0)	63 (4.6)	82 (6.0)	117 (8.6)	1,359
2012	767 (50.9)	448 (29.8)	60 (4.0)	88 (5.8)	143 (9.5)	1,506
2013	762 (47.1)	547 (33.8)	80 (4.9)	94 (5.8)	135 (8.3)	1,618
2014	617 (40.8)	567 (37.5)	82 (5.4)	98 (6.5)	149 (9.9)	1,513
2015	399 (30.4)	543 (41.4)	74 (5.6)	116 (8.8)	180 (13.7)	1,312
Total	5,035 (43.7)	3,940 (34.2)	598 (5.2)	762 (6.6)	1,175 (10.2)	11,510

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

Figure Legends

Figure 1. Trends in one-year, two-year, and five-year survival rates based on treatment modality

The graph is based on the final model which included sex, age group, treatment modality, stage of disease, histology of tumor, and year of diagnosis. The model also included interaction terms between age group and stage, age group and histology, treatment modality and year of diagnosis, treatment modality and stage, and stage and year of diagnosis. In addition, age group, stage of tumor, treatment modality, histology of tumor, and year of diagnosis were included as time dependent variables.

Figure 2. Trends in five-year survival rate based on sex and age group

The graph is based on the final model which included sex, age group, treatment modality, stage of disease, histology of tumor, and year of diagnosis. The model also included interaction terms between age group and stage, age group and histology, treatment modality and year of diagnosis, treatment modality and stage, and stage and year of diagnosis. In addition, age group, stage of tumor, treatment modality, histology of tumor, and year of diagnosis were included as time dependent variables.

Figure 3. Trends in one-year, two-year, and five-year survival rates based on stage of tumor

The graph is based on the final model which included sex, age group, treatment modality, stage of disease, histology of tumor, and year of diagnosis. The model also included interaction terms between age group and stage, age group and histology, treatment modality and year of diagnosis, treatment modality and stage, and stage and year of diagnosis. In addition, age group, stage of

tumor, treatment modality, histology of tumor, and year of diagnosis were included as time dependent variables.

Figure A.1. Adjusted Kaplan-Meier survival curves based on the preliminary treatment classification

This graph was adjusted based on age group, stage of disease, and histology. This was used to combine sublobar only and lobectomy only groups into one group (Sub/Lob only). Also, we combined sublobar with radiation or chemotherapy and lobectomy with radiation or chemotherapy into one group (Sub/Lob + RT/CT).




Figure 1. Trends in one-year, two-year, and five-year survival rates based on treatment modality

157x114mm (150 x 150 DPI)



Figure 2. Trends in five-year survival rate based on sex and age group

157x114mm (150 x 150 DPI)




Figure 3. Trends in one-year, two-year, and five-year survival rates based on stage of tumor

157x114mm (150 x 150 DPI)




Figure A.1. Adjusted Kaplan-Meier survival curves based on the preliminary treatment classification

157x114mm (150 x 150 DPI)