

# I. Markov chains

A Markov chain is a discrete-time stochastic process  $\Theta_1, \Theta_2, \dots, \Theta_t, \dots$  that satisfies the Markov property

$$p(\Theta_t | \Theta_{t-1}, \dots, \Theta_1) = p(\Theta_t | \Theta_{t-1}) \quad \forall t.$$

↓  
states  
of the  
chain



A Markov chain is fully defined by

- $\pi_0(\Theta)$ , the initial distribution.
- $T(\Theta_{t-1}, \Theta_t) = p(\Theta_t | \Theta_{t-1})$ , the transition kernel.

## Basic limit theorem

$$\pi_n(\theta) = \int \pi_{n-1}(\theta') T(\theta'; \theta) d\theta'$$

Let  $\pi_n = \pi_0 \underbrace{TT \dots T}_{n \text{ times}}$  be the probability over the states after  $n$  iterations.

Let  $\pi_*(\theta)$  be the stationary distribution of the Markov chain with kernel  $T$ , i.e. such that  $\pi_* T = \pi_*$ .

Then  $\lim_{n \rightarrow \infty} \pi_n(\theta) = \pi_*(\theta) \quad \forall \theta$

$\leadsto$  " $\pi_n$  converges to  $\pi_*$ "

Assumptions:

- $\pi_*$  exists
- The chain is irreducible, i.e. if any state can be reached from any other state with positive probability in a finite number of steps.

• The chain is aperiodic.

A state  $\theta$  is periodic with period  $k$  if the number of steps to return to  $\theta$  is always divisible by  $k \geq 2$ .

A Markov chain is aperiodic if none of its states is periodic with  $k \geq 2$ .

## Time reversibility

A Markov chain is time reversible if

$$(\theta_0, \theta_1, \dots, \theta_n) \stackrel{D}{=} (\theta_n, \theta_{n-1}, \dots, \theta_0)$$

↘ "equal in distribution"

$$\Rightarrow (\theta_0, \theta_1) \stackrel{D}{=} (\theta_1, \theta_0)$$

$$\text{and } (\theta_0) \stackrel{D}{=} (\theta_1)$$



$$\pi_0 = \pi_1$$

$$= \pi_0 T$$

$\Rightarrow \pi_0$  is the stationary distribution

The number of states moving forward is equal in distribution to the number of states moving backward.

$$\text{Also } (\theta_0, \theta_1) \stackrel{D}{=} (\theta_1, \theta_0)$$

$$\Leftrightarrow P(\theta_0 = i, \theta_1 = j) = P(\theta_1 = i, \theta_0 = j) \quad \forall i, j$$

$$\begin{aligned} \Leftrightarrow P(\theta_0 = i) P(\theta_1 = j | \theta_0 = i) \\ = P(\theta_0 = j) P(\theta_1 = i | \theta_0 = j) \end{aligned}$$

$$\Leftrightarrow \pi_0(i) T(i, j) = \pi_0(j) T(j, i)$$

Local balanced equations

If the local balanced equation hold for  $\pi$  and  $T$ , then  $\pi$  is the stationary distribution governed by  $T$ .

## II. Metropolis - Hastings (MH)

MH sampling works by driving a Markov chain  $\theta_1, \theta_2, \dots$  whose stationary distribution is the target  $\pi(\theta)$ .

①  $\theta' \sim q(\theta' | \theta_t)$

Annotations:  
-  $\theta'$ : candidate  
-  $q(\theta' | \theta_t)$ : proposal distribution  
-  $\theta_t$ : current state

②  $\alpha(\theta' | \theta_t) = \min \left\{ \frac{\pi(\theta')}{\pi(\theta_t)} \frac{q(\theta_t | \theta')}{q(\theta' | \theta_t)}, 1 \right\}$

↳ acceptance ratio

③  $\theta_{t+1} = \begin{cases} \theta' & \text{with probability } \alpha(\theta' | \theta_t) \\ \theta_t & \text{otherwise} \end{cases}$

↳  
 $u \sim U[0, 1]$   
if  $u < \alpha$ , accept

⚠ MH only requires the ratio of the target density  $\frac{\pi(\theta')}{\pi(\theta_t)}$

$\Rightarrow$  We can use an unnormalized density  $\pi^*(\theta) = \frac{1}{2} \pi(\theta)$  with an unknown normalizer  $\frac{1}{2}$ .

Random walk MH (= "Metropolis algorithm")

$\theta' = \theta_t + \epsilon$   $\rightarrow$  from a symmetric and centered distribution  $g$  (e.g.  $N$ )

In this case,  $q(\theta' | \theta_t) = g(\epsilon)$   
 $= \begin{cases} q(\theta_t | \theta') = g(-\epsilon) = g(\epsilon) \end{cases}$

$$\alpha(\theta' | \theta_t) = \min \left\{ \frac{\pi(\theta')}{\pi(\theta_t)} \frac{\cancel{q(\theta_t | \theta')}}{\cancel{q(\theta' | \theta_t)}}, 1 \right\}$$

Why does it work?

Thm: The MH acceptance probability  $\alpha(\theta'|\theta_t)$  for the proposal  $q(\theta'|\theta_t)$  and target  $\pi(\theta)$  defines a Markov chain with the transition kernel

$$T(\theta_t, \theta') = \alpha(\theta'|\theta_t) q(\theta'|\theta_t).$$


This chain satisfies the detailed balanced condition.

Proof:

$$\begin{aligned}\pi(\theta_t) T(\theta_t, \theta') &= \pi(\theta_t) \alpha(\theta'|\theta_t) q(\theta'|\theta_t) \\&= \min \{ \pi(\theta') q(\theta_t|\theta'), \pi(\theta_t) q(\theta'|\theta_t) \} \\&= \min \{ \pi(\theta_t) q(\theta'|\theta_t), \pi(\theta') q(\theta_t|\theta') \} \\&= \pi(\theta') \alpha(\theta_t|\theta') q(\theta_t|\theta') \\&= \pi(\theta_t) T(\theta'|\theta_t)\end{aligned}\quad \square$$

## III. Bayesian inference with MCMC

$$p(\theta|x) = \frac{p(x|\theta) p(\theta)}{p(x)}$$



$$= \int p(x|\theta) p(\theta) d\theta$$

Difficult to evaluate!

$$= \frac{1}{Z(x)} p(x|\theta) p(\theta) = \pi^*(\theta)$$

$\Rightarrow$  Use MCMC for posterior sampling

$$\log \pi^*(\theta) = \log p(x|\theta) + \log p(\theta) + C$$

Diagnostics:

 Gelman-Rubin diagnostic

- \* Convergence checking based on  $m$  chains from widely diverse starting points.



Assuming  $\theta_{ij}$  ( $i=1 \dots m$ ,  $j=1 \dots m$ )

- Between-chain variance

$$B = \frac{n}{m-1} \sum_{j=1}^m (\theta_{:j} - \theta_{::})^2 \quad (\text{variance of the mean})$$

$$\text{where } \theta_{:j} = \frac{1}{n} \sum_{i=1}^n \theta_{ij}$$

$$\theta_{::} = \frac{1}{m} \sum_{j=1}^m \theta_{:j}$$

- Within-chain variance

$$W = \frac{1}{m} \sum_{j=1}^m s_j^2 \quad (\text{mean of chain variances})$$

$$\text{where } s_j^2 = \frac{1}{n-1} \sum_{i=1}^n (\theta_{ij} - \theta_{:j})^2$$

$$\Rightarrow V[\theta] = \frac{n-1}{n} W + \frac{1}{n} B$$

$$\Rightarrow \hat{R} = \sqrt{\frac{V[\theta]}{W}} \geq 1 \quad \begin{array}{l} \text{overestimates } \pi' \text{'s variance} \\ \hat{R} \text{ approaches } 1 \\ \text{as } n \text{ increases} \end{array}$$

underestimate  $\pi'$ 's variance  $\leftarrow$

$\rightarrow$  Stop when  $\hat{R} < 1.1$

\* Effective sample size

$$\rho_k = \frac{\mathbb{E}[(\theta_i - \mu)(\theta_{i+k} - \mu)]}{\sigma^2}$$

over  
the time  
index  $i$

$$ESS = \frac{M}{1 + 2 \sum_{k=1}^{\infty} \rho_k} \quad \left. \vphantom{\sum_{k=1}^{\infty} \rho_k} \right\} = T \text{ autocorrelation}$$

$\Rightarrow$  Uncorrelated samples  $\rightarrow \rho_k = 0$

and  $ESS = M$

$\Rightarrow$  Correlated samples  $\rightarrow$  small  $ESS$