

## On The Semantic Position of properties, primitive particles and semantic structures


D. Gueorguiev 12/23/2021

Each property has inherent [semantic signature](#) which describes its semantic meaning in isolation, *in situ*. It also serves as its *initial semantic position* in [Semantic Space](#). Each property also has specific semantic mass based on which it is determined its path from the root in the particle property tree. The semantic mass of a property is determined based on the semantic information content a property conveys:

*Property Mass*  $\sim$  *Property Valence*  $\times$  *Information Content* or in symbol notation

$M_p = |V_p| \times IC$  where  $|V_p|$  is the valence and  $IC$  is the information content of the property. Certain properties have the affinity to bind to multiple child properties which reveal additional details for the semantic information provided by the parent. The more child properties a parent property can bind to - the higher will be its property valence.

Let us assume we have two properties  $P_1$  and  $P_2$  which are attracted by a binding force  $F_b(P_1, P_2)$ . The masses of each of the properties are  $M_{P_1}$  and  $M_{P_2}$ .



On the Figure above are depicted the two properties positioned in the  $N$ -dimensional semantic space *in situ* with grey color. Each of the properties is represented by blue oval where the oval with larger diameter corresponds to the larger semantic mass, in this case  $P_1$ . After the binding force  $F_b$  is applied to those *in-situ* properties they move to a bound state as depicted in the Figure. Note that in their bound state the semantic positions of the two properties are still  $\varepsilon(P_1, P_2)$  distance apart. Generally, the semantic distance  $\varepsilon$  setting apart any two bound properties will depend on the binding force between the properties which in turn depends on the presence of specific regions in each of the properties. For details follow the discussion in [Note on Binding of Association Property to Semantic Properties](#). The parameters  $d_1$  and  $d_2$  represent the semantic distances which each of the properties  $P_1$  and  $P_2$  has travelled accordingly until reaching bound state. The semantic distances are inversely proportional to the ratio of the semantic masses for each property:

$$\frac{d_1}{d_2} \sim \left( \frac{M_{P_2}}{M_{P_1}} \right)^\alpha \text{ where } \alpha > 0$$

//TODO: finish the description