SYMMETRIES, SPACETIME, AND RELATIVITY

Abstract. These notes embark on an exploration of the geometric and algebraic foundations underpinning special and general relativity. Starting with the fundamental concepts of symmetry groups and their associated algebras, we build from basic spatial rotations to the full symmetries of flat spacetime and their implications for particle properties. The journey then progresses to the curved spacetime of general relativity, aiming to elucidate key physical phenomena such as the anomalous precession of Mercury's orbit, the relativistic intricacies of satellite timekeeping, and the fascinating physics of rotating black holes. The treatment seeks to bridge abstract formalism with physical insight, providing a foundation for readers new to manifolds, tangent space algebras, and the geometric description of gravity.

1. Introduction

The goal of these notes is to develop a clear and mathematically grounded understanding of the principles underlying classical mechanics, special relativity, and general relativity. We aim to build the formalism necessary to explore key physical phenomena, such as the precession of Mercury's orbit, the intricacies of satellite timekeeping, and the nature of spinning black holes.

We begin in Section 2 by introducing continuous symmetries and their algebraic structure using the familiar example of spatial rotations and the classical symmetry group of spacetime. Section 3 then transitions to the fabric of flat spacetime and the symmetries of special relativity, exploring the homogeneous isometry group of flat spacetime, its algebra, and the fundamental kinematic consequences. The full symmetry group of flat spacetime and its role in defining particle properties like mass and spin through its Casimir invariants, is discussed in Section 4.

Following this, Section 5 discusses the conceptual leap from special to general relativity, motivating the idea of gravity as spacetime curvature. The mathematical language of curved spacetime, differential geometry, is developed in Section 6. This leads to Einstein's field equations in Section 7, which describe how matter and energy dictate this curvature.

With these tools, Section 8 examines the spacetime geometry around a non-rotating, uncharged spherical mass, applying this to understand phenomena like the perihelion precession of Mercury, relativistic effects on clocks, and gravitational lensing. More complex scenarios involving rotating and charged massive objects, including the concept of an ergosphere, are explored in Section 9. Finally, Section 10 briefly touches upon further striking phenomena and future directions in the study of spacetime and gravity.

Throughout, the emphasis is on building intuition and formalism simultaneously, focusing on precise mathematical definitions and constructive examples. The intended audience is readers with some background in calculus and linear algebra, but no prior exposure to group theory or differential geometry is assumed.

Date: September 11, 2025.

2. Continuous Symmetries and Their Algebraic Structure

In physics and mathematics, continuous symmetry groups describe transformations that preserve certain structures. These groups are smooth spaces with operations that vary smoothly. The algebra associated with such a group is the tangent space at the identity element, equipped with a bracket operation that encodes the infinitesimal structure of the group.

2.1. **Foundational Example: The Group of Spatial Rotations ($SO(3)$).** A key example is the group of rotations in three dimensions, consisting of all orthogonal 3×3 matrices with determinant 1. These rotations preserve lengths and orientations in three-dimensional space. We begin with the group of rotations in three dimensions and its algebra, which provides an accessible example to understand generators, commutation relations, and matrix exponentiation. This lays the groundwork for the more complex groups appearing in relativity.

Definition 2.1. The *orthogonal group* is the group that preserves the identity tensor δ under conjugation:

$$\mathrm{O}(n) := \left\{ R \in GL(n, \mathbb{R}) \mid R^{\mu}{}_{\alpha} \delta_{\mu\nu} R^{\nu}{}_{\beta} = \delta_{\alpha\beta} \right\}.$$

This group contains both rotations, reflections, and improper rotations (i.e., compositions of a rotation and a reflection). To restrict to only proper rotations, the *special orthogonal group* is the subgroup that contains only matrices with determinant $+1$

$$\mathrm{SO}(n) = \{ R \in \mathrm{O}(n) \mid \det(R) = +1 \}.$$

Lemma 2.2. *Rotation matrices in $SO(2)$ can be written as*

$$R = \begin{pmatrix} \cos\theta & \sin\theta \\ -\sin\theta & \cos\theta \end{pmatrix}$$

Proof. Let $R \in \mathrm{SO}(2)$ such that

$$R = \begin{pmatrix} a & b \\ c & d \end{pmatrix}$$

Orthogonality $R^{\mu}{}_{\alpha} \delta_{\mu\nu} R^{\nu}{}_{\beta} = \delta_{\alpha\beta}$ gives

$$\begin{pmatrix} a & b \\ c & d \end{pmatrix} \begin{pmatrix} 1 & 0 \\ 0 & 1 \end{pmatrix} \begin{pmatrix} a & c \\ b & d \end{pmatrix} = \begin{pmatrix} a^2 + b^2 & ac + bd \\ ac + bd & c^2 + d^2 \end{pmatrix} = \begin{pmatrix} 1 & 0 \\ 0 & 1 \end{pmatrix},$$

which gives the following three equations,

$$a^2 + b^2 = 1,$$
$$c^2 + d^2 = 1,$$
$$ac + bd = 0,$$

The restriction that $\det(R) = 1$ gives

$$ad - bc = 1.$$

Since the first two equations are equations for the unit circle, the first equation can be parameterized with an arbitrary angle $\theta \in [0, 2\pi)$ such that $a = \cos\theta$ and $b = \sin\theta$. The second equation can be parameterized by a second angle $\phi \in [0, 2\pi)$ such that $c = \sin\phi$ and $d = \cos\phi$. The determinant equation becomes

$$\cos\theta\cos\phi - \sin\theta\sin\phi = 1,$$

which is the cosine of the sum of two angles

$$\cos\theta\cos\phi - \sin\theta\sin\phi = \cos(\theta+\phi) = 1.$$

This implies that $\theta + \phi = 0 \pmod{2\pi}$ or $\phi = -\theta$. Substituting into the definitions of c and d:

$$c = \sin(-\theta) = -\sin\theta, \text{ and}$$
$$d = \cos(-\theta) = \cos\theta,$$

which gives the rotation matrix

$$R = \begin{pmatrix} \cos\theta & \sin\theta \\ -\sin\theta & \cos\theta \end{pmatrix}.$$

□

Definition 2.3. The *generalized orthogonal group* is the group that preserves some metric g:

$$\mathrm{O}(n,\mathbb{F}) := \left\{ R \in GL(n,\mathbb{F}) \mid R^{\mu}{}_{\alpha} g_{\mu\nu} R^{\nu}{}_{\beta} = g_{\alpha\beta} \right\},$$

where $GL(n,\mathbb{F})$ is the general linear group of order n over a field $\mathbb{F}$, and g is a symmetric, non-degenerate bilinear form. In other words, the (0,2)-tensor g has the following properties for all vectors u, v, w and scalar λ:

- Bilinearity:

$$g_{\mu\nu}\left(\lambda u^{\mu} + v^{\mu}\right) w^{\nu} = \lambda g_{\mu\nu} u^{\mu} w^{\nu} + g_{\mu\nu} v^{\mu} w^{\nu}$$
$$= g_{\mu\nu} w^{\nu}\left(\lambda u^{\mu} + v^{\mu}\right) = g_{\mu\nu} v^{\mu} w^{\nu} + \lambda g_{\mu\nu} u^{\mu} w^{\nu},$$

- Symmetry: $g_{\mu\nu} = g_{\nu\mu}$,
- Non-degeneracy: $g_{\mu\nu} u^{\mu} = 0$ implies that $u^{\mu} = 0$.

Theorem 2.4. *The associated tangent algebra for an orthogonal group consists of matrices skew with respect to g*

$$\mathfrak{o}(n,\mathbb{F}) = \left\{ X \in M_n(\mathbb{F}) \mid X_{\alpha\beta} + X_{\beta\alpha} = 0 \right\}.$$

Proof. Consider a smooth curve $R(t) : \mathbb{R} \to \mathrm{O}(n,\mathbb{F})$ with $R^{\mu}{}_{\nu}(0) = \delta^{\mu}{}_{\nu}$. Since $R(t)$ is orthogonal,

$$R^{\mu}{}_{\alpha} g_{\mu\nu} R^{\nu}{}_{\beta} = g_{\alpha\beta}.$$

Differentiating,

$$\frac{d}{dt}\left[R^{\mu}{}_{\alpha} g_{\mu\nu} R^{\nu}{}_{\beta}\right] = \frac{d}{dt}\left[g_{\alpha\beta}\right].$$

Using the product rule:

$$\dot{R}^{\mu}{}_{\alpha} g_{\mu\nu} R^{\nu}{}_{\beta} + R^{\mu}{}_{\alpha} \dot{g}_{\mu\nu} R^{\nu}{}_{\beta} + R^{\mu}{}_{\alpha} g_{\mu\nu} \dot{R}^{\nu}{}_{\beta} = \dot{g}_{\alpha\beta}.$$

At $t = 0$, $R^{\mu}{}_{\nu} = \delta^{\mu}{}_{\nu}$

$$\dot{R}^{\mu}{}_{\alpha} g_{\mu\nu} \delta^{\nu}{}_{\beta} + \delta^{\mu}{}_{\alpha} \dot{g}_{\mu\nu} \delta^{\nu}{}_{\beta} + \delta^{\mu}{}_{\alpha} g_{\mu\nu} \dot{R}^{\nu}{}_{\beta} = \dot{g}_{\alpha\beta}$$
$$\dot{R}^{\mu}{}_{\alpha} g_{\mu\beta} + \dot{g}_{\alpha\beta} + g_{\alpha\nu} \dot{R}^{\nu}{}_{\beta} = \dot{g}_{\alpha\beta}.$$

If we define, $X^{\mu}{}_{\nu} := \dot{R}^{\mu}{}_{\nu}$

$$X^{\mu}{}_{\alpha} g_{\mu\beta} + g_{\alpha\nu} X^{\nu}{}_{\beta} = 0.$$

Lowering the index of the first term and the metric in the second tensor lowers the index.

$$\begin{aligned} g^{\mu\rho} X_{\rho\alpha} g_{\mu\beta} + X_{\alpha\beta} &= 0 \\ \delta^{\rho}{}_{\beta} X_{\rho\alpha} + X_{\alpha\beta} &= 0 \\ X_{\beta\alpha} + X_{\alpha\beta} &= 0, \end{aligned}$$

which means that X is skew-symmetric with respect to g. □

Definition 2.5. The basis elements of tangent space algebra $\mathfrak{so}(3)$ are

$$J_1 = \begin{pmatrix} 0 & 0 & 0 \\ 0 & 0 & -1 \\ 0 & 1 & 0 \end{pmatrix}, \quad J_2 = \begin{pmatrix} 0 & 0 & 1 \\ 0 & 0 & 0 \\ -1 & 0 & 0 \end{pmatrix}, \quad J_3 = \begin{pmatrix} 0 & -1 & 0 \\ 1 & 0 & 0 \\ 0 & 0 & 0 \end{pmatrix}.$$

Theorem 2.6. *These satisfy the commutation relations:*

$$[J_i, J_j] = \epsilon_{ijk} J_k,$$

where ϵ_{ijk} is the fully antisymmetric symbol with

$$\epsilon_{ijk} = \begin{cases} +1 & \text{if } (i,j,k) \text{ is an even permutation of } (1,2,3), \\ -1 & \text{if } (i,j,k) \text{ is an odd permutation of } (1,2,3), \\ 0 & \text{if any two indices are equal.} \end{cases}$$

Proof. This can be proven explicitly by multiplying each pair of generators and computing their commutators. For example,

$$[J_1, J_2] := J_1 J_2 - J_2 J_1 = J_3,$$

and similarly for the other pairs, confirming the relation

$$[J_i, J_j] = \epsilon_{ijk} J_k.$$

□

Theorem 2.7. *The image of the exponential map of $\mathfrak{so}(3)$ is contained in* $\mathrm{SO}(3)$*:*

$$\forall X \in \mathfrak{so}(3) : \ \exp(X) \in \mathrm{SO}(3).$$

Proof. Let $R = \exp(X)$ for some $X \in \mathfrak{so}(3)$. Since $\mathfrak{so}(3)$ consists of skew-symmetric matrices, we have $X^\top = -X$. Using the identity $(\exp(X))^\top = \exp(X^\top)$, it follows:

$$R^\top R = \exp(X^\top) \exp(X) = \exp(-X) \exp(X).$$

Since X commutes with $-X$ (as $[X, -X] = 0$ for all matrices), we may combine the exponents:

$$R^\top R = \exp(-X + X) = \exp(0) = I,$$

so R is orthogonal. Next, we use the identity $\det(\exp(X)) = \exp(\mathrm{tr}(X))$, valid for all square matrices X. Since every $X \in \mathfrak{so}(3)$ is skew-symmetric, we have $\mathrm{tr}(X) = 0$, so:

$$\det(\exp(X)) = \exp(\mathrm{tr}(X)) = \exp(0) = 1.$$

Therefore, $R = \exp(X)$ is a special orthogonal matrix: $\exp(X) \in \mathrm{SO}(3)$. □

2.2. **Broader Example: The Classical Symmetry Group of Spacetime.**

Definition 2.8. The *classical symmetry group*, denoted CSG(3), is the group associated with classical physics. It acts on $\mathbb{R}^4$ of the form:

$$(t, \vec{x}) \mapsto (t + \tau, R\vec{x} + \vec{v}t + \vec{a}),$$

where $R \in \mathrm{SO}(3)$ is a rotation described in the previous subsection, $\vec{v} \in \mathbb{R}^3$ is a velocity, $\vec{a} \in \mathbb{R}^3$ is a spatial translation, and $\tau \in \mathbb{R}$ is a time translation.

Definition 2.9. If we use an augmented spacetime 5-vector in homogeneous coordinates, $(x^0, x^1, x^2, x^3, x^4) \equiv (t, x, y, z, h)$ where $h := 1$ is the homogeneous coordinate, we can define the classical symmetry algebra $\mathfrak{csg}(3)$ in terms of its generators. The generators are defined by their non-zero matrix elements $(M)_{rc}$ (row r, column c, 0-indexed) as follows:

- Rotation generators:
  - J_1: $(J_1)_{23} = -1, (J_1)_{32} = 1$
  - J_2: $(J_2)_{13} = 1, (J_2)_{31} = -1$
  - J_3: $(J_3)_{12} = -1, (J_3)_{21} = 1$
- Boost generators:
  - K_1: $(K_1)_{01} = 1$
  - K_2: $(K_2)_{02} = 1$
  - K_3: $(K_3)_{03} = 1$
- Spatial translation generators:
  - P_1: $(P_1)_{14} = 1$
  - P_2: $(P_2)_{24} = 1$
  - P_3: $(P_3)_{34} = 1$
- Time translation generator:
  - H: $(H)_{04} = 1$

All other matrix elements are zero for each generator.

Theorem 2.10 (Commutator relations of the classical symmetry algebra $\mathfrak{csg}(3)$)**.** *The generators J_i (rotations), K_i (boosts), P_i (spatial translations), and H (time translation) satisfy the following commutation relations:*

$$[J_i, J_j] = \epsilon_{ijk} J_k, \quad [J_i, K_j] = \epsilon_{ijk} K_k, \quad [J_i, P_j] = \epsilon_{ijk} P_k,$$
$$[K_i, K_j] = 0, \quad [P_i, P_j] = 0, \quad [K_i, P_j] = 0,$$
$$[H, J_i] = 0, \quad [H, P_i] = 0, \quad [H, K_i] = P_i,$$

where ϵ_{ijk} is the fully antisymmetric symbol.

Proof. Similar to the SO(3) relations found in theorem 2.6, this can be proven explicitly by multiplying each pair of generators and computing their commutators. For example,

$$[J_1, J_2] := J_1 J_2 - J_2 J_1 = J_3,$$

The other pairs can be easily checked the same way. □

3. The Fabric of Flat Spacetime and its Symmetries

3.1. **Redefining Space and Time.** The theory of *special relativity* revolutionized our understanding of space and time. It is built upon two fundamental postulates:

(1) **The Principle of Relativity:** The laws of physics take the same form in all inertial frames of reference. This means there is no "absolute" rest frame; any inertial frame is equally valid for describing physical phenomena.
(2) **The Principle of Invariant Light Speed:** The speed of light in a vacuum, denoted c, has the same value for all inertial observers, regardless of the motion of the light source or the motion of the observer.

These postulates, particularly the second one, have profound consequences, leading to a departure from classical notions of space and time and requiring a new set of transformation laws between inertial frames. The invariance of the speed of light implies the invariance of the spacetime interval, which is the cornerstone for defining the geometry of flat spacetime and its symmetry group.

3.2. **Symmetries of Flat Spacetime.**

Definition 3.1 (Flat spacetime metric tensor)**.** Let η be the *flat spacetime metric tensor*:

$$\eta = \operatorname{diag}(-1, +1, +1, +1).$$

Definition 3.2 (Orthogonal symmetry group)**.** The group of linear operators that preserve the flat spacetime metric tensor is an *orthogonal group*

$$\mathrm{O}(1,3) := \left\{\Lambda \in \mathbb{R}^{4\times 4} : \Lambda^{\mu}{}_{\alpha}\eta_{\mu\nu}\Lambda^{\nu}{}_{\beta} = \eta_{\alpha\beta}\right\}$$

The subgroup of these matrices that prevent reflections is the *proper* orthogonal group

$$\mathrm{SO}(1,3) := \{\Lambda \in \mathrm{O}(1,3) : \det\Lambda = +1\}$$

The subgroup of this subgroup with that prevents time reversal is called the *proper orthochronous orthogonal group*

$$\mathrm{SO}^{+}(1,3) = \left\{\Lambda \in \mathrm{SO}(1,3) : \Lambda^{0}{}_{0} \geq 1\right\}.$$

Theorem 3.3. *Rotations in* $\mathrm{SO}(3)$ *are a subgroup of* $\mathrm{SO}^{+}(1,3)$*. For* $R_3 \in \mathrm{SO}(3)$*, the* 4×4 *matrix*

$$\Lambda = \begin{pmatrix} 1 & 0 \\ 0 & R_3 \end{pmatrix} \text{ is in } \mathrm{SO}^{+}(1,3).$$

Proof. We check the conditions for $\Lambda \in \mathrm{SO}^{+}(1,3)$:

(1) $\Lambda^{\top}\eta\Lambda = \eta$:

$$\begin{aligned}
\Lambda^{\top}\eta\Lambda &= \begin{pmatrix} 1 & 0 \\ 0 & R_3 \end{pmatrix}^{\top} \begin{pmatrix} -1 & 0 \\ 0 & I_3 \end{pmatrix} \begin{pmatrix} 1 & 0 \\ 0 & R_3 \end{pmatrix} \\
&= \begin{pmatrix} 1 & 0 \\ 0 & R_3^{\top} \end{pmatrix} \begin{pmatrix} -1 & 0 \\ 0 & I_3 \end{pmatrix} \begin{pmatrix} 1 & 0 \\ 0 & R_3 \end{pmatrix} \\
&= \begin{pmatrix} 1 & 0 \\ 0 & R_3^{\top} \end{pmatrix} \begin{pmatrix} -1 & 0 \\ 0 & R_3 \end{pmatrix} \\
&= \begin{pmatrix} -1 & 0 \\ 0 & R_3^{\top}R_3 \end{pmatrix}.
\end{aligned}$$

Since $R_3 \in \mathrm{SO}(3)$, we have $R_3^{\top}R_3 = I_3$. Thus,

$$\Lambda^{\top}\eta\Lambda = \begin{pmatrix} -1 & 0 \\ 0 & I_3 \end{pmatrix} = \eta.$$

(2) $\det(\Lambda) = 1$:

$$\det(\Lambda) = \det(1) \cdot \det(R_3) = 1 \cdot 1 = 1,$$

since $\det(R_3) = 1$ for $R_3 \in \mathrm{SO}(3)$.

(3) ${\Lambda^0}_0 \geq 1$: The $(0,0)$ component of Λ is $\Lambda^0_{R0} = 1$, which satisfies the requirement.

All conditions are met, so $\Lambda \in \mathrm{SO}^+(1,3)$. □

Lemma 3.4. *For a boost in the x-direction that preserves the interval defined by the flat metric, the transformation matrix $\Lambda \in \mathrm{SO}(1,1)$ can be written as*

$$\Lambda = \begin{pmatrix} \gamma & -\gamma v \\ -\gamma v & \gamma \end{pmatrix},$$

where $\gamma := \frac{1}{\sqrt{1-v^2}}$ and the speed *$v \in (-1,1)$ is expressed in natural units. Alternatively, in terms of the* rapidity *parameter $\phi := \operatorname{arctanh} v$, the matrix becomes*

$$\Lambda = \begin{pmatrix} \cosh\phi & -\sinh\phi \\ -\sinh\phi & \cosh\phi \end{pmatrix}.$$

Proof. Consider a transformation acting only on the t and x coordinates:

$$\begin{pmatrix} t' \\ x' \end{pmatrix} = \begin{pmatrix} A & B \\ C & D \end{pmatrix} \begin{pmatrix} t \\ x \end{pmatrix}.$$

Let $\Lambda \in \mathrm{SO}(1,1)$

$$\Lambda = \begin{pmatrix} A & B \\ C & D \end{pmatrix}.$$

We require this transformation to preserve the 2D flat spacetime interval, defined by $\eta := \mathrm{diag}(-1,1)$. The condition is $\Lambda^\top \eta \Lambda = \eta$. Expanding this:

$$\begin{pmatrix} A & C \\ B & D \end{pmatrix} \begin{pmatrix} -1 & 0 \\ 0 & 1 \end{pmatrix} \begin{pmatrix} A & B \\ C & D \end{pmatrix} = \begin{pmatrix} -A^2 + C^2 & -AB + CD \\ -AB + CD & -B^2 + D^2 \end{pmatrix} = \begin{pmatrix} -1 & 0 \\ 0 & 1 \end{pmatrix}.$$

This yields the system of equations:

$$\begin{aligned} -A^2 + C^2 &= -1 \\ -AB + CD &= 0 \\ -B^2 + D^2 &= 1 \end{aligned}$$

And the determinant condition for a special transformation (unit determinant):

$$AD - BC = 1$$

From $-A^2 + C^2 = -1$, we rearrange to $A^2 = 1 + C^2$. Since $C^2 \geq 0$, this implies $A^2 \geq 1$, so $|A| \geq 1$. To ensure that the transformation does not reverse the direction of time, we impose the physical condition $A > 0$. Combined with $|A| \geq 1$, this leads to the requirement $A \geq 1$. Similarly, from $-B^2 + D^2 = 1$, we rearrange to $D^2 = 1 + B^2$. Since $B^2 \geq 0$, this implies $D^2 \geq 1$, so $|D| \geq 1$. The choice between $D \geq 1$ and $D \leq -1$ determines the relative orientation of the spatial x'-axis. For a "standard" boost along the positive x-direction, we choose the convention $D \geq 1$. Given $A \geq 1$, the equation $A^2 - C^2 = 1$ allows us to parameterize $A = \cosh\phi_A$ and $C = s_C \sinh\phi_A$ for some real parameter ϕ_A and $s_C = \pm 1$. Similarly, given $D \geq 1$, the equation $D^2 - B^2 = 1$ allows us to parameterize $D = \cosh\phi_D$ and

$B = s_B \sinh\phi_D$ for some real parameter ϕ_D and $s_B = \pm 1$. Now substitute these parameterizations into $-AB + CD = 0$ ($AB = CD$):

$$(\cosh\phi_A)(s_B \sinh\phi_D) = (s_C \sinh\phi_A)(\cosh\phi_D).$$

Assuming a non-trivial boost (so that $\sinh\phi_A \neq 0$ and $\sinh\phi_D \neq 0$), we can divide by the product $\cosh\phi_A \cosh\phi_D$. Since $\cosh\phi_A \geq 1$ and $\cosh\phi_D \geq 1$, this yields:

$$s_B \tanh\phi_D = s_C \tanh\phi_A.$$

For a single, simple transformation (represented by a single rapidity ϕ), we expect ϕ_A and ϕ_D to be the same or directly related. If we set $\phi_A = \phi_D =: \phi$, then this equation implies $s_B = s_C =: s$. So our parameters become: $A = \cosh\phi$, $C = s\sinh\phi$, $D = \cosh\phi$, $B = s\sinh\phi$. Next, use the determinant condition $AD - BC = 1$:

$$(\cosh\phi)(\cosh\phi) - (s\sinh\phi)(s\sinh\phi) = \cosh^2\phi - s^2\sinh^2\phi = 1.$$

Since $s^2 = (\pm 1)^2 = 1$, this simplifies to $\cosh^2\phi - \sinh^2\phi = 1$. This is the fundamental hyperbolic identity, which is always true for any real ϕ. Thus, the parameters $A = \cosh\phi, D = \cosh\phi, B = s\sinh\phi, C = s\sinh\phi$ satisfy all conditions derived from metric preservation and unit determinant. The matrix is

$$\Lambda = \begin{pmatrix} \cosh\phi & s\sinh\phi \\ s\sinh\phi & \cosh\phi \end{pmatrix}.$$

The specific form stated in the lemma,

$$\Lambda = \begin{pmatrix} \cosh\phi & -\sinh\phi \\ -\sinh\phi & \cosh\phi \end{pmatrix},$$

corresponds to choosing the sign factor $s = -1$. This means $A = \cosh\phi, B = -\sinh\phi, C = -\sinh\phi, D = \cosh\phi$. This is the standard parameterization for a boost. To relate ϕ to the speed v: consider the origin of the S' frame ($x' = 0$). It moves with velocity v in the S frame, so its worldline is $x = vt$. Substituting into $x' = Ct + Dx$:

$$0 = Ct + D(vt) = (C + Dv)t.$$

For this to hold for all $t \neq 0$, we need $C + Dv = 0$, so $v = -C/D$. Using $C = -\sinh\phi$ and $D = \cosh\phi$:

$$v = -(-\sinh\phi)/(\cosh\phi) = \tanh\phi.$$

Thus, the rapidity $\phi = \operatorname{arctanh} v$. Given $v = \tanh\phi$:

$$1 - v^2 = 1 - \tanh^2\phi = \operatorname{sech}^2\phi = \frac{1}{\cosh^2\phi}.$$

Letting $\gamma := \frac{1}{\sqrt{1-v^2}}$, $\cosh\phi = \frac{1}{\sqrt{1-v^2}} = \gamma$ and $\sinh\phi = \tanh\phi\cosh\phi = v\gamma$. This implies that the speed $|v| < 1$ (since ϕ is real, $\tanh\phi \in (-1, 1)$). Substituting $\cosh\phi = \gamma$ and $\sinh\phi = v\gamma$ back into the matrix form $A = \cosh\phi, B = -\sinh\phi, C = -\sinh\phi, D = \cosh\phi$ gives

$$\Lambda = \begin{pmatrix} \gamma & -\gamma v \\ -\gamma v & \gamma \end{pmatrix}.$$

□

Theorem 3.5. *A general boost in an arbitrary direction $\hat{n}$ (a unit 3-vector) with speed v is given by the 4×4 matrix $\Lambda_B \in \mathrm{SO}^+(1,3)$:*

$$\Lambda_B = \begin{pmatrix} \gamma & -\gamma v \hat{n}^\top \\ -\gamma v \hat{n} & I_3 + (\gamma - 1)\, \hat{n}\hat{n}^\top \end{pmatrix}$$

where $\gamma = \frac{1}{\sqrt{1-v^2}}$. In terms of rapidity $\phi = \operatorname{arctanh} v$:

$$\Lambda_B = \begin{pmatrix} \cosh\phi & -\sinh\phi \hat{n}^\top \\ -\sinh\phi \hat{n} & I_3 + (\cosh\phi - 1)\hat{n}\hat{n}^\top \end{pmatrix}.$$

3.3. The Algebra of Spacetime Symmetries ($\mathfrak{so}(1,3)$).

Definition 3.6. The algebra $\mathfrak{so}(1,3)$ is associated with the group $\mathrm{SO}^+(1,3)$. It consists of all 4×4 real matrices X such that a curve $\Lambda(s) = \exp(sX)$ is in $\mathrm{SO}^+(1,3)$ for small s. Differentiating $\Lambda(s)^\top \eta \Lambda(s) = \eta$ at $s = 0$ (where $\Lambda(0) = I$) yields the condition for elements $X \in \mathfrak{so}(1,3)$:

$$X^\top \eta + \eta X = 0.$$

This condition means X is "η-antisymmetric" or "η-skew-adjoint". A convenient basis for this 6-dimensional algebra, acting on coordinates

$$(x^0, x^1, x^2, x^3) \equiv (t, x, y, z),$$

is given by the following generators:

- Rotation generators:
  - J_1: $(J_1)_{23} = -1, (J_1)_{32} = 1$
  - J_2: $(J_2)_{13} = 1, (J_2)_{31} = -1$
  - J_3: $(J_3)_{12} = -1, (J_3)_{21} = 1$
- Boost generators:
  - K_1: $(K_1)_{01} = -1, (K_1)_{10} = -1$
  - K_2: $(K_2)_{02} = -1, (K_2)_{20} = -1$
  - K_3: $(K_3)_{03} = -1, (K_3)_{30} = -1$

Note: These generators X indeed satisfy $X^\top \eta + \eta X = 0$ with $\eta = \mathrm{diag}(-1,1,1,1)$. The verification for any specific generator involves direct matrix multiplication and addition, confirming the condition.

Theorem 3.7 (Commutator relations of the algebra $\mathfrak{so}(1,3)$)**.** *The generators J_i (rotations) and K_i (boosts) satisfy the following commutation relations:*

$$[J_i, J_j] = \epsilon_{ijk} J_k,$$
$$[J_i, K_j] = \epsilon_{ijk} K_k,$$
$$[K_i, K_j] = -\epsilon_{ijk} J_k,$$

where ϵ_{ijk} is the fully antisymmetric symbol.

Proof. Similar to the SO(3) relations found in theorem 2.6, this can be proven explicitly by multiplying each pair of generators and computing their commutators. For example,

$$[J_1, J_2] := J_1 J_2 - J_2 J_1 = J_3,$$

The other pairs can be easily checked the same way. □

Finite Lorentz transformations can be constructed by exponentiating linear combinations of these generators:

$$\Lambda = \exp\left(\sum_i \theta_i J_i + \sum_i \phi_i K_i\right),$$

where $\vec{\theta} = (\theta_1, \theta_2, \theta_3)$ are rotation parameters (angle and axis) and $\vec{\phi} = (\phi_1, \phi_2, \phi_3)$ are boost parameters (rapidity and direction).

3.4. **Kinematic Consequences.** The Lorentz transformations have profound implications for how different inertial observers measure time intervals and spatial lengths. These effects are not mere illusions but are fundamental to the nature of spacetime.

4. Symmetries in Special Relativity: The Inhomogeneous Special Orthogonal Group and Particle States

This section explores the full symmetry group of flat spacetime, the Poincaré group, and its profound implications for defining fundamental particle properties.

4.1. **Structure of Flat Spacetime.**

Definition 4.1. Let $\eta : \mathbb{R}^{1,3} \times \mathbb{R}^{1,3} \to \mathbb{R}$ be the *flat spacetime inner product* of two vectors

$$\eta(x, y) := \eta_{\mu\nu} x^\mu y^\nu$$

where η is still the flat spacetime metric tensor $\eta = \mathrm{diag}(-1, +1, +1, +1)$.

Remark. Note that η is a bilinear form but *not* a metric in the metric space sense, since it is not positive-definite. In particular, $\eta(x, x)$ can be negative, zero, or positive depending on whether x is time-like, null, or space-like.

Definition 4.2 (Inhomogeneous Special Orthogonal Group)**.** The *inhomogeneous special orthogonal group*, $\mathrm{ISO}(1,3)$, is the group of transforms that preserves the squared spacetime interval

$$\mathrm{ISO}(1,3) = \left\{T : \mathbb{R}^{1,3} \to \mathbb{R}^{1,3} \mid \eta\left(T(x) - T(y), T(x) - T(y)\right) = \eta(x - y, x - y)\right\}$$

Theorem 4.3. *Let Λ be an element of the proper orthochronous orthogonal group, $\mathrm{SO}^+(1,3)$ and let $a \in \mathbb{R}^{1,3}$ be any constant spacetime translation vector. The affine transformation $T : \mathbb{R}^{1,3} \to \mathbb{R}^{1,3}$ defined by*

$$T(x) = \Lambda x + a$$

is an element of $\mathrm{ISO}(1,3)$.

Proof. Let $T : \mathbb{R}^{1,3} \to \mathbb{R}^{1,3}$ defined by $x \mapsto \Lambda x + a$ where $\Lambda \in \mathrm{SO}^+(1,3)$ and $a \in \mathbb{R}^{1,3}$. For two events $x, y \in \mathbb{R}^{1,3}$, the difference of their image is

$$T(y) - T(x) = (\Lambda y + a) - (\Lambda x + a) = \Lambda y - \Lambda x$$

Since Λ is a linear operator, it has distributivity over addition:

$$T(y) - T(x) = \Lambda(y - x)$$

Since $\Lambda \in \mathrm{SO}(1,3)$, $\Lambda^{\mu}{}_{\alpha}\eta_{\mu\nu}\Lambda^{\nu}{}_{\beta} = \eta_{\alpha\beta}$, and since all products are contractions over scalars, they commute

$$\begin{aligned}\eta\left(T(y) - T(x), T(y) - T(x)\right) &= \eta_{\mu\nu}(\Lambda(y-x))^{\mu}(\Lambda(y-x))^{\nu} \\ &= \eta_{\mu\nu}\Lambda^{\mu}{}_{\alpha}(y-x)^{\alpha}\Lambda^{\nu}{}_{\beta}(y-x)^{\beta} \\ &= \eta_{\alpha\beta}(y-x)^{\alpha}(y-x)^{\beta}\end{aligned}$$

Hence, the flat spacetime interval is preserved under the transform T, so $T \in \mathrm{ISO}(1,3)$ by definition. □

Theorem 4.4. *All members $T \in \mathrm{ISO}(1,3)$ can be written as $T(x) = \Lambda x + a$ where $\Lambda \in \mathrm{O}(1,3)$ and $a \in \mathbb{R}^4$.*

Proof. Let $T \in \mathrm{ISO}(1,3)$, be a map $T : \mathbb{R}^{1,3} \to \mathbb{R}^{1,3}$ that preserves the flat spacetime interval

$$\eta\left(T(y) - T(x), T(y) - T(x)\right) = \eta(x-y, x-y) \text{ for all } x, y \in \mathbb{R}^{1,3}.$$

Fix a point at $x = 0$ such that $a := T(0)$. Then define $\Lambda(x) := T(x) - a$. If T preserves the interval then Λ also preserves the interval:

$$\begin{aligned}\eta(\Lambda(x) + a - \Lambda(y) - a, \Lambda(x) + a - \Lambda(y) - a) &= \eta(x-y, x-y) \\ \eta(\Lambda(x) - \Lambda(y), \Lambda(x) - \Lambda(y)) &= \eta(x-y, x-y)\end{aligned}$$

Since $\Lambda(0) = T(0) - a = a - a = 0$:

$$\begin{aligned}\eta\left(\Lambda(y) - \Lambda(0), \Lambda(y) - \Lambda(0)\right) &= \eta(y-0, y-0) \\ \eta(\Lambda(y), \Lambda(y)) &= \eta(y,y) \text{ for all } y \in \mathbb{R}\end{aligned}$$

So Λ preserves the norm. Using the polarization identity in the lemma above:

$$\eta(\Lambda(a), \Lambda(b)) = \frac{1}{2}\left[\eta(\Lambda(a) - \Lambda(b), \Lambda(a) - \Lambda(b)) - \eta(\Lambda(a), \Lambda(a)) - \eta(\Lambda(b), \Lambda(b))\right]$$

Since Λ preserves the interval:

$$\eta(\Lambda(a), \Lambda(b)) = \frac{1}{2}\left[\eta(a-b, a-b) - \eta(\Lambda(a), \Lambda(a)) - \eta(\Lambda(b), \Lambda(b))\right]$$

Since Λ preserves the norm:

$$\eta(\Lambda(a), \Lambda(b)) = \frac{1}{2}\left[\eta(a-b, a-b) - \eta(a,a) - \eta(b,b)\right]$$

Which, if we apply the polarization identity above again gives:

$$\eta(\Lambda(a), \Lambda(b)) = \eta(a,b)$$

Which, implies that Λ preserves the bilinear map η.

$$\Lambda(x) + \Lambda(y) = \Lambda(x+y)$$

□

Corollary 4.5. *The composition of two elements $T_1(x) = \Lambda_1 x + a_1$ and $T_2(x) = \Lambda_2 x + a_2$ in $ISO(1,3)$ satisfies*

$$T_1 \circ T_2(x) = (\Lambda_1\Lambda_2)x + (\Lambda_1 a_2 + a_1),$$

so that group multiplication corresponds to

$$(\Lambda_1, a_1) \circ (\Lambda_2, a_2) = (\Lambda_1\Lambda_2, \Lambda_1 a_2 + a_1),$$

Hence, $ISO(1,3)$ is isomorphic to the semidirect product $SO(1,3) \rtimes \mathbb{R}^4$, where $SO(1,3)$ acts linearly on $\mathbb{R}^4$.

Proof. Let $T_1(x) = \Lambda_1 x + a_1$ and $T_2(x) = \Lambda_2 x + a_2$. Then:

$$T_1 \circ T_2 = T_1(\Lambda_2 x + a_2) = \Lambda_1(\Lambda_2 x + a_2) + a_1 = \Lambda_1\Lambda_2 x + \Lambda_1 a_2 + a_1.$$

Thus the composition of two group elements in $ISO(1,3)$ corresponds to:

$$(\Lambda_1, a_1)(\Lambda_2, a_2) = (\Lambda_1\Lambda_2, \Lambda_1 a_2 + a_1),$$

which matches the semidirect product structure $SO(1,3) \rtimes \mathbb{R}^4$. □

4.2. **The Tangent Space Algebra in a Flat Universe.**

4.3. **Noether's Theorem and Conservation Laws from Poincaré Invariance.**

4.4. **Casimir Invariants: Defining Intrinsic Particle Properties.** This subsection discusses operators that commute with all generators of the Poincaré algebra, leading to fundamental invariant properties of particles.

4.4.1. *What are Casimir Invariants?*

4.4.2. *The P^2 Invariant: Mass.* The first key Casimir invariant is $P^2 = P_\mu P^\mu$.
Proof that P^2 is a Casimir Invariant: *[Proof to be filled in by user based on our previous discussion.]*
Physical Significance of P^2: The eigenvalue of P^2 is m^2 (or $-m^2c^2$ depending on metric and c conventions), where m is the invariant rest mass of the particle. This demonstrates that mass is a fundamental Poincaré-invariant characteristic.

4.4.3. *The W^2 Invariant: Spin.* The second key Casimir invariant is $W^2 = W_\mu W^\mu$, where $W_\mu = \frac{1}{2}\epsilon_{\mu\nu\rho\sigma} M^{\nu\rho} P^\sigma$ is the Pauli-Lubanski pseudovector.

4.5. **The Role of Schur's Lemma: Why Casimir Eigenvalues Label Particles.** This subsection explains the mathematical theorem that underpins why Casimir invariants provide constant labels for irreducible representations.

4.5.1. *Statement of Schur's Lemma.*

4.5.2. *Proof of Schur's Lemma. [Proof to be filled in by user based on our previous discussion. This involves showing that the eigenspace of the commuting operator is an invariant subspace, then using irreducibility.]*

4.5.3. *Implication for Casimir Invariants.* Since Casimir invariants commute with all generators (and thus all elements of a representation), Schur's Lemma implies they act as a constant scalar (their eigenvalue) on any irreducible representation.

4.6. **Consequence: Classification of Particles in Special Relativity.**

5. Transition to Curved Spacetime (General Relativity)

The theory of special relativity, while revolutionary, is fundamentally a theory of flat spacetime. It does not incorporate gravity. Newtonian gravity, on the other hand, describes gravity as a force acting instantaneously at a distance, which is incompatible with the finite speed of light posited in special relativity. General relativity emerges from the need to reconcile these ideas.

5.1. **The Principle of Equivalence.**

5.2. **Gravity as Spacetime Curvature.**

6. The Mathematical Language of Curved Spacetime (Differential Geometry)

To describe gravity as the curvature of spacetime, we need the tools of differential geometry. Spacetime is no longer assumed to be flat but is instead a more general mathematical object called a manifold, equipped with a metric tensor that can vary from point to point.

6.1. **Manifolds and Charts.**

6.2. **Tangent Spaces, Vectors, and Tensors in GR.**

6.3. **The Metric Tensor $g_{\mu\nu}$.**

6.4. **Covariant Derivative and Parallel Transport.**

6.5. **Geodesics: Paths of Freely Falling Particles.**

6.6. **Curvature: Riemann Tensor, Ricci Tensor, Ricci Scalar.**

7. The Field Equations of Gravity: How Matter and Energy Dictate Spacetime Curvature

7.1. **The Stress-Energy Tensor $T_{\mu\nu}$.**

7.2. **The Einstein Tensor $G_{\mu\nu}$.**

7.3. **The Equations: $G_{\mu\nu} = \frac{8\pi G}{c^4} T_{\mu\nu}$.**

8. Gravity of a Non-Rotating, Uncharged Spherical Mass (and its Observational Tests)

One of the first and most important solutions to Einstein's field equations describes the spacetime outside a static, spherically symmetric, uncharged massive object. This is known as the Schwarzschild solution.

8.1. **The Metric for a Spherical Mass (Schwarzschild Metric).**

8.2. **Properties: Event Horizon, Gravitational Singularity.**

8.3. **Application 1: Perihelion Precession of Orbits (e.g., Mercury).** The perihelion precession of Mercury's orbit is a seminal observational phenomenon that challenged Newtonian gravity and provided empirical support for the relativistic description of gravitation. Classical Newtonian mechanics predicts orbital motion based on an inverse-square central force, resulting in closed elliptical orbits. However, observations show an additional precession unaccounted for by classical perturbations from other planets. General relativity explains this anomaly through the curvature of spacetime caused by the Sun's mass. By modeling Mercury's trajectory as a geodesic in the Schwarzschild metric, one derives corrections to the Newtonian orbital elements, resulting in a predicted advance of the perihelion that matches observations to high precision. This section presents the derivation of the relativistic corrections to Mercury's orbit, starting from the Schwarzschild solution to Einstein's field equations and applying perturbation techniques to the geodesic equations of motion.

8.4. **Application 2: Relativistic Effects on Clocks and Timekeeping (e.g., for Satellites/GPS).**

8.5. **Application 3: The Bending of Light and Gravitational Lensing.**

8.5.1. *The "Einstein Cross" and other lensing phenomena.*

9. Beyond Simple Cases: Rotating and Charged Massive Objects

9.1. **Need for More General Solutions.**

9.2. **Rotating Massive Objects (e.g., Kerr solution).**

9.2.1. *The Ergosphere.*

9.3. **Charged, Non-Rotating Massive Objects (e.g., Reissner-Nordström solution).**

9.4. **Charged AND Rotating Massive Objects (e.g., Kerr-Newman solution).**

10. Further Striking Phenomena and Future Directions

Appendix A. Tensor Algebra

This appendix summarizes the algebraic properties of basic tensor operations. It assumes all indices are appropriately matched (contravariant vs. covariant).

Notation. Let U, V, W be tensors of the same type, and a, b scalars. Summation is implied over any index that appears once in a raised (contravariant) position and once in a lowered (covariant) position. A summation is *well-defined* if it meets the following criteria:

(1) *Appears exactly once up and once down (paired indices)*
Example:

$$A^\mu B_\mu \quad \text{(valid)} \qquad A^\mu B^\mu \quad \text{(invalid: both raised)}$$

(2) *Variance is compatible (contravariant/covariant pairing)*
Example:

$g_{\mu\nu} A^\nu$ (valid: lowers index) $\qquad g_{\mu\nu} A_\nu$ (invalid: both covariant)

(3) *Summation index appears only once per term (no double use)*
Example:

$$A^\mu B_\mu C^\nu \quad \text{(valid)} \qquad A^\mu B_\mu C_\mu \quad \text{(invalid: reused } \mu\text{)}$$

(4) *Free indices appear only once per term and match across the equation*
Example:

$$A^\mu{}_\nu B^\nu = C^\mu \quad \text{(valid)} \qquad A^\mu{}_\nu B^\nu = C^\rho \quad \text{(invalid: mismatch)}$$

(5) *All summed tensors belong to compatible spaces*
Example:

$$v^\mu w_\mu \quad \text{(valid: dual pairing)}$$
$$F^\mu{}_\nu A^\nu \quad \text{(valid: matrix acting on a vector)}$$
$$g^{\mu\nu} F^\rho{}_\nu \quad \text{(questionable if metric acts on a purely contravariant index)}$$

Addition.

$$U + V = W$$

Properties:

- *Associative*: $(U + V) + W = U + (V + W)$
- *Commutative*: $U + V = V + U$
- *Distributive over scalar multiplication*: $a(U + V) = aU + aV$
- *Identity*: $\exists\ 0$ such that $U + 0 = U$
- *Inverse*: $\exists\ -U$ such that $U + (-U) = 0$

Scalar Multiplication.

$$(aU)^\mu = aU^\mu$$

Properties:

- *Associative*: $a(bU) = (ab)U$
- *Distributes over tensor addition*: $a(U + V) = aU + aV$
- *Distributes over scalar addition*: $(a + b)U = aU + bU$

Tensor Product.

$$(T \otimes S)^{\mu\nu} = T^\mu S^\nu$$

Properties:

- *Associative*: $(T \otimes S) \otimes R = T \otimes (S \otimes R)$
- *Not commutative*: Generally $T \otimes S \neq S \otimes T$
- *Distributes over addition*: $T \otimes (S + R) = T \otimes S + T \otimes R$
- *Compatible with scalar multiplication*: $a(T \otimes S) = (aT) \otimes S = T \otimes (aS)$

Contraction.

$$U^\mu V_\mu = \text{scalar (or lower-rank tensor)}$$

Properties:

- *Generally commutative*: $U^\mu V_\mu = V_\mu U^\mu$ as long as the underlying tensor algebra is over a *commutative ring*
- *Distributive*: $(U + V)^\mu W_\mu = U^\mu W_\mu + V^\mu W_\mu$

Metric Tensor.

$$U_\mu = g_{\mu\nu}U^\nu, \quad U^\mu = g^{\mu\nu}U_\nu$$

Properties:

- *Linear*: $g_{\mu\nu}(aU^\nu + bV^\nu) = aU_\mu + bV_\mu$
- *Inverse operation*: $g^{\mu\nu}g_{\nu\rho} = \delta^\mu_\rho$

Substitution Tensor.

$$\delta^\mu{}_\nu V^\nu = V^\mu$$

Properties:

- *Acts as identity under contraction*
- *Compatible with index placement*

Volume Tensor. The *volume tensor* is denoted by $\epsilon^{\mu\nu\rho\cdots}$ or $\epsilon_{\mu\nu\rho\ldots}$. It is defined by the sign of the permutation of its indices:

$$\epsilon^{\mu_1\mu_2\cdots\mu_n} = \begin{cases} +1 & \text{even permutation of } (0,1,2,\ldots,n{-}1) \\ -1 & \text{odd permutation} \\ 0 & \text{any repeated index} \end{cases}$$

Appendix B. Linear Algebra

A place to hide the scary linear algebra proofs.

B.1. Bilinear Algebra.

Definition B.1 (Bilinear Form)**.** Let V be a vector space over a field $\mathbb{F}$. A *bilinear form* is a map

$$g : V \times V \to \mathbb{F}$$

such that for $\lambda \in \mathbb{F}$ and $u, v, w \in V$:

(1) *Compatibility with scalars in the left argument*:

$$\lambda g(u, v) = g(\lambda u, v)$$

(2) *Compatibility with scalars in the right argument*:

$$\lambda g(u, v) = g(u, \lambda v)$$

(3) *Additivity in the left argument*:

$$g(u + v, w) = g(u, w) + g(v, w)$$

(4) *Additivity in the right argument*:

$$g(u, v + w) = g(u, v) + g(u, w)$$

A *symmetric bilinear map* additionally has the following property

$$g(u, v) = g(v, u),$$

while an *anti-symmetric bilinear map* has the following property

$$g(u, v) = -g(v, u).$$

Theorem B.2. *Let V be a finite-dimensional vector space over a field $\mathbb{F}$, and let $\{e_\mu\}$ be a basis of V. Any bilinear form $g : V \times V \to \mathbb{F}$ can be represented in terms of its components as*

$$g(u, v) = G_{\mu\nu} u^\mu v^\nu,$$

where $G_{\mu\nu} := g(e_\mu, e_\nu)$, and u^μ, v^ν are the components of u, v in the chosen basis.

Proof. Let $\{e_\mu\}_{\mu=0}^{n-1}$ be a basis of V. Then for any $u, v \in V$, there exists components $u^\mu, v^\nu \in \mathbb{F}$ such that:

$$u = u^\mu e_\mu \quad v = v^\nu e_\nu.$$

Since g is bilinear, we can expand

$$g(u, v) = g(u^\mu e_\mu, v^\nu e_\nu) = u^\mu v^\nu g(e_\mu, e_\nu).$$

Let $G_{\mu\nu} := g(e_\mu, e_\nu)$. Then the equation becomes

$$g(u, v) = G_{\mu\nu} u^\mu v^\nu$$

□

Theorem B.3 (Symmetric Polarization Identity)**.** *Let g be a symmetric bilinear form on a vector space V. Then for all $x, y \in V$,*

$$g(x - y, x - y) = g(x, x) - 2g(x, y) + g(y, y).$$

Proof. Let g be a symmetric bilinear form on a vector space V and $x, y \in V$. Expanding into its tensor form.

$$g(x - y, x - y) = g_{\mu\nu}(x - y)^\mu (x - y)^\nu$$

Expanding

$$g_{\mu\nu}(x - y)^\mu (x - y)^\nu = g_{\mu\nu} x^\mu x^\nu - g_{\mu\nu} x^\mu y^\nu - g_{\mu\nu} y^\mu x^\nu + g_{\mu\nu} y^\mu y^\nu$$

After swapping the dummy indices, and since $g_{\mu\nu}$ is symmetric:

$$g_{\mu\nu}(x - y)^\mu (x - y)^\nu = g_{\mu\nu} x^\mu x^\nu - 2g_{\mu\nu} x^\mu y^\nu + g_{\mu\nu} y^\mu y^\nu$$

Or

$$g(x - y, x - y) = g(x, x) - 2g(x, y) + g(y, y).$$

□