

# 5<sup>th</sup> Semester project summery

Dinesh Beniwal (1811061)

February 2021

## 1 Random Walk

Random walks are the most general yet important phenomena in real life. In this project we demonstrated random walk in 2-D plane and visualise the results. To code our random walk we choose any random angle  $\theta$  and move one step in that direction. We continued this process for different number of step and observed these results :

- As we increase the numbers of steps we drifts further from origin in every direction.
- Calculated average displacement in x and y is almost equal to zero( $\Delta X = \Delta Y \approx 0$ ), this means probability to move in any direction is equal.
- Our  $R_{rms}$  values increases with the N(number of steps). Appended figure( $R_{rms}$  vs  $\sqrt{N}$ ) gives us the slope = 0.9671, close to 1 and hence proved that  $R_{rms} \approx \sqrt{N}$ .
- Result  $R_{rms} \approx \sqrt{N}$  tends to agree with equation with the increase in number of steps.

## 2 Ellipsoid

We use Monte Carlo method to determine the volume of ellipsoid. Idea being that throw random points in a cuboid, dimensions equal to axis of ellipsoid and find the probability how many points are satisfying equation of ellipsoid hence inside the ellipsoid.

$$\text{Volume of ellipsoid} = \frac{\text{Number of points inside ellipsoid}}{\text{Total number of points}} \times \text{Volume of cuboid} \quad (1)$$

So to find out which points are inside the ellipsoid we apply and equation:

$$\frac{x^2}{a^2} + \frac{y^2}{b^2} + \frac{z^2}{c^2} \leq 1 \quad (2)$$

After performing the simulation results are:

- Converse rate of Monte Carlo is  $1/\sqrt{N}$ .
- As we increase the number of points our volume converges to analytical volume.
- We also get the convergence of fractional error to zero when we increase the number of points.