

CLINICAL TRIAL PROTOCOL NCT04987211

SECTION 1: STUDY OBJECTIVES

Phase 3 randomized evaluation of Glucotran in adult subjects with Type 2 Diabetes.

SECTION 2.1: INCLUSION CRITERIA

1. Male or female patients aged 18 to 75 years inclusive at time of screening.
2. Documented clinical diagnosis of Type 2 Diabetes Mellitus for at least 180 days prior.
3. Glycated hemoglobin (HbA1c) between 7.0% and 10.5% inclusive at screening.
4. Stable dosage of metformin monotherapy or combination therapy for at least 60 days.

SECTION 2.2: EXCLUSION CRITERIA

1. Estimated Glomerular Filtration Rate (eGFR) less than 30 mL/min/1.73m² at screening.
2. Severe hepatic impairment (Child-Pugh Class C) or AST/ALT greater than 3x ULN.
3. Documented history of acute diabetic ketoacidosis or hyperosmolar coma within 6 months.
4. Known hypersensitivity or allergy to study investigational medicinal product.

SECTION 3: CONTRAINDICATIONS AND SAFETY WARNINGS

Patients presenting with eGFR less than 30 mL/min/1.73m² or end-stage renal disease are strictly contraindicated due to heightened risk of severe lactic acidosis.