Warming and survey effort contribute comparably to new bird distribution records when dated to discovery

Running title: Warming and effort in new bird records

Chenchen Ding[1], et al.

[1] Institute of Ecology, College of Urban and Environmental Sciences, Peking University, Beijing 100871, China

Correspondence: Chenchen Ding (chenchen.ding@pku.edu.cn)

Abstract

New distribution records are widely read as evidence that species' ranges are shifting, yet a record exists only when a species is present, someone detects it and the detection is published. In mainland China, birding visits in the main checklist databases rose about 175-fold between 2002–2004 and 2022–2024, so the ecological and observational signals in these records are hard to separate. We analysed 1,029 provincial new bird records from the China Bird New Record dataset with a discrete-time hazard model of first records across 175,901 species × province × year cells (2002–2024), dating each record to its discovery and modelling the observation process explicitly. Publication lagged discovery by 2.09 years on average. Dating records to publication, the usual practice, inflated the survey-effort hazard ratio by 37% (1.79 versus 1.31) and created a spurious effect of annual climate variability, while accumulated warming was unaffected. After correction, warming of a province relative to each species' historical range (hazard ratio 1.36 per SD, 95% CI 1.22–1.52) and survey effort (1.40, 1.25–1.58) contributed comparably and interacted negatively (0.85, 0.76–0.95). Permutation nulls that preserve the data

structure centred the warming estimate on 1.157, so the species-specific component is about 1.18; the effort estimate fell to 0.980 when its timing was scrambled. A province-by-year random intercept, the largest variance component, raised conditional discrimination from 0.72 to 0.85. The return on effort fell from 2.18 in 2002 to 1.07 in 2024, whereas the return on warming showed no detectable change. The apparent eastward bias of new records relative to species' range centroids was reproduced by geography alone; against an effort-weighted null, records lay slightly further north than expected. New records thus reflect climate and observation in similar measure, and the date attached to a record decides which appears to dominate.

Keywords: biodiversity monitoring; China; citizen science; climate change; discrete-time hazard model; new distribution records; observation bias; range shift; survey effort

Introduction

Records of species in places where they had not been documented before are among the most widely used evidence that ranges are shifting with climate. Range boundaries have moved poleward and upslope in many taxa as temperatures have risen (Parmesan & Yohe, 2003; Chen et al., 2011; Lenoir & Svenning, 2015), and the resulting redistribution affects ecosystems and human well-being (Pecl et al., 2017). In China, compilations of provincial new bird records have been read as evidence of northward and poleward shifts (Xing et al., 2024; Chen et al., 2025). A new record, however, is not an observation of movement. It is the last link in a chain of three events: a species occupies a place, someone detects it there, and the detection is published. Only the first link is ecological, and imperfect detection and uneven sampling can create apparent range changes where none occurred (Tingley & Beissinger, 2009; Hughes et al., 2021).

The difficulty is that observation has grown as quickly as the climate has changed. Documented distributions reflect where people look as well as where organisms live (Boakes et al., 2010; Hortal et al., 2015; Oliver et al., 2021), and many records reported as new fill gaps in knowledge, the Wallacean shortfall, without any change in range (Moura & Jetz, 2021; Diniz-Filho et al., 2023). In mainland China, the annual number of birding visits logged in the eBird and China Bird Report databases rose about 175-fold between 2002–2004 and 2022–2024 (Materials and Methods), the period over which the shifts above were reported. Any series of new records from these years will correlate with the growth of observation, whatever the contribution of climate. Opportunistic records yield unbiased trends only when the recording process is modelled (Isaac et al., 2014), and gaps and biases in such data are best treated as a missing-data problem (Bowler et al., 2025), yet compilations of first records have not been analysed in this way.

Each link in the chain leaves its own signature, and analyses of these compilations have separated none of them. The first concerns time. Compilations date each record by its year of publication, the field that sources report most consistently, but publication lags discovery. Because observation expanded year on year, a record dated to publication is credited to a year in which more people were looking than when it was made, and because recent discoveries are still unpublished, the latest years are incomplete. The size and direction of the resulting bias have not been measured. The second concerns scale. Observation grows in pulses: a survey campaign, a birding festival, a new reserve or a new reporting platform raises documentation in one province in one year. Models with random effects for species and region alone cannot represent such pulses, so their magnitude is unknown, and so is whether they distort estimates of the drivers. The third concerns the benchmark for climate. A warming effect measured relative to each species' own range should be compared with what shared regional climate would produce if

species identity carried no information, and that benchmark need not be unity. A fourth problem follows: projections of record generation under climate scenarios are rarely checked against the range of conditions to which the models were fitted (Elith et al., 2010; Mesgaran et al., 2014; Yates et al., 2018).

We asked whether warming measurably raises the rate at which new provincial bird records are generated in China once the publication and observation steps are modelled explicitly, and how its contribution compares with that of survey effort. We used version 2 of the China Bird New Record dataset (CBNR), which gives dates of discovery as well as of publication for 1,029 provincial records (Ding et al., 2026; Ding et al., in review). Every species × province × year in which a first record could have occurred enters a risk set, and the annual hazard of a first record is modelled with a complementary log-log link, the discrete-time form of a proportional-hazards model (Allison, 1982; Singer & Willett, 1993). Records are dated to discovery, with an offset for the incompleteness of recent years. Survey effort enters as a covariate and province-year pulses as a random intercept, and warming is expressed relative to each species' historical Chinese range and evaluated against permutation nulls that preserve the structure of the data. A decomposition that changes one analytical choice at a time shows what each correction does. None of these steps is specific to birds or to China; each applies to any dated compilation of first records.

We tested four predictions, specified before the robustness analyses were run. (H1) Dating records to publication inflates the effort coefficient and creates a spurious effect of year-to-year climate variability but leaves accumulated warming unchanged, because a lag of one or two years moves a record into a year with more observers and different weather while barely altering a 15-year mean. (H2) Province-year pulses are the largest source of unexplained variation in the

hazard, yet absorbing them leaves the discrimination achieved by the fixed effects unchanged, because the pulses are short-lived and unrelated to long-term trends. (H3) Warming and effort interact negatively. A species can be recorded for the first time in a province only once, so where effort has long been high, many of the species that warming favours will already have been recorded, and the pairs still at risk respond less. (H4) For the same reason, the return on effort declines through time as the stock of detectable but undocumented occurrences is drawn down, whereas the return on warming, which keeps adding candidates as climates shift, does not.

Materials and Methods

Event data and dating

New provincial-level bird distribution records for China were taken from version 2 of the China Bird New Record dataset (CBNR; 1,029 records covering 2000–2025; Ding et al., 2026), compiled from peer-reviewed publications with a large-language-model-assisted extraction pipeline followed by manual validation of every record (Ding et al., in review). Each record carries a discovery date and a source publication year. We parsed a four-digit year from the discovery date field, obtaining a usable year for 1,028 of 1,029 records. Nine further records reported a discovery year later than the publication year, which is internally inconsistent; for these ten records in total we substituted publication year minus the median lag of one year, and flagged the substitution in the released data.

Species names were harmonised by trying, in order, the Catalogue of Life China 2026, BirdLife International v10 and Clements v2025 names supplied with the compilation, retaining the first that matched the candidate pool; provincial names were harmonised by mapping Xizang to Tibet.

Taiwan, Hong Kong and Macao were excluded because the survey-effort compilation does not cover them.

For each species × province pair we retained the earliest event year; later reports of the same pair were treated as re-documentation. Pairs whose earliest record predates 2002 describe species already present at the start of the analysis window and were removed from the risk set as prevalent rather than incident cases. Every remaining event was forced into the risk set irrespective of its species-distribution-model status.

Reporting completeness

Let F be the empirical cumulative distribution of non-negative publication lags and $P\max = 2025$ the latest publication year in the compilation. A record discovered in year t can have appeared only if its lag is at most $P\max - t$, so the expected share of year-t discoveries already reported is $c(t) = F(P\max - t)$. Lag stationarity across discovery years was verified by Spearman correlation, a Wilcoxon comparison of the two halves of the well-reported period (2002–2018), and direct comparison of the cumulative distributions. Under incomplete reporting the observed hazard is approximately $c(t) \cdot h(t)$; with a complementary log-log link this is exactly an offset, $\log(-\log(1 - p\text{obs})) = \eta + \log c(t)$, which we include in every model. Sensitivity analyses omitting the offset and truncating the window at 2021 are reported.

Risk set

The candidate pool for each species comprises every province in which its species distribution model marks at least a threshold number of suitable grid cells, plus every province in which the species was actually recorded. Cells are those of the 2.5-arc-minute climate surface and have a median area of 18.3 km² (about 4.3 km on a side), so the three thresholds of 50, 100 and 200

cells require roughly 0.9, 1.8 and 3.7 thousand km² of suitable habitat within the province. Each species × province pair contributes one row per year from 2002 until the year of its first record, after which it exits the risk set (absorbing exit). At the 50-cell threshold this yields 8,319 pairs, 185,478 rows and 657 events.

Climate variables

Monthly temperature surfaces at 10 arc-minutes for 1980–2024 were taken from the WorldClim 2.1 downscaling of CRU TS 4.09 (Harris et al., 2020; Fick & Hijmans, 2017). We computed four indicators: annual mean temperature, winter mean temperature, warmest-month maximum temperature and coldest-month minimum temperature. Provincial series $T(p,t)$ are area-weighted means over the 100 km analysis grid, weighted by the area of overlap between each grid cell and the province; species-range series $N(s,t)$ are means over the species' Chinese range polygon (BirdLife International & Handbook of the Birds of the World, 2024). Baselines $T\text{base}$ and $N\text{base}$ are 1980–2000 means at both ends.

The species-referenced climate gradient is

$$x(s,p,t) = [T(p,t) - T\text{base}(p)] - [N(s,t) - N\text{base}(s)],$$

that is, how much more the target province has warmed than the species' own historical range. This is decomposed, strictly backward-looking, into an accumulated component and a residual:

$$\text{clim_change}(s,p,t) = \text{mean of } x \text{ over } [t - W + 1,\ t],\quad \text{clim_var}(s,p,t) = x(s,p,t) - \text{clim_change}(s,p,t),$$

with $W \in \{5, 10, 15, 20\}$ years. Because the surfaces extend back to 1980, no analysis-period rows are lost to the trailing window at any W. The main specification uses annual mean temperature with $W = 15$.

Survey effort

Provincial annual survey effort was compiled from deduplicated eBird/GBIF occurrence records and China Bird Report checklists (http://www.birdreport.cn) through four proxies: visits, distinct observers, birding days and records. Thirty-nine province-years, all in Hong Kong, Macao and Taiwan, have no coverage in the source database. A further 24 mainland province-years were recorded as exactly zero. We treat these as coverage gaps rather than true zeros: in 22 of 24, the adjacent years in the same province record between one and 27 visits. In the main analysis these cells are missing; sensitivity analyses zero-fill them (the previously published treatment) and impute them by within-province linear interpolation on the log scale, which is bounded by the neighbouring observed values. Each proxy is log1p-transformed and standardised within the analysis scope separately for each missing-data rule. Summed over mainland provinces, annual visits rose from a mean of 493 in 2002–2004 to 86,380 in 2022–2024.

Model

We model the annual hazard that species s is first recorded in province p in year t as a discrete-time proportional-hazards model with a complementary log-log link (Allison, 1982; Singer & Willett, 1993; Brooks et al., 2017):

$$\log(-\log(1 - h(s,p,t))) = \beta_0 + \beta_1 \cdot \text{clim_change_z} + \beta_2 \cdot \text{effort_z} + \beta_3 \cdot (\text{clim_change_z} \times \text{effort_z}) + \beta_4 \cdot \text{clim_var_z} + \log c(t) + u_s + v_p + w_{pt},$$

with $u_s \sim N(0, \sigma^2_{species})$, $v_p \sim N(0, \sigma^2_{province})$ and $w_{pt} \sim N(0, \sigma^2_{province \times year})$. All predictors are standardised on the analysis sample, so coefficients are hazard ratios per standard deviation. Models were fitted with glmmTMB 1.1 (Brooks et al., 2017).

Six random-effect structures were compared on four criteria: AIC and BIC; explanatory power (McFadden pseudo-R^2, Nakagawa marginal and conditional R^2 (Nakagawa & Schielzeth, 2013), and marginal and conditional AUC); variance-component identifiability; and the ecological interpretability of each term. The structure reported above was frozen before the robustness analyses were run.

Resampling and null models

Three design-preserving permutation nulls each break exactly one association: N1 shuffles warming among species within a province-year; N2 shuffles effort across years within a province; N3 shuffles the event indicator among species at risk within a province-year. Each leaves risk-set size, per-province-year event counts, the marginals of effort and completeness, and the random-effect hierarchy identical to the data. Each null was run 199 times and two-sided empirical *P* values were computed from the permutation quantiles. Cluster bootstraps resampled species (392 clusters) and provinces (31 clusters) with replacement, 300 times each; only the resampled grouping variable was renamed, so that the crossed random-effect structure being bootstrapped is preserved. A parametric bootstrap simulated 150 response vectors from the fitted model, including fresh random-effect draws, and refitted. Influence was assessed by refitting with each province, and each of ten randomly assigned species deciles, held out.

A small number of resamples converge to degenerate optima. These leave percentile intervals unaffected but corrupt any statistic based on a mean or a standard deviation, so draws with any |log hazard ratio| exceeding 2 were excluded and their number is reported (3 of 199 in N3; 3 of 300 in the species bootstrap; none elsewhere). Inference is based on percentile quantities

throughout. All refits used glmmTMB with `profile = TRUE`, which changes coefficients by less than 10^{-3}.

Diagnostics

Residual diagnostics used simulation-based scaled residuals with 500 simulations (Hartig, 2024): uniformity, dispersion, outliers, zero inflation, and grouped uniformity by year and by province. Spatial autocorrelation was assessed as Moran's *I* on province-mean residuals with inverse-distance weights and a 999-permutation null. Proportional hazards were tested by adding interactions between each focal term and standardised year.

Projections

CMIP6 four-model ensemble median deltas for annual mean temperature were computed for SSP2-4.5 and SSP5-8.5 at 2030, 2050 and 2080, separately for each province and each species range, and added to both ends of the climate gradient. Effort was grown at 0.30 (SSP2-4.5) and 0.60 (SSP5-8.5) standard deviations per decade. Predictions set the province-by-year random effect to its expectation and the completeness offset to log 1, so the projected quantity is the latent generation hazard. Support was defined as the 1st–99th percentile of the fitted accumulated warming and effort; main projection figures aggregate only cells inside this range, and the share of qualifying cells is reported on every panel.

To separate the two scenario axes we also projected the frozen model over a 2 × 2 × 2 grid: SSP2-4.5 or SSP5-8.5 deltas; effort frozen at 2024 or growing at 0.6 SD per decade (observed pace 1.19); horizons 2030 and 2050. Province-by-year effects for future years were set to their expectation, species and province effects retained, and provincial expectations summed over species. Future county surfaces hold the 2024 allocation shares fixed.

Model-family comparison

The projection comparison used a gradient-boosted model (XGBoost, 400 rounds, learning rate 0.05, depth 4) (Chen & Guestrin, 2016) on the same three predictors, with exact TreeSHAP attributions (Lundberg et al., 2020). For the cross-validated comparison, a random forest (ranger, 800 trees, class-balanced down-sampling) and XGBoost (300 rounds, learning rate 0.05, depth 5, positive-class weight n_0/n_1) were fitted to the province-level risk set under the main model's four validation schemes with identical folds and to the prefecture- and county-level allocation sets under temporal and leave-one-province splits with within-stratum rank skill. Hyperparameters were fixed, not tuned per fold.

Protected areas

Protected-area analyses (Supporting Information Note S6) used the 1,028 mapped nature reserves in China (China Nature Reserve Specimen Resource Sharing Platform, 2024).

Direction of new records

Bearings from the centroid of each species' global range (AVONET; Tobias et al., 2022) to each record, computed as the planar angle of the longitude and latitude differences, were binned into eight 45° sectors. Because a uniform expectation ignores where China lies relative to these centroids, observed sector counts were compared with two Monte Carlo nulls (999 draws) that reassign every record to one of its species' candidate provinces, located at the mean position of that province's new records, with equal probability (geometry) or with probability proportional to the province's 2002–2024 survey visits (geometry + effort). Sector-wise two-sided *P* values were Holm-corrected, and an omnibus χ^2 against the null expectation was referred to its Monte Carlo distribution. A sensitivity analysis placed each province at a point on its surface.

Results

The risk set and the dating problem

Of the 1,029 records, 1,028 (99.9%) carry a parseable discovery date and 684 match a species × province cell in the candidate pool; 657 fall within the 2002–2024 window (Supplementary Table S7; Supplementary Fig. S16). Twenty-six records predate 2002 and describe species already present at its start, so they were removed from the risk set as prevalent cases, whereas eleven pairs published in 2025 but discovered by 2024 enter it only under discovery-year dating.

Publication lags discovery by a median of one year and a mean of 2.09; 81.7% of records lag by at least one year and 26.8% by three or more (Figure 2a). Re-dating moves 526 of 633 comparable events (83.1%), by a median of one year. The two conventions give qualitatively different series (Figure 2b): the publication-year series rises monotonically to 67 events in 2024, whereas the discovery-year series peaks at 59 in 2019 and falls to 14. Neither can be used as it stands. The first aggregates delayed reports, and the second is right-censored because many records from 2023–2024 are not yet in print.

The censoring can be quantified directly. The compilation extends to publications of 2025, and the pooled lag distribution, which is stationary across discovery years, implies that the share of discoveries already reported falls from 99.7% for 2002 to 89.5% for 2021, 83.4% for 2022, 73.2% for 2023 and 56.7% for 2024. This completeness enters every model as an offset (Materials and Methods; Supplementary Note S1).

After requiring non-missing effort and climate components, the risk set at the 50-cell threshold (8,319 species × province pairs, 185,478 rows and 657 events) yields modelling data of 175,901 rows, 649 events, 392 species and 31 provincial units.

Publication-year dating inflates the apparent role of survey effort

We decomposed the difference between the previously published specification and the corrected one into five steps (each changing one analytical choice; step S3 also recovers 215 rows and 11 events whose unweighted provincial climate series were missing, so S0–S2 and S3–S5 differ slightly in sample) (Figure 2c; Table 1).

Re-dating alone moves the survey-effort hazard ratio from 1.788 to 1.305, so that the publication-year estimate is 37% larger than the corrected one, and moves the annual-variability hazard ratio from 0.850 ($P = 1.8 \times 10^{-4}$) to 0.971 ($P = 0.48$). The accumulated-warming coefficient barely changes (1.394 → 1.437), and neither does the interaction (0.876 → 0.859), as H1 predicted. The remaining corrections are comparatively minor: treating structural zeros in the effort database as coverage gaps moves effort by −0.038; area-weighting the provincial climate series moves it by +0.022; the completeness offset restores +0.159, recovering signal that right-censoring had attenuated; and the province-by-year random intercept removes a further 0.044.

Both shifts follow from the timing of publication. Publication year and the size of the observational database increase together, so a record assigned to its publication year sits in a cell whose recorded effort exceeds the effort that produced the observation. The same shift detaches the record from the weather of the year in which the bird was seen, which creates the spurious climate-variability effect. A 15-year trailing mean of warming barely moves when a record shifts by one or two years.

Warming and effort contribute comparably

In the frozen main model, both drivers raise the hazard and the effect sizes are similar (Figure 1a; Table 2). One standard deviation of accumulated warming relative to a species' historical range

multiplies the annual hazard by 1.362 (95% CI 1.216–1.524; $P = 8.3 \times 10^{-8}$); one standard deviation of survey effort multiplies it by 1.404 (1.245–1.583; $P = 2.9 \times 10^{-8}$). Annual climate variability has no effect (0.995, 0.909–1.090; $P = 0.92$). The two main effects interact negatively (0.849, 0.761–0.948; $P = 3.7 \times 10^{-3}$): the marginal effect of warming is steepest where effort is low and flattens where coverage is already dense (Figure 1b).

The warming coefficient must be read against a benchmark above unity. Permuting warming among species within a province-year, or the event indicator among species at risk, still returns 1.157–1.167, because both permutations preserve the covariance between provincial-annual climate and event counts. The observed 1.362 lies outside that null ($P = 0.005$ for both), but only the increment, $1.362/1.157 \approx 1.18$, is attributable to referencing warming to each species' range; the remainder is a baseline that the null shares. Effort behaves differently. Permuting it across years within a province collapses its hazard ratio to 0.980, with a null interval covering unity, so the effort effect arises from the timing of effort within provinces, and contrasts between provinces contribute little. A province-cluster bootstrap widens the Wald intervals to 1.20–1.72 for warming and 1.21–1.76 for effort; both lower bounds exceed unity, and the interaction stays below it (0.74–0.99).

In natural units (Figure 4c), one standard deviation of accumulated warming is 0.179 °C: a province that has warmed by that much more than a species' own historical Chinese range carries a 36% higher annual hazard for that species. Across the observed range of the variable (−0.65 to +0.61 °C), the implied hazard spans roughly ninefold. One standard deviation of survey effort corresponds to a 7.75-fold increase in annual provincial visits, from a median of 60 visits per year to about 465.

Much of the variation in the climate variable lies among species within the same province and year (Figure 4a, b): 39% of the variance in accumulated warming and 55% of that in the raw gradient. In Yunnan in 2020, for example, accumulated warming ranges from +0.309 °C for *Aethopyga christinae*, whose Chinese range has warmed less than Yunnan has, to −0.280 °C for *Coracias garrulus*, whose range has warmed more. A province-level warming index would give both species the same value.

Four independent importance criteria agree that warming and effort are of comparable weight, with warming slightly ahead on discrimination (Figure 1c). Dropping accumulated warming costs 28.7 AIC units and 0.053 of marginal AUC; dropping survey effort costs 29.5 AIC units and 0.032 of marginal AUC. The interaction costs 6.5 AIC units. Dropping annual variability improves AIC by 2.0 units; we retain it as one half of a pre-specified decomposition.

The discrimination ladder shows the same balance (Figure 1d). With the random structure held at the main model, fixed-effect-only AUC rises from 0.411 for the null model to 0.559 with effort alone, 0.580 with accumulated warming alone, and 0.610 with both. Climate alone discriminates better than effort alone; neither alone approaches the pair.

Separating the observation process requires a province-by-year level

We evaluated six random-effect structures against four criteria, of which AIC is only one (Table 3; Supplementary Fig. S1). Adding a province-by-year intercept to the conventional species + province structure improves AIC by 93.4 units, raises conditional R^2 from 0.274 to 0.411, raises conditional AUC from 0.721 to 0.854, and raises the intraclass correlation from 0.172 to 0.358. Marginal AUC, the discrimination attributable to the fixed effects, is flat across all six structures

(0.607–0.613), so the added level absorbs observational heterogeneity without redistributing the fixed-effect signal, as H2 predicted.

Two further structures were tested. A species-specific random slope on accumulated warming (ΔAIC 4.5 relative to the best) and a province-specific random slope on effort (ΔAIC 0.0, the AIC optimum) both fit marginally better, but both *reduce* conditional R^2 (0.403 and 0.395 respectively, against 0.411 for the simpler structure), and their fixed effects are indistinguishable from those of the main model (effort 1.397 and 1.372; warming 1.348 and 1.366). We therefore froze the three-level intercept structure, the most parsimonious of the three statistically indistinguishable options and the one with the highest explanatory power. No variance component collapses in any structure (minimum standard deviation 0.274).

What each level of the model represents

Each random term corresponds to a distinct process, and the fitted magnitudes are themselves results (Figure 4d, Figure 5); the ecological rationale for every component is given in Supplementary Table S2.

Species (SD 0.405). Species differ in intrinsic detectability for reasons no covariate in the model captures. One standard deviation multiplies the hazard by 1.50, and the fitted species span a 3.7-fold range (Figure 5a).

Province (SD 0.326). One standard deviation multiplies the hazard by 1.39 (Figure 5b).

Province by year (SD 0.804). This is the largest variance component in the model. One standard deviation multiplies the hazard by 2.23, more than either fixed effect, and the fitted levels span a thirteenfold range (Figure 5c). It represents the scale at which observation operates: a regional

survey campaign, a provincial birding festival, a newly established protected area or a new local reporting channel raises the documentation rate for one province in one year. Because such pulses are not shared nationally, a model with only species and province intercepts cannot absorb them.

Reporting completeness offset. This known correction consumes no degrees of freedom; omitting it attenuates the effort coefficient from 1.372 to 1.245, because censoring is concentrated in the recent, high-effort years.

Migratory strategy shifts the baseline hazard but moderates neither driver

New provincial records could arise by different routes in species with different migratory strategies. Residents disperse slowly, whereas long-distance migrants can appear far outside their range as vagrants or storm-displaced individuals and are visible only on passage. We therefore tested whether strategy moderates either driver (Supplementary Table S3; Supplementary Fig. S3).

The analysis uses the three groups with a known strategy: 137 resident, 72 partial-migrant and 107 long-distance-migrant species, carrying 218, 156 and 175 events. Species with no recorded strategy are excluded. Their event rate is 0.791% against 0.29–0.42% in the other groups, because a species without trait data entered the candidate pool only by virtue of having a record.

Migratory strategy shifts the baseline hazard: adding it as a main effect improves fit (ΔAIC 6.2) and is the best model in the ladder. It does not moderate either driver: adding warming × strategy, effort × strategy, both, or the full three-way term each *worsens* AIC, and none is supported by a likelihood-ratio test ($P = 0.693$, 0.854, 0.899 and 0.256 respectively).

Fitting each group separately agrees. Accumulated warming raises the hazard in every group: 1.363 (95% CI 1.158–1.605) in residents, 1.507 (1.244–1.826) in partial migrants and 1.514 (1.265–1.812) in long-distance migrants. Survey effort does too (1.293, 1.270 and 1.344). The intervals overlap almost completely; long-distance migrants, for which we expected warming to matter least, have the highest point estimate, though it is indistinguishable from the others. A warming × effort interaction confined to long-distance migrants, and the identifiability constraints of the stratified fits, are reported in Supplementary Note S2; because the test of group differences is not significant, we treat that pattern as a hypothesis still to be tested.

Which species, and which provinces

Two coarser questions complement the hazard model: which species are more likely to yield a new provincial record at all, and which provinces accumulate more of them. We followed the framework of the companion mammal study (Ding et al., 2025), adapted to birds (Supplementary Figs S4, S5; Supplementary Tables S4, S5).

Species level. The pool is the 1,445 Chinese bird species of the national ecological trait database; 1,312 match the dated Clements phylogeny, and 361 of these acquired at least one new provincial record, carrying 636 of the 657 events (Supplementary Note S3).

Two effects hold under all three phylogenetic treatments: partial migrants are about twice as likely as residents to yield a new provincial record (odds ratio 1.91–2.50, all $P < 0.001$), and range size raises the odds by 1.23–1.46 per standard deviation. Partial migrants also carry the highest raw event rate in the hazard model (0.417% against 0.29–0.33%). Body mass, clutch size, habitat breadth, diet breadth, number of congeners, hand-wing index, IUCN status, endemism and trophic niche show no detectable effect. In mammals, by contrast, smaller-bodied,

nocturnal and data-deficient species were more often newly recorded (Ding et al., 2025). A stratified fit recovers two group-specific patterns, including a vagrancy signature among full migrants (Supplementary Table S6); refitting without species whose traits were imputed leaves every conclusion intact (Supplementary Note S3).

Range size has to be measured as an area (Supplementary Table S4). Two area measures, both independent of the response, agree: global range area gives an odds ratio of 1.456 (95% CI 1.204 –1.762, $P = 1.1 \times 10^{-4}$, n = 1,298) and Chinese range area, taken from BirdLife polygons clipped to China, gives 1.653 (1.420–1.925, $P = 9.0 \times 10^{-11}$, n = 1,003). The count of provinces occupied does not qualify: published in 2022, it already contains the 2002–2021 records used as the response, returns an inflated odds ratio of 2.09 ($P = 5 \times 10^{-15}$), and absorbs the area effect entirely when both are entered (area 0.96, $P = 0.70$).

Province level. Counts of new records per province are overdispersed relative to Poisson (dispersion 5.60, $P = 2.3 \times 10^{-17}$); a negative binomial is adequate (1.24, $P = 0.20$, $\theta = 6.11$) and passes simulation-based diagnostics. Only regional species richness is significant in the count model (incidence rate ratio 1.27, $P = 0.036$). Treating recent effort as an offset makes the response a discovery rate per unit effort: provinces already well covered in 2002–2008 yield fewer records per unit of recent effort (0.71, $P = 0.028$), as do those with higher GDP per capita (0.61, $P = 0.0027$). Hierarchical partitioning attributes 51.2% of explained deviance to species richness, 33.3% to area and 9% to the effort terms; the full model explains 20.5%.

At the provincial scale, then, effort appears as diminishing returns, the same signal that the hazard model shows as an effort coefficient falling through time. Supplementary Note S3 documents the one departure from the mammal framework.

Records lie slightly north of where geometry and effort place them

Bearings from the centroid of each species' global range (Tobias et al., 2022) to its new records are far from uniform ($\chi^2 = 580.1$, df = 7, $P = 4.6 \times 10^{-121}$; Figure 6a): east holds 237 of 642 records (36.9%) and northeast 179 (27.9%), whereas south and west hold 20 and 19. A uniform expectation is the wrong benchmark, however. Many of these species have ranges centred west or south of China, so a record anywhere in China lies east or north of the centroid by position alone. Reassigning each record at random among its species' candidate provinces reproduces the pattern closely: 234.8 records are expected to the east and 158.6 to the northeast, no sector departs from expectation after Holm correction, and the omnibus departure is small ($\chi^2 = 10.8$, Monte Carlo $P = 0.039$; Figure 6a, b; Supplementary Tables S9, S10).

Weighting the candidate provinces by their 2002–2024 survey visits adds the observation process to the null, and the departure that remains points north ($\chi^2 = 25.8$, $P = 0.001$). Records fall to the northeast (179 against 153.2 expected) and northwest (31 against 17.9) more often, and to the east (237 against 263.9) and south (20 against 32.1) less often, than the distribution of survey effort predicts (Holm-adjusted $P = 0.016$–0.036). The three northern sectors hold 40.2% of records against 33.0% expected. Of the five best-sampled orders, only Passeriformes departs from both nulls under both placements of provinces ($P \leq 0.007$), again with an excess in its northern sectors (38.7% against 30.4%); the four smaller orders are consistent with geometry or depart from it under only one placement (Supplementary Table S10; Supplementary Fig. S15).

The eastward concentration of raw bearings therefore reflects China's position relative to species' global ranges and carries no information about observers or climate. Once geometry and effort are accounted for, new records lie modestly poleward of expectation, and placing each

province at a point on its surface instead of at the mean position of its records gives the same result ($\chi^2 = 12.0$ and 22.5).

The return on survey effort is declining

The proportional-hazards assumption holds for accumulated warming (interaction with time $\beta = -0.022$, $P = 0.82$) but not for effort ($\beta = -0.211$, $P = 7.7 \times 10^{-4}$), and the departure is itself a result. Allowing the effort coefficient to vary through time, its hazard ratio falls from 2.18 (95% CI 1.63–2.90) in 2002 to 1.07 (0.88–1.30) in 2024, crossing into non-significance in the early 2020s (Figure 5d). An additional standard deviation of effort now yields far fewer first records than it did at the start of the period. The warming term shows no such decline, as H4 predicted, although we did not test the two time interactions against each other.

Robustness

None of the conclusions depends on the analytical choices we were free to vary (Figure 3).

Species-distribution-model threshold. Raising the requirement from 50 to 200 suitable cells (about 0.9 to 3.7 thousand km²) changes the risk set from 175,901 to 155,435 rows but leaves the coefficients essentially fixed (effort 1.404, 1.405, 1.411; warming 1.362, 1.376, 1.372; interaction 0.849, 0.851, 0.850).

Climate indicator. Annual mean temperature carries the signal; warmest-month maximum temperature carries a weaker version of it (hazard ratio 1.30, ΔAIC 9.0 behind the best cell) but shows no interaction with effort; coldest-month minimum temperature and winter mean temperature show no signal at any window (hazard ratios 0.96–1.08, all $P > 0.12$).

Accumulation window and climate baseline. We tested windows of 5, 10, 15 and 20 years against 1980–2000 and 1970–2000 baselines (Supplementary Note S5; Supplementary Fig. S2). The warming coefficient rises with window length in every series, from 1.16–1.21 at five years to 1.36–1.40 at twenty, as expected of a process that integrates climate over decades. A 3–23-year grid (Supplementary Table S8) places the optimum at 18–20 years; the pre-specified 15 years lies within 2.2–5.6 AIC units of it. The effort coefficient is flat at 1.34–1.41 and the interaction negative throughout (0.82–1.00), so the two terms are separately identified. The baseline changes the estimate only modestly: at $W = 15$ the hazard ratio is 1.361 under CRU 1970–2000 and 1.307 under CRU 1980–2000, and the main estimate (1.362) is within 0.001 of the former. Because the 0.5° CRU grid smooths topography, one standard deviation of accumulated warming is 0.058 °C in the CRU series against 0.179 °C in the main one; natural-unit statements refer to the latter.

Effort proxy and missing-data rule. All twelve combinations of four proxies (visits, observers, birding days, records) and three treatments of the 24 implausible zero cells (coverage gap, bounded within-province interpolation, zero-filling) give effort hazard ratios between 1.25 and 1.52 and warming hazard ratios between 1.35 and 1.37.

Model adequacy

Simulation-based residual diagnostics indicate a well-calibrated model (Supplementary Table S1; Supplementary Fig. S9). Scaled residuals are uniform (Kolmogorov–Smirnov $P = 0.74$), dispersion is correct ($P = 0.59$) and there is no zero inflation ($P = 0.59$). Two of 23 year levels and two of 31 province levels show within-group deviation at $P < 0.05$, close to the nominal rate. Province-mean residuals show no spatial autocorrelation (Moran's $I = -0.038$ against an

476 expectation of −0.033; permutation $P = 0.86$). The outlier test is significant (95 outliers, $P <$
477 0.001), as expected for a binary outcome with an event rate of 0.37%.

478 Three resampling analyses complete the assessment (Supplementary Fig. S11). The design-
479 preserving nulls place every reported coefficient except annual climate variability outside its
480 null. A 300-resample cluster bootstrap shows that provincial clustering inflates standard errors
481 1.26–1.59-fold and species clustering does not (0.86–1.17); Table 2 reports both intervals. A
482 150-simulation parametric bootstrap recovers the coefficients with bias below 0.01 on the log-
483 hazard scale and coverage of 91.3–95.3%. Leaving out each province, or each of ten species
484 deciles, moves the coefficients by at most 4.2%, so no single province or species group carries
485 the result. A refit allowing both focal effects to vary through time is reported in Supplementary
486 Note S4.

487 **How far can these relationships be projected?**

488 We projected the fitted hazard under SSP2-4.5 and SSP5-8.5 for 2030, 2050 and 2080 (Materials
489 and Methods) and asked how much of each scenario lies inside the range over which the model
490 was fitted (Figure 7b).

491 The historical species-specific gradient has a standard deviation of only 0.18 °C, because
492 provincial and range-wide anomalies largely move together. Future province-minus-range
493 warming differentials reach 2 °C, more than ten standard deviations beyond the fitted range.
494 Consequently, only 47.8% of species-province cells fall inside the fitted covariate support under
495 SSP2-4.5 in 2030, 10.4% in 2050 and 2.1% in 2080; under SSP5-8.5 the corresponding figures
496 are 34.2%, 2.1% and 1.9%.

Restricted to cells inside the support, the mechanistic model projects hazard ratios relative to 2024 of 0.79 to 2.94 (Supplementary Fig. S6). Applied without the mask, the same model produces ratios up to 78 (Supplementary Fig. S10).

A gradient-boosted model fitted to the same three predictors shows why the support matters. Within the observed range the two model families track each other across province-years (Spearman $\rho = 0.54$); outside it they agree on the direction of change in only 23.7% of cells, falling to 6.5% under SSP5-8.5 in 2080 (Figure 7c). The cause is structural: tree ensembles saturate once covariates leave the training support, whereas the complementary log-log model extrapolates linearly on the log-hazard scale. TreeSHAP attributions within the training range are monotone for both drivers (Supplementary Fig. S8).

With climate and survey effort as independent scenario axes (Supplementary Fig. S17), freezing effort at its 2024 level and switching from SSP2-4.5 to SSP5-8.5 leaves the national expectation essentially unchanged (23.2 versus 23.1 first records per year in 2030; 23.6 versus 23.7 in 2050), whereas effort growing at 0.6 SD per decade, half the observed pace, raises the 2050 expectation to 57–62 records per year with only 2% of species–province cells still inside the fitted support (40–44% under frozen effort). The projected number of records is governed by the effort scenario, and its increase lies almost entirely outside the fitted support.

Random forest and XGBoost models fitted under identical validation designs lead the mechanistic model on leave-species folds but trail it on temporal and leave-one-province prediction, and all three families agree on where the next records will fall (Spearman ρ 0.62–0.90; Supplementary Note S7; Supplementary Figs S18, S19). On the same scenario grid they diverge sharply (Supplementary Figs S20, S21): calibrated to a common 2024 total, the mechanistic model reaches 57.4 records per year under SSP2-4.5 with continued effort growth in

2050, the random forest stays at 23.0 and XGBoost falls to 6.1, differences produced by boundary leaves that the data cannot adjudicate.

Records inside nature reserves track the observation process

Twenty-six per cent of new records fall inside one of the 1,028 mapped Chinese nature reserves, against a land-area share of 10.4%, an apparent 2.5-fold enrichment that would ordinarily be read as evidence that the network is receiving redistributing species. Birding effort, however, is not concentrated in reserves: the checklist-weighted share inside them is 10.8%, indistinguishable from the area share. Matching each event to 200 control locations from the same province and year, weighted by the checklists recorded there, reduces the enrichment to an odds ratio of 1.71 (95% CI 1.39–2.11), and to 1.57 (1.26–1.96) after adding elevation, longitude and two bioclimatic covariates (Supplementary Note S6; Supplementary Fig. S12).

Three features argue that even this residual association is largely observational. It rests on few sites: the 172 inside-reserve events fall in 68 reserves, the top two holding 27.3%; dropping those two leaves 1.16 (0.92–1.47). It is confined to the period of rapid growth in birdwatching (2.05, 1.61–2.59, for 2013–2023 against 0.97, 0.62–1.50, for 2002–2012), although the mapped network gained no area after 2012. And checklists inside reserves record 18.9 species against 15.3 outside, a 24% difference in per-visit yield that stratification cannot remove. Because within-site observation bias cannot be conditioned out, we declared in advance how the result would be read: a null would be strong evidence against an ecological contribution, a positive one weak evidence for it. The 1.57 odds ratio is therefore an upper bound, and record counts inside reserves should not index reserve performance.

Two further results point the same way. Of all first records, 41.1% were re-detected in the same province at least once and 22.5% in three or more later years, so provincial first records are neither mostly vagrants nor mostly established colonisations; whether a record persists is unrelated to reserve status once later survey effort is controlled (odds ratios 1.01, 0.62–1.64, and 1.10, 0.61–1.99). And the network shows no measurable mismatch with observed warming: with warming recomputed at cell level from CRU TS 4.09, area-weighted coverage across warming quintiles spans only 8.5–11.8%, against the 8.5-fold range implied by the provincial field broadcast to cells, and a stratified permutation places the observed correlation inside the null (P = 0.48). What the network lacks is monitoring: the upper warming tercile crossed with the lower recent-effort tercile identifies 3.93×10^5 km² inside existing reserves, where monitoring could be grafted onto current management, and 1.60×10^6 km² outside any mapped reserve (Supplementary Figs S13, S14). All statements here apply only to the 1,028 mapped reserves established before 2012, and we make no causal claim.

Discussion

New provincial bird records in China reflect both the movement of birds and the growth of birdwatching, in similar measure. Once records are dated to discovery and observation is modelled explicitly, a standard deviation of warming relative to a species' historical range raises the annual hazard of a first record about as much as a standard deviation of survey effort (hazard ratios 1.36 and 1.40). Which of the two appears to dominate depends on a field that analyses of such compilations seldom question: the date. We had expected effort to dominate, given the growth of birdwatching and the association of new mammal records with survey effort (Ding et al., 2025); that expectation held only under publication-year dating.

Publication dating biases estimates in a predictable direction (H1). Publication lags discovery by about two years and observation grew every year, so a record dated to publication is credited to a year in which more people were looking than when the bird was found, and the error favours whichever covariate trends with observer numbers. The same shift detaches a record from the weather of its discovery year, which is how a spurious effect of annual climate variability arises. The problem extends beyond new records: most new plant species are described from specimens collected years or decades earlier (Bebber et al., 2010), and assessments of alien-species accumulation rest on dated first records (Seebens et al., 2017). Any analysis relating such series to trending covariates is exposed to the same bias. The first remedy is to record the date of observation, as CBNR does. Where that is impossible, a completeness offset like ours, the discrete-time counterpart of the reporting-delay adjustments used in epidemiological surveillance (Lawless, 1994), corrects the expected hazard for records not yet published.

The two drivers also behave differently through time. The return on effort fell from 2.18 in 2002 to 1.07 in 2024, and provinces that were well surveyed in 2002–2008 later yielded fewer records per unit of effort, whereas the return on warming showed no detectable change (H4). We read this as the difference between a stock and a flow. Effort draws down a stock of occurrences that were detectable but undocumented, the Wallacean shortfall; warming keeps adding to it as the climate of each province moves relative to species' ranges. The negative interaction (H3) fits the same reading: where effort has long been high, many of the species that warming favours have already been recorded, and the pairs left at risk respond less. Depletion of this kind is expected in hazard analyses, because the most susceptible units leave the risk set first (Hernán, 2010). Geography offers an alternative: sparsely surveyed provinces are disproportionately western and montane, where steep gradients may make a given amount of warming more consequential, and

this alone could produce a negative interaction. Nor did we test the two time trends against each other. If the stock-and-flow reading is correct, the share of new records that reflects redistribution will grow as the stock is exhausted, and new-record series will become more informative about climate, provided that their dating and the observation process are modelled.

The warming coefficient also needs the right benchmark. Permutations that retain provincial-annual climate but scramble species identity still return a hazard ratio of about 1.16, so the component attributable to referencing warming to each species' own range is about 1.18. The rest is an association between provincial climate and the number of records in a province-year, which may be climatic or observational and which a species-blind analysis would reproduce. Direction makes the same point more sharply: the apparent eastward concentration of new records is what China's position relative to species' global ranges produces when records fall at random among candidate provinces. Against a null that also follows survey effort, records lie further north than expected, with 40.2% in the three northern sectors against 33.0%. Directional readings of new records (Xing et al., 2024; Chen et al., 2025) therefore need a geometric and observational benchmark; against one, the poleward signal in these data is detectable but modest.

Projection meets a different limit. Because the species-referenced gradient varied little in the past, late-century scenarios push about 98% of species–province cells outside the fitted support, where the mechanistic and machine-learning models agree on the direction of change in fewer than a quarter of cells. The scenario projections also apply the effort coefficient estimated over the whole period, although its value had fallen to 1.07 by 2024, so the effort-driven rise they imply is an upper bound. Projections of record generation, and of other indicators built on species-specific climate differentials, should report the share of projected cells inside the fitted

support (Elith et al., 2010; Mesgaran et al., 2014; Yates et al., 2018). Where that share is small, a near-term surface with an explicit support mask is more defensible than a century-scale map.

Several limitations bound these conclusions. The events are published records, and the model has no explicit detection component, so the coefficients describe the generation of documented records; a species present but unreported is counted as absent, and no coefficient measures range expansion itself. The province-by-year intercept absorbs provincial climate anomalies as well as survey pulses, which protects the fixed effects from observational confounding but, if anything, biases the warming estimate towards zero. Effort is measured per province and year, far more coarsely than the sites observers visit, and the directional nulls likewise place records at province level. The candidate pool rests on species-distribution models whose thresholds we varied but whose structure we did not, and vagrants without established Chinese ranges were excluded, so the results describe dynamics at range margins. The most recent years remain right-censored: the completeness offset corrects the expected hazard but cannot recover unpublished records. The design is observational, and no coefficient should be read as causal.

A province-year pulse of one standard deviation multiplies the hazard of a first record by 2.23, more than either driver, so survey campaigns change the biodiversity record quickly and should be placed deliberately. Because effort acts on a logarithmic scale and well-surveyed provinces yield fewer records per unit of new effort, additional visits document most where surveys are sparse, which is also where warming raises the hazard most; existing reserves in the upper tercile of warming and the lower tercile of recent effort (3.93×10^5 km²) are a practical starting point. Because the return on effort has fallen within two decades, a plateau in new records should not be taken to mean that redistribution has slowed. Records inside nature reserves track where people watch birds and should not be used to score reserve performance. Finally, compilers of

new-record data can make every later analysis more reliable, at little cost, by publishing both the date of observation and the date of publication for each record.

Acknowledgements

We thank the birdwatchers who contributed observations to eBird and China Bird Report, and the authors of the publications compiled in CBNR. [Funding sources and grant numbers to be added by the authors.] The authors used Claude (Anthropic) to assist with drafting and editing text, checking numerical consistency and writing figure code; they reviewed and verified all content and take full responsibility for it.

Conflict of interest statement

The authors declare no conflicts of interest.

Data availability statement

The new-record data analysed in this study are version 2 of the China Bird New Record dataset (CBNR), openly available under a CC BY 4.0 licence in Zenodo at https://doi.org/10.5281/zenodo.20809949 (Ding et al., 2026). Nature-reserve boundaries are openly available in Zenodo at https://doi.org/10.5281/zenodo.14875797 (China Nature Reserve Specimen Resource Sharing Platform, 2024). Historical monthly climate (WorldClim 2.1 downscaling of CRU TS 4.09; Fick & Hijmans, 2017; Harris et al., 2020) and CMIP6 projections are publicly available from WorldClim (https://www.worldclim.org). Species range maps (BirdLife International & Handbook of the Birds of the World, 2024) are available on request from BirdLife International (http://datazone.birdlife.org/species/requestdis); their licence does not permit redistribution. Survey-effort indicators were derived from eBird/GBIF occurrence

records and China Bird Report checklists (http://www.birdreport.cn); the raw records remain subject to their providers' terms, and the province-year effort panel is provided in the project repository. All analysis code (R 4.5.1), the risk set, the derived climate and effort panels and every result table are available at https://github.com/dingchenchen6/bird-newrecord-discovery-dated (Ding, 2026) and will be archived in Zenodo with a DOI upon acceptance; the model equations and all parameter estimates are given in Materials and Methods and Supporting Information.

References

Allison, P. D. Discrete-time methods for the analysis of event histories. *Sociol. Methodol.* **13**, 61–98 (1982). https://doi.org/10.2307/270718

Bebber, D. P. et al. Herbaria are a major frontier for species discovery. *Proc. Natl Acad. Sci. USA* **107**, 22169–22171 (2010). https://doi.org/10.1073/pnas.1011841108

BirdLife International & Handbook of the Birds of the World. Bird species distribution maps of the world, version 2024. http://datazone.birdlife.org/species/requestdis (2024).

Boakes, E. H. et al. Distorted views of biodiversity: spatial and temporal bias in species occurrence data. *PLoS Biol.* **8**, e1000385 (2010). https://doi.org/10.1371/journal.pbio.1000385

Bowler, D. E. et al. Treating gaps and biases in biodiversity data as a missing data problem. *Biol. Rev.* **100**, 50–67 (2025). https://doi.org/10.1111/brv.13127

Brooks, M. E. et al. glmmTMB balances speed and flexibility among packages for zero-inflated generalized linear mixed modeling. *R J.* **9**, 378–400 (2017). https://doi.org/10.32614/RJ-2017-066

673 Chen, I.-C., Hill, J. K., Ohlemüller, R., Roy, D. B. & Thomas, C. D. Rapid range shifts of
674 species associated with high levels of climate warming. *Science* **333**, 1024–1026 (2011).
675 https://doi.org/10.1126/science.1206432

676 Chen, S. et al. Chinese provincial-level new records for 96 resident bird species reveal poleward
677 range shifts. *Avian Res.* **16**, 100310 (2025). https://doi.org/10.1016/j.avrs.2025.100310

678 Chen, T. & Guestrin, C. XGBoost: a scalable tree boosting system. In *Proc. 22nd ACM SIGKDD*
679 *Int. Conf. Knowledge Discovery and Data Mining* 785–794 (2016).
680 https://doi.org/10.1145/2939672.2939785

681 China Nature Reserve Specimen Resource Sharing Platform. List and vector boundaries of
682 nature reserves in China. Zenodo https://doi.org/10.5281/zenodo.14875797 (2024).

683 Ding, C. Analysis code for 'Warming and survey effort contribute comparably to new bird
684 distribution records when dated to discovery'. GitHub https://github.com/dingchenchen6/bird-
685 newrecord-discovery-dated (2026).

686 Ding, C. et al. A dataset of provincial-level new distribution records for birds in China from
687 2000 to 2025. *Sci. Data* (in review).

688 Ding, C., Ding, J., Qiao, H., Jiang, Z. & Wang, Z. Taxonomic and spatiotemporal patterns and
689 ecological correlates of new mammal distribution records in China. *Glob. Ecol. Biogeogr.* **34**,
690 e70165 (2025). https://doi.org/10.1111/geb.70165

691 Ding, C., Zhao, H., Wang, Z., Li, L., Ding, J., Jiang, Z., Zhang, Z. & Wang, Z. A dataset of
692 provincial-level new distribution records for birds in China from 2000 to 2025 (China Bird New
693 Record dataset, CBNR), version 2. Zenodo https://doi.org/10.5281/zenodo.20809949 (2026).

694 Diniz-Filho, J. A. F. et al. Macroecological links between the Linnean, Wallacean, and
695 Darwinian shortfalls. *Front. Biogeogr.* **15**, e59566 (2023). https://doi.org/10.21425/F5FBG59566

696 Elith, J., Kearney, M. & Phillips, S. The art of modelling range-shifting species. *Methods Ecol.*
697 *Evol.* **1**, 330–342 (2010). https://doi.org/10.1111/j.2041-210X.2010.00036.x

698 Fick, S. E. & Hijmans, R. J. WorldClim 2: new 1-km spatial resolution climate surfaces for
699 global land areas. *Int. J. Climatol.* **37**, 4302–4315 (2017). https://doi.org/10.1002/joc.5086

700 Harris, I., Osborn, T. J., Jones, P. & Lister, D. Version 4 of the CRU TS monthly high-resolution
701 gridded multivariate climate dataset. *Sci. Data* **7**, 109 (2020). https://doi.org/10.1038/s41597-
702 020-0453-3

703 Hartig, F. DHARMa: residual diagnostics for hierarchical (multi-level/mixed) regression models.
704 R package version 0.4 (2024). https://CRAN.R-project.org/package=DHARMa

705 Hernán, M. A. The hazards of hazard ratios. *Epidemiology* **21**, 13–15 (2010).
706 https://doi.org/10.1097/EDE.0b013e3181c1ea43

707 Hortal, J. et al. Seven shortfalls that beset large-scale knowledge of biodiversity. *Annu.*
708 *Rev. Ecol. Evol. Syst.* **46**, 523–549 (2015). https://doi.org/10.1146/annurev-ecolsys-112414-
709 054400

710 Hughes, A. C. et al. Sampling biases shape our view of the natural world. *Ecography* **44**, 1259–
711 1269 (2021). https://doi.org/10.1111/ecog.05926

712 Isaac, N. J. B., van Strien, A. J., August, T. A., de Zeeuw, M. P. & Roy, D. B. Statistics for
713 citizen science: extracting signals of change from noisy ecological data. *Methods Ecol. Evol.* **5**,
714 1052–1060 (2014). https://doi.org/10.1111/2041-210X.12254

715 Lawless, J. F. Adjustments for reporting delays and the prediction of occurred but not reported
716 events. *Can. J. Stat.* **22**, 15–31 (1994). https://doi.org/10.2307/3315820

717 Lenoir, J. & Svenning, J.-C. Climate-related range shifts – a global multidimensional synthesis
718 and new research directions. *Ecography* **38**, 15–28 (2015). https://doi.org/10.1111/ecog.00967

719 Lundberg, S. M. et al. From local explanations to global understanding with explainable AI for
720 trees. *Nat. Mach. Intell.* **2**, 56–67 (2020). https://doi.org/10.1038/s42256-019-0138-9

721 Mesgaran, M. B., Cousens, R. D. & Webber, B. L. Here be dragons: a tool for quantifying
722 novelty due to covariate range and correlation change when projecting species distribution
723 models. *Divers. Distrib.* **20**, 1147–1159 (2014). https://doi.org/10.1111/ddi.12209

724 Moura, M. R. & Jetz, W. Shortfalls and opportunities in terrestrial vertebrate species discovery.
725 *Nat. Ecol. Evol.* **5**, 631–639 (2021). https://doi.org/10.1038/s41559-021-01411-5

726 Nakagawa, S. & Schielzeth, H. A general and simple method for obtaining R^2 from generalized
727 linear mixed-effects models. *Methods Ecol. Evol.* **4**, 133–142 (2013).
728 https://doi.org/10.1111/j.2041-210x.2012.00261.x

729 Oliver, R. Y., Meyer, C., Ranipeta, A., Winner, K. & Jetz, W. Global and national trends, gaps,
730 and opportunities in documenting and monitoring species distributions. *PLoS Biol.* **19**, e3001336
731 (2021). https://doi.org/10.1371/journal.pbio.3001336

732 Parmesan, C. & Yohe, G. A globally coherent fingerprint of climate change impacts across
733 natural systems. *Nature* **421**, 37–42 (2003). https://doi.org/10.1038/nature01286

734 Pecl, G. T. et al. Biodiversity redistribution under climate change: impacts on ecosystems and
735 human well-being. *Science* **355**, eaai9214 (2017). https://doi.org/10.1126/science.aai9214

736 Seebens, H. et al. No saturation in the accumulation of alien species worldwide. *Nat. Commun.* **8**,
737 14435 (2017). https://doi.org/10.1038/ncomms14435

738 Singer, J. D. & Willett, J. B. It's about time: using discrete-time survival analysis to study
739 duration and the timing of events. *J. Educ. Stat.* **18**, 155–195 (1993).
740 https://doi.org/10.3102/10769986018002155

741 Tingley, M. W. & Beissinger, S. R. Detecting range shifts from historical species occurrences:
742 new perspectives on old data. *Trends Ecol. Evol.* **24**, 625–633 (2009).
743 https://doi.org/10.1016/j.tree.2009.05.009

744 Tobias, J. A. et al. AVONET: morphological, ecological and geographical data for all birds.
745 *Ecol. Lett.* **25**, 581–597 (2022). https://doi.org/10.1111/ele.13898

746 Xing, X. et al. Where are the provincial-level new records in China from the past 20 years, and
747 which traits determine their shift directions? *Front. Ecol. Evol.* **12**, 1415268 (2024).
748 https://doi.org/10.3389/fevo.2024.1415268

749 Yates, K. L. et al. Outstanding challenges in the transferability of ecological models. *Trends*
750 *Ecol. Evol.* **33**, 790–802 (2018). https://doi.org/10.1016/j.tree.2018.08.001

751 ## Tables

752 **Table 1.** Step-by-step decomposition from the previously published specification to the corrected
753 main model.

Step	Change	*n*	Events	HR effort	HR warming	HR variability	HR interaction
S0	previously published	182,485	655	1.788	1.394	0.850	0.876
S1	+ discovery-year dating	181,134	646	1.305	1.437	0.971	0.859

Step	Change	*n*	Events	HR effort	HR warming	HR variability	HR interaction
S2	+ coverage-gap effort	175,686	638	1.267	1.432	0.958	0.863
S3	+ area-weighted climate	175,901	649	1.289	1.432	0.958	0.863
S4	+ completeness offset	175,901	649	1.448	1.428	0.962	0.862
S5	+ province × year effect	175,901	649	**1.404**	**1.362**	**0.995**	**0.849**

Table 2. Main model. Discrete-time proportional hazards, complementary log-log link; n = 175,901; 649 events; 392 species; 31 provinces; AIC 8,313.2. Cluster bootstrap: 300 resamples of provinces. Permutation null medians are from the strongest constrained null (N3, events shuffled among the species at risk within each province-year) for the climate terms and the interaction, and from N2 (effort shuffled across years within province) for effort; 199 permutations each. Because these nulls preserve the provincial-annual structure, their medians rather than unity are the correct reference for the corresponding effect sizes.

Term	HR per SD	95% CI (Wald)	95% CI (province cluster bootstrap)	Permutation null median	*P*
Accumulated warming	1.362	1.216–1.524	1.20–1.72	1.157	8.3×10^{-8}
Survey effort	1.404	1.245–1.583	1.21–1.76	0.980	2.9×10^{-8}
Annual climate variability	0.995	0.909–1.090	0.90–1.13	0.955	0.92
Warming × effort	0.849	0.761–0.948	0.74–0.99	0.955	3.7×10^{-3}
σ species	0.405				
σ province	0.326				
σ province × year	0.804				

Table 3. Random-effect structure comparison on four criteria.

Structure	ΔAIC	McFadden R^2	Marginal R^2	Conditional R^2	Marginal AUC	Conditional AUC	ICC
species	163.1	0.0142	0.108	0.183	0.613	0.687	0.084
species + province	98.7	0.0220	0.124	0.274	0.612	0.721	0.172
+ year	74.8	0.0250	0.069	0.277	0.607	0.737	0.224

Structure	ΔAIC	McFadden R^2	Marginal R^2	Conditional R^2	Marginal AUC	Conditional AUC	ICC
+ province × year (main)	**5.25**	**0.0331**	**0.082**	**0.411**	**0.612**	**0.854**	**0.358**
+ species warming slope	4.48	0.0334	0.080	0.403	0.612	0.865	0.351
+ province effort slope	0.00	0.0339	0.082	0.395	0.611	0.852	0.341

Figure legends




Figure 1 | Survey effort and accumulated warming contribute comparably to the hazard of a new provincial record. a, Hazard ratios per standard deviation with 95% Wald confidence intervals, for the four fixed effects, at three species-distribution-model thresholds (main threshold: n = 175,901 species × province × year cells, 649 events). The dashed line marks unity. Note that unity is not the appropriate null for the warming terms: design-preserving permutation nulls that retain the provincial-annual climate and event structure are centred on 1.16 for

accumulated warming and 0.96 for the interaction (Supplementary Fig. S11a), so the increment attributable to species-specific referencing is smaller than the distance from unity suggests. **b**, Marginal predictions of the annual hazard against accumulated warming at the 10th, 50th and 90th percentiles of survey effort; ribbons are 95% confidence intervals on the fixed-effect linear predictor. **c**, Relative importance of each term under four independent criteria: loss of AIC, loss of fixed-effect AUC, loss of conditional R^2, and the absolute standardised coefficient. Larger values indicate greater importance. **d**, Discrimination (area under the ROC curve, AUC) attributable to fixed effects alone as terms are added, with the random structure held at the main model.




Figure 2 | Dating records to publication rather than to discovery is a systematic source of bias. a, Distribution of the delay between discovery and publication (bars, left axis as per cent of the 1,028 records with a parseable discovery year) and the resulting reporting completeness $c(t)$

implied by a compilation extending to 2025 publications (line). **b**, Annual counts of new provincial records under the two dating conventions. The shaded band marks years in which the discovery-year series is right-censored because the corresponding papers have not yet appeared. **c**, Hazard ratios (point estimates) for the four fixed effects across the five-step decomposition from the previously published specification (S0) to the corrected main model (S5). The shaded column marks the re-dating step.




Figure 3 | The result does not depend on any freely chosen analytical decision. a, Hazard ratio for accumulated warming across four climate indicators and four accumulation windows; asterisks denote two-sided Wald $P < 0.05$ (*), < 0.01 (**) ***and* < 0.001** (***). **b**, ΔAIC across the same grid, all cells fitted on identical rows. **c**, Hazard ratios with 95% Wald confidence intervals for survey effort (triangles) and accumulated warming (circles) across four effort proxies and three

treatments of the 24 implausible zero cells. AIC is not comparable across missing-data rules because the sample sizes differ, so only coefficients are shown.




Figure 4 | Anatomy of the model, in units a reader can picture. a, Construction of the species-referenced climate gradient, shown as 15-year trailing means for one province and two species. Yunnan has warmed more than the Chinese range of *Aethopyga christinae* and less than that of *Coracias garrulus*, so the same province in the same year yields opposite values of x for the two species. **b**, Variance of each climate term partitioned into variation between province-years and variation within a province-year, the latter arising purely from differences among species. **c**, Partial effects in natural units: annual hazard against how much more the province has warmed than the species' range, at three levels of survey effort expressed as annual visits. **d**,

Every variance component and both focal fixed effects expressed as the hazard multiplier per one standard deviation, on a common axis.




Figure 5 | The observation process made visible. a, Species random intercepts as hazard multipliers (n = 392 species), grouped by migratory strategy; diamonds are group medians. **b**, Province random intercepts mapped, after controlling for effort and climate; grey denotes provincial units outside the analysis scope. Base map GS(2019)1822. **c**, Province-by-year random intercepts, the largest variance component in the model. Provinces are ordered by their mean. **d**, Hazard ratio for survey effort as a function of year, from the model that allows effects to vary through time; the ribbon is the 95% confidence interval (effort × time interaction, Wald P

= 7.7×10^{-4}). Map lines delineate study areas and do not necessarily depict accepted national boundaries. Base map: standard map GS(2019)1822.




Figure 6 | Direction of new records relative to species' range centroids, against geometric and observational nulls. a, Share of the 642 records with a range centroid in each of eight 45° sectors of bearing from the centroid of the species' global range (solid line), with the shares expected if each record fell at random among its species' candidate provinces (dashed;

geometry) or among them in proportion to their 2002–2024 survey visits (dotted; geometry + effort), from 999 Monte Carlo draws. **b**, Ratio of observed to expected records per sector under the two nulls (log scale); bars show the 95% Monte Carlo interval of the null expressed as a ratio, and filled points mark sectors with Holm-adjusted two-sided $P < 0.05$. Against a uniform expectation direction departs strongly ($\chi^2 = 580.1$, df = 7, $P = 4.6 \times 10^{-121}$); against the geometric null only weakly ($\chi^2 = 10.8$, Monte Carlo $P = 0.039$); against the geometry + effort null the departure points north ($\chi^2 = 25.8$, $P = 0.001$). **c**, Observed shares for the six orders with most records; each polygon gives the share of that order's own records per sector, and the key gives the numbers of records and species, because one species can contribute records in several provinces. Wind roses for the sixteen best-sampled orders are provided in the Supplementary Information.




Figure 7 | The two model families agree inside the fitted range and diverge outside it. a, Province-year mean predicted hazard, 2002–2024, from the mechanistic and gradient-boosted models (Spearman $\rho = 0.54$). **b**, First-to-99th-percentile range of each predictor in the fitted data and in each scenario-horizon combination. **c**, Province-level projected hazard ratios, with rank correlation and directional agreement per panel.