

# ROSC: Robust Spectral Clustering on Multi-scale Data

Anonymous Author(s)

This technical report proves the grouping effect of  $|Z^*|, |(Z^*)^T|$  and  $\tilde{Z}$ . In the following discussion, we use  $z_q^*$  to denote the  $q$ -th column vector of  $Z^*$ .

LEMMA 1. *Given a set of objects  $X$ , the matrix  $X \in \mathbb{R}^{p \times n}$  that is composed of the pseudo-eigenvectors as rows, the reachability matrix  $\mathcal{W}$ , and the optimal solution  $Z^*$ ,*

$$Z_{ip}^* = \frac{\mathbf{x}_i^T (\mathbf{x}_p - Xz_p^*) + \alpha_2 \mathcal{W}_{ip}}{\alpha_1 + \alpha_2}, \quad \forall 1 \leq i, p \leq n. \quad (1)$$

PROOF. For  $1 \leq p \leq n$ , let  $J(z_p) = \|\mathbf{x}_p - Xz_p\|_2^2 + \alpha_1 \|z_p\|_2^2 + \alpha_2 \|\mathbf{z}_p - \mathbf{w}_p\|_2^2$ . Since  $Z^*$  is the optimal solution, we have  $\frac{\partial J}{\partial Z_{ip}}|_{z_p=Z_{ip}^*} = 0 \quad \forall 1 \leq i \leq n$ . Hence,  $-2\mathbf{x}_i^T (\mathbf{x}_p - Xz_p^*) + 2\alpha_1 Z_{ip}^* + 2\alpha_2 (Z_{ip}^* - \mathcal{W}_{ip}) = 0$ , which induces Equation 1.  $\square$

LEMMA 2.  $\forall 1 \leq i, j, p \leq n$ ,

$$|Z_{ip}^* - Z_{jp}^*| \leq \frac{c\sqrt{2(1-r)} + \alpha_2 |\mathcal{W}_{ip} - \mathcal{W}_{jp}|}{\alpha_1 + \alpha_2}, \quad (2)$$

where  $c = \sqrt{1 + \alpha_2 \|\mathbf{w}_p\|_2^2}$  and  $r = \mathbf{x}_i^T \mathbf{x}_j$ .

PROOF. From Equation 1, we have

$$Z_{ip}^* - Z_{jp}^* = \frac{(\mathbf{x}_i^T - \mathbf{x}_j^T)(\mathbf{x}_p - Xz_p^*) + \alpha_2 (\mathcal{W}_{ip} - \mathcal{W}_{jp})}{\alpha_1 + \alpha_2}.$$

That implies

$$\begin{aligned} |Z_{ip}^* - Z_{jp}^*| &\leq \frac{|(\mathbf{x}_i^T - \mathbf{x}_j^T)(\mathbf{x}_p - Xz_p^*)| + \alpha_2 |\mathcal{W}_{ip} - \mathcal{W}_{jp}|}{\alpha_1 + \alpha_2} \\ &\leq \frac{\|(\mathbf{x}_i^T - \mathbf{x}_j^T)\|_2 \|\mathbf{x}_p - Xz_p^*\|_2 + \alpha_2 |\mathcal{W}_{ip} - \mathcal{W}_{jp}|}{\alpha_1 + \alpha_2} \end{aligned} \quad (3)$$

Since the column vectors of  $X$  are normalized (i.e.,  $\mathbf{x}_q^T \mathbf{x}_q = 1 \quad \forall 1 \leq q \leq n$ ), we have  $\|(\mathbf{x}_i^T - \mathbf{x}_j^T)\|_2 = \sqrt{2(1-r)}$ , where  $r = \mathbf{x}_i^T \mathbf{x}_j$ . As  $Z^*$  is the optimal solution, we have

$$\begin{aligned} J(z_p^*) &= \|\mathbf{x}_p - Xz_p^*\|_2^2 + \alpha_1 \|z_p^*\|_2^2 + \alpha_2 \|z_p^* - \mathbf{w}_p\|_2^2 \leq \\ J(\mathbf{0}) &= \|\mathbf{x}_p\|_2^2 + \alpha_2 \|\mathbf{w}_p\|_2^2 = 1 + \alpha_2 \|\mathbf{w}_p\|_2^2. \end{aligned} \quad (4)$$

Hence,  $\|\mathbf{x}_p - Xz_p^*\|_2 \leq \sqrt{1 + \alpha_2 \|\mathbf{w}_p\|_2^2} = c$ . Equation 3 can be further simplified as

$$|Z_{ip}^* - Z_{jp}^*| \leq \frac{c\sqrt{2(1-r)} + \alpha_2 |\mathcal{W}_{ip} - \mathcal{W}_{jp}|}{\alpha_1 + \alpha_2}.$$

$\square$

Permission to make digital or hard copies of part or all of this work for personal or classroom use is granted without fee provided that copies are not made or distributed for profit or commercial advantage and that copies bear this notice and the full citation on the first page. Copyrights for third-party components of this work must be honored. For all other uses, contact the owner/author(s).  
Conference'17, July 2017, Washington, DC, USA  
© 2017 Copyright held by the owner/author(s).  
ACM ISBN 978-x-xxxx-xxxx-x/YY/MM.  
<https://doi.org/10.1145/nnnnnnnn.nnnnnnnn>

LEMMA 3.  $Z^*$  and  $|Z^*|$  have grouping effect.

PROOF. Given two objects  $x_i$  and  $x_j$  such that  $x_i \rightarrow x_j$ , we have, (1)  $\mathbf{x}_i^T \mathbf{x}_j \rightarrow 1$  and (2)  $\|\mathbf{w}_i - \mathbf{w}_j\|_2 \rightarrow 0$ . These imply  $r = \mathbf{x}_i^T \mathbf{x}_j \rightarrow 1$  and  $|\mathcal{W}_{ip} - \mathcal{W}_{jp}| \rightarrow 0$ . Hence, the two terms of the numerator of the R.H.S of Equation 2 are close to 0. Therefore,  $|Z_{ip}^* - Z_{jp}^*| \rightarrow 0$  and thus  $Z^*$  has grouping effect. Further, we have  $||Z_{ip}^*| - |Z_{jp}^*|| \leq |Z_{ip}^* - Z_{jp}^*|$ , so  $||Z_{ip}^*| - |Z_{jp}^*|| \rightarrow 0$  and  $|Z^*|$  has grouping effect.  $\square$

LEMMA 4.  $(Z^*)^T$  and  $|(Z^*)^T|$  have grouping effect.

PROOF. The problem is equivalent to if  $x_i \rightarrow x_j$ ,  $Z_{pi}^* \rightarrow Z_{pj}^*$ . From Lemma 1,

$$z_{pi}^* - z_{pj}^* = \frac{\mathbf{x}_p^T (\mathbf{x}_i - \mathbf{x}_j - X(z_i^* - z_j^*)) + \alpha_2 (\mathcal{W}_{pi} - \mathcal{W}_{pj})}{\alpha_1 + \alpha_2}.$$

Since  $z_i^* = (X^T X + \alpha_1 I + \alpha_2 I)^{-1} (X^T \mathbf{x}_i + \alpha_2 \mathbf{w}_i)$ ,  $z_j^* = (X^T X + \alpha_1 I + \alpha_2 I)^{-1} (X^T \mathbf{x}_j + \alpha_2 \mathbf{w}_j)$ , let  $Y = X(X^T X + \alpha_1 I + \alpha_2 I)^{-1}$ . Then

$$\begin{aligned} |Z_{pi}^* - Z_{pj}^*| &\leq \frac{||\mathbf{x}_p^T (\mathbf{x}_i - \mathbf{x}_j)||_2 + ||\mathbf{x}_p^T X(z_i^* - z_j^*)||_2 + \alpha_2 |\mathcal{W}_{pi} - \mathcal{W}_{pj}|}{\alpha_1 + \alpha_2} \\ &\leq \frac{||\mathbf{x}_p^T (\mathbf{x}_i - \mathbf{x}_j)||_2 + ||\mathbf{x}_p^T Y X^T (\mathbf{x}_i - \mathbf{x}_j)||_2}{\alpha_1 + \alpha_2} \\ &\quad + \frac{\alpha_2 ||\mathbf{x}_p^T Y (\mathbf{w}_i - \mathbf{w}_j)||_2 + \alpha_2 |\mathcal{W}_{pi} - \mathcal{W}_{pj}|}{\alpha_1 + \alpha_2} \end{aligned} \quad (5)$$

If  $x_i \rightarrow x_j$ , i.e.,  $\mathbf{x}_i^T \mathbf{x}_j \rightarrow 1$  and  $\|\mathbf{w}_i - \mathbf{w}_j\|_2 \rightarrow 0$ , we have  $\|\mathbf{x}_i - \mathbf{x}_j\|_2 \rightarrow 0$  and  $|\mathcal{W}_{pi} - \mathcal{W}_{pj}| \rightarrow 0$ . Then  $|Z_{pi}^* - Z_{pj}^*| \rightarrow 0$  and  $(Z^*)^T$  has grouping effect. Since  $||Z_{pi}^*| - |Z_{pj}^*|| \leq |Z_{pi}^* - Z_{pj}^*|$ ,  $|(Z^*)^T|$  also has grouping effect.  $\square$

LEMMA 5. Matrix  $\tilde{Z}$  has grouping effect.

PROOF. From Lemma 3 and 4, both  $|Z^*|$  and  $|(Z^*)^T|$  have the grouping effect. Since  $\tilde{Z} = (|Z^*| + |(Z^*)^T|)/2$ ,

$$\begin{aligned} |\tilde{Z}_{ip} - \tilde{Z}_{jp}| &= \frac{|(|Z_{ip}^*| + |Z_{pi}^*|) - (|Z_{jp}^*| + |Z_{pj}^*|)|}{2} \\ &\leq \frac{|Z_{ip}^* - Z_{jp}^*| + |Z_{pi}^* - Z_{pj}^*|}{2} \end{aligned} \quad (6)$$

If  $x_i \rightarrow x_j$ , both  $|Z_{ip}^* - Z_{jp}^*| \rightarrow 0$  and  $|Z_{pi}^* - Z_{pj}^*| \rightarrow 0$ , so  $|\tilde{Z}_{ip} - \tilde{Z}_{jp}| \rightarrow 0$  and  $\tilde{Z}$  has grouping effect.  $\square$