
# Homework 2 R Problem

```
set.seed(2)
```

### a) ACF

```
# a) AR(1) series with phi = 0.5
y.50 <- arima.sim(model = list(ar = 0.5), n = 100)
# plot ACF of AR(1) series with phi = 0.5
acf(y.50)</pre>
```



```
y_acf_values <- acf(y.50, plot = FALSE)
y_acf_values$acf</pre>
```


```
## , , 1
##
## [,1]
## [1,] 1.000000000
```

```
[2,] 0.456407732
##
##
    [3,] 0.177230916
    [4,] -0.071981494
    [5,] -0.111073841
##
##
    [6,] -0.079874933
##
    [7,] -0.058616143
##
    [8,] -0.218855549
    [9,] -0.138868166
##
##
   [10,] -0.072900659
   [11,]
         0.004573513
##
   [12,]
          0.024174182
   [13,]
          0.048179630
   [14,]
          0.082197678
##
  [15,]
          0.137261651
## [16,]
          0.105199109
## [17,]
          0.072913649
## [18,] -0.028337238
## [19,] -0.073045319
## [20,] -0.047920471
## [21,] -0.065887861
```

Comparing the ACF of phi = 0.5 with answer from Exercise 3.1, both have similar values.

### **PACF**

```
# plot PACF of AR(1) series with phi = 0.5
acf(y.50, type = "partial")
```




```
y_pacf_values <- acf(y.50, plot = FALSE, type = "partial")
y_pacf_values$acf</pre>
```

```
, , 1
##
##
                 [,1]
##
  [1,] 0.456407732
  [2,] -0.039254032
## [3,] -0.174744252
##
   [4,] -0.007050781
## [5,] 0.011324608
## [6,] -0.041286401
##
   [7,] -0.255048863
## [8,] 0.068537680
## [9,] 0.027009750
## [10,] -0.037883358
## [11,] -0.030948631
## [12,] 0.050008586
## [13,] 0.083653500
## [14,] 0.021588056
## [15,] -0.004644291
## [16,] 0.041094359
## [17,] -0.052483404
## [18,] -0.041966874
## [19,] 0.041631822
## [20,] -0.039006680
```

The PACF with k > 2 are equal to 0 from Exercise 3.1. Here, it's slightly more or less than 0 when k > 2, which is still relatively similar to answer from Exercise 3.1.

### b) ACF

```
# a) AR(1) series with phi = 0.98
y.98 <- arima.sim(model = list(ar = -0.98), n = 100)
# plot ACF of AR(1) series with phi = 0.98
acf(y.98)</pre>
```




```
y_acf_values <- acf(y.98, plot = FALSE)
y_acf_values$acf</pre>
```

```
## , , 1
##
##
                [,1]
   [1,] 1.00000000
   [2,] -0.97677891
   [3,] 0.94996095
   [4,] -0.91377189
   [5,] 0.87719861
##
   [6,] -0.83490631
   [7,] 0.78355674
   [8,] -0.73174023
   [9,] 0.67866957
## [10,] -0.62207870
## [11,] 0.56073667
## [12,] -0.49430312
## [13,] 0.43098663
## [14,] -0.37626696
## [15,] 0.32456442
## [16,] -0.27387787
## [17,] 0.21991694
## [18,] -0.16794373
## [19,] 0.11717676
## [20,] -0.07042299
## [21,] 0.02105614
```

The ACF for phi = 0.98 in R, the values are also similar to Exercise 3.1.

#### **PACF**

```
# plot PACF of AR(1) series with phi = 0.98
acf(y.98, type = "partial")
```



```
y_pacf_values <- acf(y.98, plot = FALSE, type = "partial")
y_pacf_values$acf</pre>
```

```
##
##
##
                  [,1]
##
   [1,] -0.9767789129
    [2,] -0.0901051283
    [3,] 0.2136131363
##
   [4,] -0.0002978241
##
   [5,] 0.1107155689
    [6,] -0.2240810830
##
   [7,] -0.0206345138
##
   [8,] 0.0080086916
## [9,] 0.1329421091
## [10,] -0.1073070447
## [11,] 0.0926112262
## [12,] 0.0437971662
```

```
## [13,] -0.1956436941

## [14,] 0.0280227474

## [15,] 0.0552841946

## [16,] -0.1198905029

## [17,] 0.0641688190

## [18,] -0.0649022461

## [20,] -0.0923450575
```

The values in R here are also the same as PACF in Exercise  $3.1\mathrm{b}$