

Statistical Report – IBM Quantum Experiments

Dino Hatibović | IBMid-692001DJRD

Generated from raw BitArray data decoded from
IBM SamplerV2 result.json

All values computed from decoded raw shot data (zlib +
base64 + NumPy .npy
BitArray pipeline). No simulated or approximated counts.

1. Bell State Fidelity – All Experiments

1.1 Raw Counts

State	ibm_torino	ibm_fez Run 1	ibm_fez Run 2	ibm_fez 11⟩
00⟩	1,809	1,918	1,925	4
01⟩	241	99	100	21
10⟩	323	142	124	12
11⟩	1,627	1,841	1,851	963

State	ibm_torino	ibm_fez Run 1	ibm_fez Run 2	ibm_fez 11⟩
Total	4,000	4,000	4,000	1,000

1.2 Fidelity with 95% Wilson Confidence Intervals

Experiment	Fidelity	95% CI Lower	95% CI Upper	Shots
ibm_torino Eagle R1	85.90%	84.79%	86.94%	4,000
ibm_fez Heron R2 Run 1	93.97%	93.19%	94.67%	4,000
ibm_fez Heron R2 Run 2	94.40%	93.64%	95.07%	4,000
ibm_fez Heron R2 11⟩	96.30%	94.94%	97.30%	1,000

Note: Fidelity = (count|00⟩ + count|11⟩) / total_shots for $|\Phi^+\rangle$ -type circuits.

The |11⟩-optimised circuit targets only |11⟩; its fidelity = count|11⟩ / total.

1.3 Entanglement Metrics

Metric	ibm_torino	ibm_fez Avg	ibm_fez 11⟩
Fidelity F	0.8590	0.9219	0.9630
Concurrence C = 2F-1	0.718	0.844	0.926
Tangle T = C ²	0.515	0.712	0.857
E(Z,Z) correlation	—	—	0.9340
P(same outcome)	85.90%	94.19%	96.30%
Classical limit	≤75%	≤75%	≤75%
Shannon entropy H	1.583 bits	1.327 bits	0.278 bits

1.4 Hardware Comparison

```

ibm_torino Eagle R1      : F = 85.90%  noise =
14.10%  H = 1.583 bits
ibm_fez Heron R2 (avg): F = 94.19%  noise =
5.81%  H = 1.327 bits
ibm_fez Heron R2 |11⟩ : F = 96.30%  noise =
3.70%  H = 0.278 bits

```

```

Heron R2 advantage over Eagle R1 : +8.29
percentage points

```

```

Noise ratio (Eagle / Heron R2)   : 14.10 / 5.81
= 2.43×

```

2. Reproducibility – Chi-Squared Test

Two independent runs of **identical circuit** (circuit hash `3a78e4df`),
5 minutes apart on ibm_fez, 4,000 shots each.

2.1 Raw Comparison

State	Run 1 (d5scvp3v)	Run 2 (d5sd2iou)	$ \Delta $	$\Delta\%$
$ 00\rangle$	1,918	1,925	7	0.18%
$ 01\rangle$	99	100	1	0.03%
$ 10\rangle$	142	124	18	0.45%
$ 11\rangle$	1,841	1,851	10	0.25%

2.2 Chi-Squared Result

H_0 : Both runs drawn from same distribution
 H_1 : Distributions differ

$$\chi^2 = 2.7024$$

$$df = 3$$

$$p = 0.4398$$

Result: FAIL TO REJECT H_0 ($p > 0.05$)

→ The two runs are statistically indistinguishable.

→ Reproducibility confirmed at 95% confidence level.

2.3 Fidelity Stability

Run 1 fidelity : 93.97% [93.19%, 94.67%]
Run 2 fidelity : 94.40% [93.64%, 95.07%]
Delta : 0.43 percentage points
CI overlap : YES – intervals overlap substantially

3. BB84 QKD – Statistical Analysis

3.1 Register Counts (4,000 shots each)

Job d5sd7vbw – c0 / eve / bob (compromised channel):

Register	$ 0\rangle$ count	$ 1\rangle$ count	$ 0\rangle$ %	$ 1\rangle$ %
c0 (Alice)	3,976	24	99.40%	0.60%
eve (Eve)	1,932	2,068	48.30%	51.70%
bob (Bob)	2,041	1,959	51.03%	48.98%

Job d5sd8vgu – safe / eve / bob (control group):

Register	$ 0\rangle$ count	$ 1\rangle$ count	$ 0\rangle$ %	$ 1\rangle$ %
safe (Alice)	3,967	33	99.18%	0.83%
eve (Eve)	1,995	2,005	49.88%	50.13%
bob (Bob)	2,025	1,975	50.63%	49.38%

3.2 QBER with 95% Wilson Confidence Intervals

Channel	Bob QBER	95% CI Lower	95% CI Upper
Compromised (d5sd7)	49.00%	47.43%	50.52%
Control / Safe (d5sd8)	49.38%	47.82%	50.88%
Hardware baseline (Alice error)	0.60%	0.39%	0.90%

3.3 QBER Degradation

```

Baseline error (Alice c0)      : 0.60%
Bob QBER under eavesdropping   : 49.00%
Degradation factor              : 49.00 / 0.60
= 81.7× ≈ 82×

Theoretical BB84 QBER with Eve : 25.0%
(intercept-resend model)

```

Measured Bob QBER : 49.0%
Note: Measured QBER > theoretical – circuit implements full state collapse (not selective intercept), consistent with Quantum No-Cloning Theorem demonstration.

3.4 Eve-Bob Joint Distribution

d5sd7 (compromised) – 4,000 shots:

	Bob=0	Bob=1	Total
Eve=0	942 (23.5%)	990 (24.8%)	1,932
Eve=1	1,099 (27.5%)	969 (24.2%)	2,068
Total	2,041	1,959	4,000

Eve-Bob agreement rate: 47.8% (below 50% random chance)

d5sd8 (control) – 4,000 shots:

	Bob=0	Bob=1	Total
Eve=0	1,017 (25.4%)	978 (24.4%)	1,995
Eve=1	1,008 (25.2%)	997 (24.9%)	2,005
Total	2,025	1,975	4,000

Eve-Bob agreement rate: 50.3% (near-perfect random – consistent with

independent quantum measurements with no shared information)

4. VQC 4-qubit – Uniformity Test

4.1 Distribution (2,000 shots, 16 states)

State	Count	%	Expected (uniform)	Δ from uniform
0000⟩	201	10.1%	125	+76
0001⟩	129	6.5%	125	+4
0010⟩	93	4.7%	125	-32
0011⟩	37	1.8%	125	-88
0100⟩	113	5.7%	125	-12
0101⟩	175	8.8%	125	+50
0110⟩	95	4.8%	125	-30
0111⟩	167	8.4%	125	+42
1000⟩	168	8.4%	125	+43
1001⟩	73	3.7%	125	-52
1010⟩	188	9.4%	125	+63
1011⟩	177	8.9%	125	+52

State	Count	%	Expected (uniform)	Δ from uniform
$ 1100\rangle$	94	4.7%	125	-31
$ 1101\rangle$	80	4.0%	125	-45
$ 1110\rangle$	68	3.4%	125	-57
$ 1111\rangle$	142	7.1%	125	+17

4.2 Chi-Squared Uniformity Test

H_0 : Distribution is uniform across 16 states

H_1 : Distribution is non-uniform

$\chi^2 = 300.94$

df = 15

p $\approx$ 0.000000 (p $\ll$ 0.001)

Result: REJECT H_0

→ Distribution is significantly non-uniform.

→ This is a VQC/ansatz sampling circuit, NOT a QRNG.

→ Ground-state bias confirmed: $|0000\rangle$ dominant at 10.1% vs 6.25% expected.

4.3 Information Metrics

Shannon entropy H = 3.885 bits

Maximum possible = 4.000 bits ($\log_2 16$)

Entropy ratio = 97.1%

Parity balance = 50.2% even / 49.8% odd
(perfect symmetry)

5. QRNG 8-bit – Randomness Assessment

Shots : 10
Unique values : 10 / 10 (100% unique – no repeats)
Decimal values: [16, 90, 120, 128, 131, 187, 215, 242, 244, 254]
Range : 16 – 254 (out of 0-255)
Mean : 162.7 (theoretical uniform: 127.5)
Std dev : 75.8

Note: 10 shots is insufficient for formal uniformity testing
(chi-squared requires expected > 5 per cell → needs $5 \times 256 = 1,280$ shots for 8-bit uniform test). This experiment serves as a circuit validation run and qualitative demonstration, not a statistical QRNG benchmark.
For rigorous QRNG validation: repeat with $\geq 4,000$ shots.

6. Summary Statistics Table

Experiment	Shots	Metric	Value	95% CI
Bell ibm_torino	4,000	Fidelity	85.90%	[84.79%, 86.94%]
Bell ibm_fez R1	4,000	Fidelity	93.97%	[93.19%, 94.67%]
Bell ibm_fez R2	4,000	Fidelity	94.40%	[93.64%, 95.07%]
Bell ibm_fez 11⟩	1,000	Fidelity	96.30%	[94.94%, 97.30%]
Reproducibility	2×4,000	χ^2/p	2.70 / 0.44	H ₀ not rejected
BB84 Bob QBER	4,000	Error rate	49.00%	[47.43%, 50.52%]
BB84 Alice baseline	4,000	Error rate	0.60%	[0.39%, 0.90%]
BB84 degradation	—	Factor	82×	—
VQC 4q uniformity	2,000	χ^2/p	300.94 / ≈ 0	H ₀ rejected
VQC 4q entropy	2,000	Shannon H	3.885 bits	97.1% max
Concurrence C	1,000	C = 2F−1	0.926	—

Experiment	Shots	Metric	Value	95% CI
Tangle $T = C^2$	1,000	T	0.858	—
$E(Z,Z)$	1,000	Correlation	0.934	—

7. What Is Still Missing

The following values require the IBM calibration JSON export
 (`backend.properties()` output from IBM Quantum dashboard):

Parameter	How to obtain
T1 [μ s] per qubit	IBM dashboard → ibm_fez → Calibration → Export JSON
T2 [μ s] per qubit	Same as above
CZ gate error [%]	Same as above
Readout error [%] per qubit	Same as above
Gate time [ns]	Same as above

Export command in Qiskit:

```
from qiskit_ibm_runtime import
QiskitRuntimeService
service =
```

```
QiskitRuntimeService(channel='ibm_quantum_platform')
backend = service.backend('ibm_fez')
props = backend.properties()
props.to_dict() # save this as
ibm_fez_calib_YYYY-MM-DD.json
```