

Lecture 06: Evaluation of evaluations

PPHA 34600

Prof. Fiona Burlig

Harris School of Public Policy
University of Chicago

From last time: RCTs are the bee's knees

Randomized controlled trials are super powerful:

- Random assignment allows us to solve our selection problem
- We can implement them with tweaks to handle challenges:
 - *Noncompliance*: Dividing τ^{ITT} by share of compliers $\rightarrow \tau^{LATE}$
 - *Spillovers*: Proper design to avoid or measure
- More opportunities than you might imagine for implementation

Moving out of RCT land

We will spend the rest of the course on other research designs:

- Randomized controlled trials (RCTs)
- Trying to control for observable things
- Panel data
- Instrumental variables
- Regression discontinuity
- Big Data and machine learning

Why leave RCT land?

RCTs are the gold standard for a reason, but:

- They can be expensive
- Some programs require evaluation at scale
- RCTs can't always be implemented
- **There's a lot to learn from non-RCTs**

The ideal experiment

Even as we move away from RCTs, it's useful to consider the
ideal experiment

The ideal experiment

Even as we move away from RCTs, it's useful to consider the
ideal experiment

- “What experiment would I run to answer this question?”
- Useful to nail down your question of interest
- Valuable to think through problems with your non-RCT

The ideal experiment

Even as we move away from RCTs, it's useful to consider the
ideal experiment

- “What experiment would I run to answer this question?”
- Can be totally feasible (RED for energy efficiency upgrades)...
- ...or totally infeasible (randomly warm one Earth while keeping the other cold)

Testing quasi-experimental methods

We want to know how well our non-RCT toolkit works...

How can we test this?

Testing quasi-experimental methods

We want to know how well our non-RCT toolkit works...

How can we test this?

→ Compare an actual RCT to other methods *on the same data*

- Requires a setting with an RCT
- (Usually) toss the control group
- Use quasi-experimental methods to estimate $\hat{\tau}$
- Compare the RCT to the quasi-experimental methods

Testing quasi-experimental methods

We want to know how well our non-RCT toolkit works...

How can we test this?

→ Compare an actual RCT to other methods *on the same data*

- Requires a setting with an RCT
- (Usually) toss the control group
- Use quasi-experimental methods to estimate $\hat{\tau}$
- Compare the RCT to the quasi-experimental methods

→ What do we learn?

Testing quasi-experimental methods

We want to know how well our non-RCT toolkit works...

How can we test this?

→ Compare an actual RCT to other methods *on the same data*

- Requires a setting with an RCT
- (Usually) toss the control group
- Use quasi-experimental methods to estimate $\hat{\tau}$
- Compare the RCT to the quasi-experimental methods

→ What do we learn?

This is often referred to as a “LaLonde exercise” after LaLonde (1986)

Testing quasi-experimental methods

Comparing RCTs with quasi-experiments teaches us:

- If the estimates are the same:

Testing quasi-experimental methods

Comparing RCTs with quasi-experiments teaches us:

- If the estimates are the same:
 - Good sign!
 - This suggests that the quasi-experimental method is working properly


Testing quasi-experimental methods

Comparing RCTs with quasi-experiments teaches us:

- If the estimates are the same:
 - Good sign!
 - This suggests that the quasi-experimental method is working properly
- If the estimates are different:


Testing quasi-experimental methods

Comparing RCTs with quasi-experiments teaches us:

- If the estimates are the same:
 - Good sign!
 - This suggests that the quasi-experimental method is working properly
- If the estimates are different:
 - 
 - Something is likely going wrong with the non-RCT

Testing quasi-experimental methods

Comparing RCTs with quasi-experiments teaches us:

- If the estimates are the same:
 - Good sign!
 - This suggests that the quasi-experimental method is working properly
- If the estimates are different:
 - 
 - Something is likely going wrong with the non-RCT

Context is really important for this!

Leveraging an RCT we know and love

Blast from the past: We'll use the SMUD pricing RCT


Policy issue:

- The cost of providing electricity is time-varying
- Prices typically aren't
- This causes large welfare losses

Program:

- SMUD (randomly) implemented time-varying pricing
- Experimental run: 2011-2013
- Two flavors: “time-of-use” (TOU) and “critical peak pricing” (CPP)
- Both opt-in and opt-out versions

Fowlie & Wolfram et al results recap



Why did we want an RCT in this context?

Research question:

What is the effect of time-varying electricity pricing on consumption?

Why did we want an RCT in this context?

Research question:

What is the effect of time-varying electricity pricing on consumption?

Potential for selection into treatment on characteristics

- People who participate in utility programs look different
- Suppose *no response* to treatment: If units who choose treatment have different consumption patterns, we'll mistakenly measure $\hat{\tau} > 0$

Why did we want an RCT in this context?

Research question:

What is the effect of time-varying electricity pricing on consumption?

Potential for selection into treatment on characteristics

- People who participate in utility programs look different
- Suppose *no response* to treatment: If units who choose treatment have different consumption patterns, we'll mistakenly measure $\hat{\tau} > 0$

Potential for selection into treatment on τ_i

- People who choose to get treated may have different price sensitivity
- We know some of this is happening! (two LATEs)

What would the naive estimator do?

Recall our naive estimator:

$$\tau^N = \bar{Y}(1) - \bar{Y}(0)$$

What would the naive estimator do?

Recall our naive estimator:

$$\tau^N = \bar{Y}(1) - \bar{Y}(0)$$

Why is this problematic for electricity pricing?

Going beyond the naive estimator

We have a good idea that the naive estimator won't work...

... so we turn to alternative **research designs**

Going beyond the naive estimator

We have a good idea that the naive estimator won't work...

... so we turn to alternative **research designs**

A research design:

- Tries to solve the selection problem without randomization
- Invokes stronger assumptions than the RCT
- Allows us to make progress without randomization
- Best-case scenario: mimics an RCT

Spurlock et al test three common methods

We will cover each of these methods in detail later:

Spurlock et al test three common methods

We will cover each of these methods in detail later:

① Difference-in-differences

- Essentially compares treated and untreated units over time
- Intuition: I am similar to myself, treated or not...
- ... but to rule out other stuff happening, I use never-treated units

Spurlock et al test three common methods

We will cover each of these methods in detail later:

① **Difference-in-differences**

- Essentially compares treated and untreated units over time
- Intuition: I am similar to myself, treated or not...
- ... but to rule out other stuff happening, I use never-treated units

② **(Propensity score) matching**

- Essentially a sophisticated way of “controlling for stuff”
- Tries to generate a (non-experimental) control group
- Goal is to make this similar to the treated group

Spurlock et al test three common methods

We will cover each of these methods in detail later:

1 Difference-in-differences

- Essentially compares treated and untreated units over time
- Intuition: I am similar to myself, treated or not...
- ... but to rule out other stuff happening, I use never-treated units

2 (Propensity score) matching

- Essentially a sophisticated way of “controlling for stuff”
- Tries to generate a (non-experimental) control group
- Goal is to make this similar to the treated group

3 Regression discontinuity

- Essentially compares just-treated units to just-untreated units
- Leverages cutoffs in policy

Difference in difference approach

The DD approach compares units to themselves over time (3 steps):

Difference in difference approach

The DD approach compares units to themselves over time (3 steps):

- 1 Compare treated unit i in time t to (pre-treatment) i in time $t - 1$
 - Intuitive part: compare me to myself pre/post treatment

Difference in difference approach

The DD approach compares units to themselves over time (3 steps):

- ① Compare treated unit i in time t to (pre-treatment) i in time $t - 1$
 - Intuitive part: compare me to myself pre/post treatment
- ② Compare untreated unit j in time t to untreated j in time $t - 1$
 - Why do this?

Difference in difference approach

The DD approach compares units to themselves over time (3 steps):

- 1 Compare treated unit i in time t to (pre-treatment) i in time $t - 1$
 - Intuitive part: compare me to myself pre/post treatment
- 2 Compare untreated unit j in time t to untreated j in time $t - 1$
 - Why do this?
→ Control for common shocks to everyone
- 3 Subtract difference (1) from difference (2):

$$y_{it} = \alpha + \tau D_{it} + \underbrace{\gamma_i}_{i \text{ to itself}} + \underbrace{\delta_t}_{i \text{ to } j \text{ over time}} + \varepsilon_{it}$$

Difference in difference approach

The DD approach compares units to themselves over time (3 steps):

- ① Compare treated unit i in time t to (pre-treatment) i in time $t - 1$
 - Intuitive part: compare me to myself pre/post treatment
- ② Compare untreated unit j in time t to untreated j in time $t - 1$
 - Why do this?
→ Control for common shocks to everyone
- ③ Subtract difference (1) from difference (2):

$$y_{it} = \alpha + \tau D_{it} + \underbrace{\gamma_i}_{i \text{ to itself}} + \underbrace{\delta_t}_{i \text{ to } j \text{ over time}} + \varepsilon_{it}$$

For this to work, we require:

- Consumption for treated units is trending similarly to untreated units

Difference in difference approach

The DD approach compares units to themselves over time (3 steps):

- ① Compare treated unit i in time t to (pre-treatment) i in time $t - 1$
 - Intuitive part: compare me to myself pre/post treatment
- ② Compare untreated unit j in time t to untreated j in time $t - 1$
 - Why do this?
 - Control for common shocks to everyone
- ③ Subtract difference (1) from difference (2):

$$y_{it} = \alpha + \tau D_{it} + \underbrace{\gamma_i}_{i \text{ to itself}} + \underbrace{\delta_t}_{i \text{ to } j \text{ over time}} + \varepsilon_{it}$$


For this to work, we require:

- Consumption for treated units is trending similarly to untreated units


Note: Spurlock et al drop encouraged-but-untreated units

→ Was this necessary?

Difference in difference intuition



Comparing experimental and diff-in-diff results



Matching approach

(Propensity score) matching is a fancy way to control for stuff (2 steps):

Matching approach

(Propensity score) matching is a fancy way to control for stuff (2 steps):

- ① Use pre-treatment consumption to find untreated units that look like treated units
 - Eliminate dis-similar untreated units from the sample

Matching approach

(Propensity score) matching is a fancy way to control for stuff (2 steps):

- ① Use pre-treatment consumption to find untreated units that look like treated units
 - Eliminate dis-similar untreated units from the sample
- ② Estimate treatment effects for all treated units and selected untreated units only:

$$y_{it} = \alpha + \tau D_{it} + \gamma_i + \delta_t + \varepsilon_{it}$$

- (You can do this in several different ways)

Matching approach

(Propensity score) matching is a fancy way to control for stuff (2 steps):

- ① Use pre-treatment consumption to find untreated units that look like treated units
 - Eliminate dis-similar untreated units from the sample
- ② Estimate treatment effects for all treated units and selected untreated units only:

$$y_{it} = \alpha + \tau D_{it} + \gamma_i + \delta_t + \varepsilon_{it}$$


- (You can do this in several different ways)

For this to work, we require:

- Our selection control soaks up everything that matters!

Note: Same approach as DD, but now controlling for more

Comparing experimental and propensity score results



Regression discontinuity approach

The RD compares unit i to a nearly-identical unit j below a cutoff:

Regression discontinuity approach

The RD compares unit i to a nearly-identical unit j below a cutoff:

- Use a policy cutoff to “randomize” treatment
 - Compare a unit with pre-period consumption just below 100 to a unit with pre-period consumption just above 100

Regression discontinuity approach

The RD compares unit i to a nearly-identical unit j below a cutoff:

- Use a policy cutoff to “randomize” treatment
 - Compare a unit with pre-period consumption just below 100 to a unit with pre-period consumption just above 100

For this to work, we require:

- Units on either side of the cutoff are otherwise similar

Regression discontinuity approach

The RD compares unit i to a nearly-identical unit j below a cutoff:

- Use a policy cutoff to “randomize” treatment
 - Compare a unit with pre-period consumption just below 100 to a unit with pre-period consumption just above 100


For this to work, we require:

- Units on either side of the cutoff are otherwise similar


Note: Spurlock et al construct fake cutoffs

- They stitch together control group units with treatment group units
- The stitching point is their artificial cutoff


Regression discontinuity intuition



Comparing experimental and regression discontinuity



Comparing experimental and regression discontinuity



Spurlock et al find:

- Difference in difference estimates *understate* treatment effects
 - Due to unabsorbed selection

Spurlock et al find:

- Difference in difference estimates *understate* treatment effects
 - Due to unabsorbed selection
- Propensity score estimates *understate* treatment effects
 - Due to unabsorbed selection (controls make it worse!)

Spurlock et al find:

- Difference in difference estimates *understate* treatment effects
 - Due to unabsorbed selection
- Propensity score estimates *understate* treatment effects
 - Due to unabsorbed selection (controls make it worse!)
- Regression discontinuity estimates *overstate* treatment effects
 - Due to unabsorbed selection
 - OR estimating a different LATE

Spurlock et al find:

- Difference in difference estimates *understate* treatment effects
 - Due to unabsorbed selection
- Propensity score estimates *understate* treatment effects
 - Due to unabsorbed selection (controls make it worse!)
- Regression discontinuity estimates *overstate* treatment effects
 - Due to unabsorbed selection
 - OR estimating a different LATE
- Opt-out treatments are less biased than opt-in treatments
 - Intuition: We do better with a less-selected treatment

Exercise caution with non-experimental results

“Even though I was unable to evaluate all non-experimental methods, this evidence suggests that policymakers should be aware that the available non-experimental evaluations...may contain large and unknown biases resulting from specification errors.” – LaLonde (1986)

TL;DR:

- ① RCTs are (still) great!
- ② Quasi-experimental methods can get things wrong
- ③ We don't usually have a good experimental benchmark (💀)