Formulário

M/M/1

$$\pi_0 = 1 - \rho$$

$$\pi_n = \rho^n \pi_0 = \rho^n (1 - \rho), n \ge 1$$

$$L_q = \frac{\rho^2}{1 - \rho}$$

$$L_s = \rho$$

$$L = \frac{\rho}{1 - \rho}$$

$$W_q = \frac{\rho}{\mu(1 - \rho)}$$

$$W_s = 1/\mu$$

$$W = \frac{1}{\mu(1 - \rho)}$$

$$W_q(t) = \begin{cases} \rho, \text{ para } t = 0 \\ \rho e^{-\mu(1 - \rho)t}, \text{ para } t \ge 0 \end{cases}$$

M/M/s

$$\pi_0 = \left[\sum_{n=0}^{s-1} \frac{(s\rho)^n}{n!} + \frac{(s\rho)^s}{s!(1-\rho)} \right]^{-1}$$

$$\pi_n = \left\{ \frac{(s\rho)^n \pi_0}{n!}, para \ 1 \le n \le s \right.$$

$$\frac{s^s \rho^n \pi_0}{s!}, para \ n \ge s$$

$$P_B = \frac{\pi_s}{1 - \rho}$$

$$L_{q} = \frac{s^{s} \rho^{s+1} \pi_{0}}{s! (1-\rho)^{2}}$$

$$L_s = \lambda / \mu$$

$$W_q = L_q / \lambda$$

$$W_s = 1/\mu$$

$$W_{q}(t) = \begin{cases} 1 - \frac{(s\rho)^{s} \pi_{0}}{s!(1-\rho)}, para \ t = 0 \\ \frac{(s\rho)^{s} \pi_{0}}{s!(1-\rho)} e^{-s\mu(1-\rho)t}, para \ t > 0 \end{cases}$$