

HUMBOLDT-UNIVERSITÄT ZU BERLIN

Faculty of Life Sciences

**Development and Evaluation of an R package for
Nutrient Solution Formulation in Horticulture**

Dilân Olmo Şakar

Program: M.Sc. International Horticultural Science

Institute: Thaer-Institute of Agricultural and Horticultural Sciences

Department: Plant Nutrition and Fertilization

Supervisors: Prof. Dr. Eckhart George, Dipl.-Ing. Dr. Olivier Duboc

Institutions: Humboldt University of Berlin (HU) & University of
Natural Resources and Life Sciences, Vienna (BOKU)

Date: 12.05.2026

Table of contents

Abstract	i
List of Figures	ii
List of Tables	iii
List of Abbreviations	iv
1 Introduction	1
1.1 Gap: lack of transparent, reproducible, flexible formulation tools	1
1.2 Background and Motivation	3
1.3 Objectives and research questions	4
2 State of the Art / Scientific Background	6
2.1 Principles of nutrient solution formulation and the structure of experimental design in plant nutrition	6
2.2 Stoichiometry and salt–ion relationships	11
2.3 Existing tools	12
3 Material and Methods	13
3.1 Development and evaluation workflow	13
3.2 Software Environment and Data Sources	14
3.2.1 R version, packages	14
3.2.2 Built-in recipe datasets	14
3.3 Data Structures and Chemical Representation	16
3.3.1 Elementary Entities	16
3.3.2 Nutrient set	17
3.3.3 Compound-by-nutrient matrix	18
3.4 Algorithmic Methods	19
3.4.1 Formula parsing and molar mass calculation	19
3.4.2 Stoichiometric conversions	20
3.4.3 NNLS optimization with weighting	22
3.4.4 Target scaling and importance weighting	23
3.4.4.1 Importance weighting of nutrient targets	23
3.4.4.2 Target-based scaling of residuals	24
3.4.5 Interpretation of the solution	25
3.4.6 Error quantities reported by NutriCalc	25
3.4.7 Two-stage NNLS solver: separation of neutral salts and acid–base reagents	26

3.5	Chemical Modeling	28
3.5.1	Water quality and input handling	28
3.5.2	Estimation of pH via charge balance	29
3.5.3	EC estimation using ionic diffusion coefficients	30
3.6	Shiny Application Workflow	33
3.6.1	Installation or launching the web application	33
3.6.2	Input parameters	34
3.6.2.1	Nutrient Targets	34
3.6.2.2	Water	34
3.6.2.3	Fertilizers	35
3.6.2.4	Recipes	35
3.6.2.5	Tissue	35
3.6.3	Output parameters	36
3.6.3.1	Result	36
3.6.3.2	pH & EC	36
3.6.3.3	Solution	37
3.6.3.4	Anions and Cations	37
3.7	Case study selection	37
3.7.1	Case Study I: Alternative Hoagland II nutrient solution formulation . . .	38
3.7.2	Case Study II: Tomato nutrient solution using Berlin-Dahlem water with acid adjustment	38
3.7.3	Case Study III: Nutrient solution with increased sodium concentration for halophytes using mixed water sources	38
4	Results	39
4.1	General evaluation across crop specific recipes	39
4.1.1	Achieved Nutrient solution formulations	39
4.2	pH and EC Outputs	41
4.2.0.1	Performance of pH Calculation	44
4.2.0.2	Performance of EC Calculation	44
4.2.1	Comparison with published formulation	45
4.3	Case Study Results	47
4.3.1	Case study 1 — Experimental scientist: classical solution reconstruction under restricted salt availability (deionized water) under different EC. . .	47
4.3.1.1	Achieved composition	47
4.3.1.2	Deviations from target	48
4.3.1.3	pH and EC	48
4.3.2	Case study 2 — Horticultural technician: tomato nutrient formulation us- ing Berlin-Dahlem water with acid adjustment	49
4.3.2.1	Achieved composition	49
4.3.2.2	Deviations from target	49
4.3.2.3	pH and EC	49

4.3.3	Case study 3 — Research scientist: nutrient formulation for a novel halo- phyte (<i>Salicornia</i>) using mixed water sources	51
4.3.3.1	Achieved composition	51
4.3.3.2	Deviations from target	52
4.3.3.3	pH and EC	52
5	Discussion	53
5.1	Solver robustness and formulation performance	53
5.2	Exact solvability, electroneutrality and non uniqueness	54
5.3	Error metrics and interpretation of residual deviation	55
5.4	Weighting and two stage optimisation as practical control tools	56
5.5	Electroneutrality and pH as coupled constraints	56
5.6	Role of pH and EC modelling in the workflow	57
5.7	Relevance of the mass balance approach	58
5.8	Comparison with existing formulation approaches	59
5.9	Limitations and scope	59
6	Conclusions and Outlook	61
6.1	Scientific contribution	61
6.2	Practical relevance for horticulture	61
6.3	Limitations	62
6.4	Future extensions	62
7	Summary (possibly in German and English)	64
8	References	65
	Appendix	vi
	Declaration of Independence	vii

Abstract

Efficient nutrient-solution formulation is fundamental to controlled horticultural production, yet in practice it is often based on fixed recipes or iterative manual adjustment. This thesis addresses that limitation through the development of a modular open-source computational framework in R for reproducible nutrient-solution design called NutriCalc.

The central objective was to formalize nutrient formulation as a mathematically defined problem that can be solved transparently under agronomic and chemical constraints. To achieve this, the formulation task was expressed as a constrained linear optimization problem and solved using non-negative least squares, ensuring physically meaningful fertilizer quantities. The framework combines stoichiometric conversion, unit transformation, charge-balance-based evaluation and estimation of pH and electrical conductivity, thereby linking nutrient targets to key physicochemical properties of the final solution.

NutriCalc was implemented both as an R package for programmable workflows and as a Shiny-based graphical user interface for interactive use. Overall, the thesis demonstrates that nutrient-solution design can be shifted from a recipe-driven practice to a reproducible, optimization-based and chemically constrained workflow. NutriCalc provides a transparent computational foundation for horticultural research, teaching and applied nutrient management.

List of Figures

2.1	Phosphate speciation (percent) as a function of pH.	8
4.1	Comparison of 50 mL of 0.1 mol L ⁻¹ KH ₂ PO ₄ + x mL of 0.1 mol L ⁻¹ NaOH (25 °C).	44
4.2	Screenshot of the fertilizer salt composition calculated in the NutriCalc Shiny app for a tomato nutrient solution in rockwool using tap water from Berlin postal code 14195. The formulation is shown separately for A-BAK, B-BAK, and micronutrients stock solutions. Fertilizer requirements are reported for the preparation of 500 L nutrient solution and the amounts for 18 L of a 50-fold concentrated stock solution.	50
4.3	Screenshot of the fertilizer salt composition calculated in the NutriCalc Shiny app for a tomato nutrient solution in rockwool using tap water from Berlin postal code 14195. The formulation is shown separately for A-BAK, B-BAK, and micronutrients stock solutions. Fertilizer requirements are reported for the preparation of 500 L nutrient solution and the amounts for 18 L of a 50-fold concentrated stock solution.	52

List of Tables

3.1	Target compositions of crop-specific nutrient solutions (Olericulture, Floriculture, Fruticulture) from Sonneveld & Straver (1994)	15
3.2	Molar masses of elementary entities used for nutrient formulation.	16
3.3	Fertilizer compounds and their molar masses.	17
3.4	Compound–nutrient stoichiometric matrix (mmol nutrient per mmol compound). Stage 1 = salts, Stage 2 = acids/bases.	18
4.1	Achieved Nutrient Solutions with SSE and SSRE. Asterisks denote two-stage optimization; parentheses indicate standard-solver errors.	40
4.2	Solver-derived pH, ionic strength, total charge equivalents (TCE), electrical conductivity (EC) and osmotic pressure for crop-specific formulations (salts-only; water = 0; KS 4.3 = 0; CO ₂ (aq) = 0). EC sonneveld and reportet EC for comparison.	42
4.3	pH and EC comparison across activity-coefficient models (salts-only; water = 0; KS 4.3 = 0; CO ₂ (aq) = 0). EC Sonneveld is the linear estimate from TCE (meq/L).	43
4.4	Electrical conductivity of KCl solutions at 25 °C.	45
4.5	Comparison of Sonneveld formulation and NutriCalc solution.	45
4.6	Comparison of fertilizer additions in the Sonneveld and NutriCalc formulations.	45
4.7	Comparison of the original Hoagland solution No. 2 fertilizer combination and three alternative fertilizer combinations achieving the same target macronutrient composition. Values are given in mmol L ⁻¹	47
4.8	Amounts of fertilizer salts required to prepare modified Hoagland solution No. 2 at target EC values from 0.5 to 3.0 mS cm ⁻¹ . Salt amounts are given in mmol L ⁻¹	48

List of Abbreviations

A-BAK Stock solution A [A-stammlösung / stock tank A]

B-BAK Stock solution B [B-stammlösung / stock tank B]

BK Base Capacity [Basenkapazität]

BOKU University of Natural Resources and Life Sciences, Vienna [Universität für Bodenkultur Wien]

BWB Berliner Wasserbetriebe

CO₂(aq) Aqueous carbon dioxide

CAN Calcium ammonium nitrate

CRC CRC Handbook of Chemistry and Physics

CSV Comma-Separated Values

DWC Deep Water Culture

EC Electrical Conductivity

EDTA Ethylenediaminetetraacetic acid

Fe-EDTA Iron ethylenediaminetetraacetate

GUI Graphical User Interface

HU Humboldt-Universität zu Berlin

meq L⁻¹ Milliequivalents per litre

mmol L⁻¹ Millimoles per litre

N_A Avogadro constant

NNLS Non-Negative Least Squares

pH Negative logarithm of the hydrogen ion activity

PHREEQC PH REdox EQUilibrium; Geochemical speciation and reaction-modeling software developed by the U.S. Geological Survey

R The R programming language and software environment for statistical computing

RDW Reuse drainage water

SI International System of Units [Système international d'unités]

SK Acid Capacity [Säurekapazität]

SSE Sum of Squared Errors

SSRE Sum of Squared Relative Errors

TCE Total Charge Equivalents

WUE Water Use Efficiency

1 Introduction

Precise nutrient-solution formulation is fundamental to controlled horticultural production, particularly in hydroponic and other soilless cultivation systems. In these systems plant nutrition depends on the chemical composition of the supplied solution, so inaccuracies in formulation can directly affect plant performance, experimental validity and resource-use efficiency.

In practice, however, nutrient-solution design remains a complex task. Fertilizer salts often supply multiple nutrients simultaneously, local irrigation water contributes additional ions and target concentrations must be translated across different chemical and commercial representations. As a result, the formulation process is error-prone, when performed manually or with opaque spreadsheet-based workflows, especially when transparency, reproducibility and extensibility are required.

These challenges are particularly relevant in horticultural research, where treatments are often designed to manipulate individual nutrients while keeping other factors as constant as possible. Because ions are chemically coupled through shared salts, electroneutrality and solution chemistry, such designs require computationally explicit methods rather than ad hoc adjustment. Despite this need, no dedicated modular open-source framework in R is currently available for transparent nutrient-solution formulation and analysis in horticultural applications.

This thesis addresses that gap through the development of NutriCalc, an open-source computational framework in R for the formulation, analysis and optimization of nutrient solutions. The work formalizes nutrient formulation as a constrained linear optimization problem solved using non-negative least squares and extends this workflow through stoichiometric conversion, charge-balance-based evaluation, water input and estimation of pH and electrical conductivity. In addition to the computational backend, the framework is implemented as a Shiny-based graphical user interface to support practical use by researchers and horticultural practitioners. The overarching objective of the thesis is to provide a reproducible chemically coherent and extensible framework that shifts nutrient-solution formulation from a recipe-based practice to a transparent computational workflow.

1.1 Gap: lack of transparent, reproducible, flexible formulation tools

A reconstruction of nutrient calculations reported in the literature further illustrates the relevance of transparent conversion pipelines. For example, J. J. Jones et al. (2024) describe a control nutrient solution for CUBES Circle derived from defined molar concentrations of fertilizer salts (5.15 mM $\text{Ca}(\text{NO}_3)_2$; 8.79 mM KNO_3 ; 2.36 mM MgSO_4 ; 2.94 mM KH_2PO_4), subsequently scaled to a

10 % strength solution. Recalculating the resulting elemental concentrations reported in the paper reveals that the final values depend critically on how stoichiometric relationships, dilution factors and ion attribution are applied. In particular, deviations may arise when dilution factors are inconsistently propagated across elements, when molar masses of salts are used instead of elemental molar masses, or when contributions of shared ions are only partially accounted for. While K, P, and S match the reported control values, the largest discrepancies occur for Ca, Mg, and NO_3^- . Recomputing the resulting concentrations using stoichiometric attribution and elemental molar masses yields:

```
compute_molar_mass("Ca")
```

```
[1] 40.078
```

$$c(\text{Ca}) = 5.15 \text{ mmol L}^{-1} \times 40.08 \text{ mg mmol}^{-1} \times 0.1 = 20.64 \text{ mg L}^{-1}.$$

```
compute_molar_mass("Mg")
```

```
[1] 24.305
```

$$c(\text{Mg}) = 2.36 \text{ mmol L}^{-1} \times 24.31 \text{ mg mmol}^{-1} \times 0.1 = 5.74 \text{ mg L}^{-1}.$$

```
compute_molar_mass("N03")
```

```
[1] 62.004
```

$$c(\text{NO}_3^-) = (2 \times 5.15 + 8.79) \text{ mmol L}^{-1} \times 62.01 \text{ mg mmol}^{-1} \times 0.1 = 118.38 \text{ mg L}^{-1}.$$

The reported Ca value (82.05 mg L^{-1}) is consistent with using the molar mass of $\text{Ca}(\text{NO}_3)_2$ (and effectively 5.0 mM instead of 5.15 mM) rather than elemental Ca:

```
compute_molar_mass("Ca(N03)2")
```

```
[1] 164.086
```

$$c(\text{Ca})_{\text{reported}} = 5.0 \text{ mmol L}^{-1} \times 164.09 \text{ mg mmol}^{-1} \times 0.1 = 82.05 \text{ mg L}^{-1}.$$

The reported Mg value (57.37 mg L^{-1}) is consistent with the 0.1 scaling factor not being applied:

$$c(\text{Mg})_{\text{reported}} = 2.36 \text{ mmol L}^{-1} \times 24.31 \text{ mg mmol}^{-1} = 57.37 \text{ mg L}^{-1}.$$

The reported nitrate value (63.87 mg L^{-1}) is consistent with counting nitrate only from $\text{Ca}(\text{NO}_3)_2$ (omitting the KNO_3 contribution):

$$c(\text{NO}_3^-)_{\text{reported}} = (2 \times 5.15) \text{ mmol L}^{-1} \times 62.01 \text{ mg mmol}^{-1} \times 0.1 = 63.87 \text{ mg L}^{-1}.$$

While such differences do not necessarily affect the overall conclusions of the study, they demonstrate how small ambiguities in conversion logic can lead to substantial divergence in reported nutrient concentrations. These observations underline that even potentially well documented experimental studies can benefit from calculation frameworks in which unit conversions, stoichiometric coefficients and scaling factors are explicitly encoded and reproducible. This is particularly relevant for integrated production systems, where nutrient streams are reused across modules and quantitative consistency is essential.

1.2 Background and Motivation

Precise nutrient solution formulation is a cornerstone of modern horticultural production, particularly in hydroponic and soilless cultivation systems. This process involves selecting and combining soluble fertilizer salts to deliver specific concentrations of essential elements in desired chemical forms and ratios (J. B. Jones, 2014). While standard formulations were published since the mid-19th century (Hoagland, 1920; Hoagland & Arnon, 1938; Knop, 1860; Shive, 1915), practitioners frequently need to tailor them to local water quality, crop-specific requirements or available fertilizers, creating a complex and often error-prone challenge (Resh, 2013; Sonneveld, 2009).

In plant nutrition experiments, treatments are typically designed to vary individual nutrients while holding others constant (*ceteris paribus*) to investigate the effect of a single nutrient element. However, due to the interdependencies among ions and their shared presence in various fertilizer salts, such isolation is rarely achievable in practice. Researchers must therefore account for the introduction of accompanying ions, which may affect plant responses and experimental validity.

Creating accurate nutrient solutions under such constraints is time-consuming and sometimes mathematically infeasible, underscoring the need for computational tools that enable efficient, accurate, and reproducible formulation under realistic chemical and experimental conditions.

The nutrient formulation problem is inherently mathematical. Many fertilizer salts contribute multiple nutrients (e.g., $\text{Ca}(\text{NO}_3)_2$ supplies both calcium and nitrate), leading to an interdependent system. This can be modeled as a system of linear equations:

$$A \cdot x = b$$

Where:

- A is the nutrient matrix (each row representing a nutrient, each column a fertilizer salt),
- x is the vector of unknown fertilizer quantities,
- b is the vector of target nutrient concentrations.

Due to overlapping contributions and limited degrees of freedom, exact solutions may not exist. Thus, approximate solutions are needed, with the additional constraint that all fertilizer quantities must be non-negative. This motivates the use of non-negative least squares (NNLS) optimization, which minimizes the squared error between the achieved and target nutrient vectors under the constraint $x \geq 0$.

Moreover, real-world use requires support for various chemical notations such as conversions between elemental and oxide forms (e.g., $P \leftrightarrow P_2O_5$, $S \leftrightarrow SO_4^{2-}$) and integration with commercially available fertilizers labeled in % w/w or expressed in units such as mmol/L or mg/L. Despite the existence of some basic Windows applications and Excel tools, there is currently no open-source, modular, and transparent software solution in R that offers this level of flexibility and chemical accuracy for horticultural applications.

1.3 Objectives and research questions

The main objective of this thesis is to design and implement a computational framework in R that enables the formulation, analysis and optimization of nutrient solutions specifically for controlled-environment horticulture. This framework is intended to support accurate and efficient nutrient planning and to facilitate reproducibility and transparency in both research and practical applications.

More specifically, the thesis aims to develop tools that can construct nutrient matrices from molecular fertilizer formulas. These tools will allow for precise conversion between fertilizer compounds and their elemental forms using stoichiometric calculations. The formulation process will be modelled mathematically as a non-negative linear system and solved using Non-Negative Least Squares optimization methods (Mullen & van Stokkum, 2024).

In addition to the computational backend, the framework will include features to visualize the distribution of nutrients across different fertilizer sources. Furthermore, it will offer the ability to approximate the pH and EC of a nutrient solution in advance, based on the concentrations and molar conductivity of respective ions (McCleskey et al., 2012).

The tool will also be designed with extensibility in mind, laying the groundwork for the import of commercial fertilizer compositions. In the long term, the framework will be equipped to benchmark nutrient formulations based on either experimental or simulated crop yield data.

Finally, to ensure accessibility and ease of use, the thesis will culminate in the development of a user-friendly graphical interface using the Shiny framework, enabling non-technical users to benefit from the computational functionalities (Chang et al., 2025).

2 State of the Art / Scientific Background

2.1 Principles of nutrient solution formulation and the structure of experimental design in plant nutrition

This chapter addresses horticultural plant nutrition experiments aimed at evaluating nutrient composition and its effects under chemically and statistically controlled conditions. It does not address other types of biological experiments, such as studies on uptake mechanisms, isotope tracing, signaling pathways, or related physiological processes.

Experiments in plant nutrition seek to identify causal relationships between specific nutrients and plant responses such as growth, yield, vitality, and physiological performance. Despite the apparent simplicity of manipulating nutrient concentrations, such experiments are fundamentally constrained by chemical laws that limit causal identifiability. In particular, electroneutrality and the statistical requirement of *ceteris paribus* jointly restrict how nutrient solutions can be manipulated and how resulting effects may be interpreted. This chapter develops a formal framework, expressed in milliequivalent (meq) notation, that derives classical experimental designs in plant nutrition from first principles and clarifies the interpretational limits of nutrient experimentation.

A nutrient solution may be represented as a vector of its ionic charge equivalents:

$$\mathbf{S}_{\text{meq}} = (\text{K}^+, \text{Ca}^{2+}, \text{Mg}^{2+}, \dots \mid \text{NO}_3^-, \text{Cl}^-, \text{SO}_4^{2-}, \dots)$$

where each component is expressed in milliequivalents per liter (meq L^{-1}). For any ion i ,

$$\text{meq}_i = |z_i| c_i,$$

with z_i denoting ionic charge and c_i the molar concentration. Expressing solution composition in meq makes ionic charge balance explicit and avoids ambiguity associated with heterovalent ions.

Not all vectors in this space are chemically admissible. All aqueous nutrient solutions must satisfy electroneutrality, which in meq notation takes the form

$$\sum \text{meq}_{\text{cat}} - \sum \text{meq}_{\text{an}} = 0.$$

The sums include all charged species present, including H^+ (formally H_3O^+) and OH^- . Electroneutrality implies that no ionic component can be altered independently of ions carrying opposite charge. Any experimental manipulation therefore induces compensatory changes elsewhere in the ionic system. In this chapter, meq values are treated as analytical charge equivalents used for experimental design; activity coefficients, complexation, and ion pairing may alter speciation but do not remove the electroneutrality constraint.

In addition to charge balance, many nutritionally relevant components participate in acid–base equilibria and therefore exist as families of chemical species. For example, phosphate occurs as H_2PO_4^- , HPO_4^{2-} , and PO_4^{3-} , while carbonate occurs as CO_2 , HCO_3^- , and CO_3^{2-} . Let c_i^{tot} denote the total analytical concentration of component i . The total charge contribution of this component to the solution can be written as

$$\text{meq}_i^{\text{tot}} = \sum_j |z_{ij}| c_{ij},$$

where c_{ij} is the concentration of species j belonging to component i . Changes in pH redistribute charge among species with different valencies as shown for phosphate in Figure 2.1, thereby altering the effective ionic composition of the solution even when total analytical concentrations are held constant. Although pH can be experimentally controlled using buffers or acid–base additions, such control necessarily introduces additional ionic components and therefore does not decouple pH from nutrient composition in a causal sense.

From a causal-inference perspective, identification of an effect requires isolating variation in a target factor while holding other causally relevant factors fixed or blocked (*ceteris paribus*). In plant nutrition experiments, strict independent variation of a single ionic species is generally not possible, because electroneutrality and acid–base equilibria couple ionic charge, pH, ionic strength, and osmotic potential. As a result, strict *ceteris paribus* conditions for individual ions cannot be achieved in bulk nutrient solutions; they can only be approximated within constrained experimental frameworks. Within this framework, classical experimental designs in plant nutrition can be derived as three distinct classes of experiments, defined by the invariants they preserve in meq-space.

Here, Δ denotes the change induced by the experimental manipulation, i.e. $\Delta x := x_{\text{treatment}} - x_{\text{control}}$ (or, more generally, the difference between two solution states). The first class consists of qualitative, or ion-substitution, experiments. In these experiments, one ionic species is replaced by another while total cation and total anion charge equivalents are kept constant:




Figure 2.1: Phosphate speciation (percent) as a function of pH.

$$\Delta \text{meq}_i + \Delta \text{meq}_j = 0.$$

When the substituted ions have the same charge, the substitution is isovalent (e.g. $\text{K}^+ \leftrightarrow \text{Na}^+$ or $\text{Cl}^- \leftrightarrow \text{NO}_3^-$). When the charges differ, the substitution is heterovalent, for example replacing 2 meq of K^+ (2 mmol) with 2 meq of Ca^{2+} (1 mmol). Qualitative substitution experiments provide the highest degree of causal specificity achievable within whole-solution manipulations at fixed charge totals, although their interpretation remains conditional on buffering and counter-ion choice.

The second class comprises quantitative, or salt addition (or removal), experiments. Here, the total amount of dissolved salt is changed, which necessarily alters both total cation and total anion charge equivalents by the same non-zero amount:

$$\Delta \sum \text{meq}_{\text{cat}} = \Delta \sum \text{meq}_{\text{an}} \neq 0.$$

Electroneutrality is preserved automatically, but ionic strength, electrical conductivity, and osmotic potential change and pH may shift depending on buffering. Because at least two ionic components vary simultaneously, plant responses in such experiments reflect composite effects of nutrient ions, counter-ions and general salinity.

The third class treats quantity as an emergent quality through solution-strength series. In this case, all charge equivalents in the solution are scaled by a constant factor k :

$$\mathbf{S}_{\text{meq},k} = k \mathbf{S}_{\text{meq},1}.$$

This transformation preserves ion ratios expressed in meq while increasing or decreasing total charge density, ionic strength and osmotic potential. Although composition remains constant at the level of charge ratios, pH and ionic speciation may shift because acid–base equilibria are non-linear. As a result, solution strength functions as an environmental condition rather than a simple quantitative variable.

Across these three experiment types, causal specificity for individual ion identity typically decreases as the number of inseparable or uncontrolled variables increases. Qualitative substitution experiments offer the highest degree of causal specificity, whereas quantitative salt manipulations and solution-strength series progressively trade causal resolution for ecological or agronomic relevance. These experiment classes are therefore best understood as distinct projections of a chemically constrained ionic system, most transparently described in milliequivalent space.

Experiment class	Core manipulation	Charge balance condition	Held constant	Necessarily changes	Example	Interpretation
I.a Qualitative (isovalent)	Equal-meq substitution	$\Delta \sum \text{meq}_{cat} = 0$ $\Delta \sum \text{meq}_{an} = 0$	Total cation & anion meq	Ion identity	$\text{K}^+ \leftrightarrow \text{Na}^+$	Conditional ion identity effect
I.b Qualitative (heterovalent)	Equal-meq substitution	$\Delta \text{meq}_i + \Delta \text{meq}_j = 0$	Charge equivalents	Molar conc., ionic strength	2 mmol $\text{K}^+ \leftrightarrow$ 1 mmol Ca^{2+}	Identity + secondary effects
II Quantitative	Salt addition or removal	$\Delta \sum \text{meq}_{cat} =$ $\Delta \sum \text{meq}_{an} \neq 0$	Electroneutrality	Ions, EC, osmotic potential	$\uparrow$ KCl	Composite salt effect
III Quantitative-as-qualitative	Solution scaling	$\mathbf{S}_{\text{meq},k} = k \mathbf{S}_{\text{meq},1}$	meq ratios	Total charge, speciation	$\frac{1}{4}\times, \frac{1}{2}\times, 1\times$	Environmental strength effect

From a plant nutrition perspective, comparisons between nutrient solutions of different provenance are only informative to the extent that differences in ionic composition are maintained or eliminated in a controlled manner. If two solutions of different origin, such as wastewater-derived (e.g. animal or human urine) and mineral solutions, are formulated to the same ionic composition, then the specifically nutritional dimension of the comparison is largely removed, reducing the question to a Ship of Theseus problem of identity. Under such conditions, any remaining differences are more appropriately interpreted in terms of economic factors associated with source, production, transport, cost and sustainability rather than nutrient composition itself.

2.2 Stoichiometry and salt–ion relationships

In hydroponics, nutrients are supplied by dissolving fertilizer salts in water to prepare a nutrient solution (Resh, 2013). The formulation of such solutions is complicated by the declaration conventions used for commercial fertilizers. Nutrient contents are commonly expressed in oxide form rather than as elemental concentrations or ionic species. Thus, potassium is typically declared as K_2O rather than as K or ionic K^+ , while phosphorus is conventionally declared as P_2O_5 rather than as P or as the pH-dependent phosphate species $H_2PO_4^-$, HPO_4^{2-} , or PO_4^{3-} . Although these conventions are historically established, they introduce an unnecessary level of abstraction and increase the susceptibility of nutrient calculations to error.

A more serious limitation is that fertilizer labels do not necessarily disclose all contained elements. As a result, attempts to recalculate complete fertilizer compositions in ionic form often fail to account for 100 % of the product mass. One reason is that elements such as chloride may be present without mandatory declaration. Under Regulation (EC) No 2003/2003, only specific nutrient categories are subject to declaration requirements, namely the primary nutrients nitrogen, phosphorus, and potassium; the secondary nutrients calcium, magnesium, sodium, and sulfur; and the micronutrients boron, cobalt, copper, iron, manganese, molybdenum, and zinc. In addition, secondary nutrients must only be declared above specified threshold concentrations. For sodium, for example, declaration is required only above 3 % Na_2O or 2.2 % Na .

For this reason, the present work was restricted to single salts or mixtures of single salts with chemically defined composition. Such compounds can be represented directly by their molecular formulas, so that molar mass and stoichiometric composition are fully determined by the sum of the constituent elements and their stoichiometric coefficients. This ensures complete compositional transparency and avoids the uncertainty associated with incomplete commercial fertilizer declarations.

Many fertilizers used in greenhouse production are in fact single salts. A representative example is Epsom salt, $MgSO_4 \cdot 7H_2O$, marketed by Yara as YaraTera KRISTA MgS and declared as containing 16 % MgO and 32.5 % SO_3 (YARA, 2026). Chemically, pure $MgSO_4 \cdot 7H_2O$ contains 9.86 % Mg and 13.01 % S , corresponding to 16.35 % MgO and 32.48 % SO_3 , or 38.98 % SO_4 . Thus, 51.16 % of the compound mass consists of water of crystallization. Another example is calcium nitrate, which may occur as anhydrous $Ca(NO_3)_2$, as the hydrate $Ca(NO_3)_2 \cdot 4H_2O$, or as the double salt $5Ca(NO_3)_2 \cdot NH_4NO_3 \cdot 10H_2O$. These distinctions are stoichiometrically relevant because each

form differs in molar mass and nutrient contribution, even where the commercial designation is similar.

2.3 Existing tools

A classical approach to nutrient-solution calculation is the universal algorithm proposed by Sonneveld et al. (1999). The method follows an iterative balancing procedure in which fertilizer salts are added stepwise to meet nutrient targets, while the residual nutrient requirements are recalculated after each addition because each salt simultaneously contributes several ions. In this way, the final formulation is obtained through successive correction of remaining deficits under the constraints imposed by salt composition, target EC and raw-water composition.

Nutrient Solution Calculator is an Excel™-based tool developed by Luca Incrocci and supported by the EUPHOROS project (Pardossi et al., 2011). It was previously available for download via Wageningen University and was still accessible in 2024, but was no longer publicly available there in 2026 (Incrocci, 2007). A Spanish translation remains available online through the University of Almeria (Incrocci et al., 2020). The spreadsheet is designed to calculate the composition and cost of fertigation stock solutions (A+B) from three principal inputs: the ionic composition of the irrigation water, the crop nutrient recipe, including target pH, EC, and nutrient concentrations and the technical characteristics of the fertigation system, particularly stock tank volume and dilution ratio. Furthermore, it evaluates the potential risk of salt precipitation in concentrated stock solutions. The spreadsheet is password-protected.

Another tool is HydroBuddy, a free and open-source software application developed by Daniel Fernández Pinto for the formulation of hydroponic nutrient solutions (Fernández Pinto, 2022). The program calculates the required quantities of fertilizers to achieve specified target nutrient concentrations. It includes a database of commonly used fertilizers whose compositions can be edited or extended by the user and allows the formulation of both direct nutrient solutions and concentrated stock solutions. The software determines the combination of fertilizer salts that best matches the desired nutrient composition through the solution of a system of linear equations and also provides an empirical estimation of the EC.

At present, no dedicated open-source R library is available for transparent, modular nutrient-solution calculation in horticultural applications.

3 Material and Methods

3.1 Development and evaluation workflow

The development of the computational framework followed a structured development and validation pipeline. For the development of the R package devtools was used (Wickham et al., 2025).

The first step involved designing the underlying data structures. A nutrient matrix was established, in which the rows represent essential nutrients (such as N, P, K, Ca, etc.) and the columns represent fertilizer salts. For readability the input matrix was transposed and converted for the calculation. Each fertilizer was parsed based on its molecular formula into its elemental components, using a database of molar masses and established stoichiometric rules.

Building on this, an R package was developed. This package includes core functions for parsing chemical formulas and computing molar masses, as well as for converting between different chemical representations such as oxides and elemental forms (e.g., P_2O_5 to P or SO_4 to S). It also includes tools for transforming concentration units (e.g., from mmol/L to mg/L) using accurate molar mass calculations. Also bidirectional transformations between compound and elemental quantities, incorporating stoichiometric logic, was integrated.

The central component of the package was the implementation of the nutrient solution formulation algorithm, using the `nnls::nnls()` function to solve the system of equations under non-negativity constraints and giving the difference between desired and obtained solution as absolute and relative error. The framework allows users to assign weights to individual nutrients, enabling exploration of different formulation priorities and trade-offs.

The framework was validated through a series of use cases. These included reproducing historical and crop-specific nutrient solutions, comparing the computed results with values from the literature and collecting some crop and author specific nutrient solutions in a base library. Nutrient distribution of anions and cations across different solutions is visualized e.g. by ternary plot for cations (K^+ , Ca^{2+} , Mg^{2+}) and anions (NO_3^- , SO_4^{2-} , PO_4^{3-}) to systematically classify different nutrient combinations as done by Steiner (1961).

3.2 Software Environment and Data Sources

3.2.1 R version, packages

The package `nutricalc` was written in R 4.5.1 (R Core Team, 2025) using Posit RStudio 2025.09.2+418 “Cucumberleaf Sunflower” (Posit team, 2025). Following packages were used: `stringr` (Wickham et al., 2025), `nnls` (Mullen & van Stokkum, 2024), `shiny` (Chang et al., 2025), `bslib` (Sievert et al., 2025), `Ternary` (Smith & Sanselme, 2025), `xml2` (Wickham et al., 2026). For the development in R `devtools` (Wickham, 2025) was used.

Code of the package is provided on GitHub under <https://github.com/diolsa/nutricalc> and the shiny-App is deployed under: <https://nutricalc.shinyapps.io/nutricalc-app/>

Code development was supported by AI-assisted LLM for debugging and optimization to ensure code quality and computational reliability with OpenAI CODEX and ChatGPT 5.2.

3.2.2 Built-in recipe datasets

Hewitt (1966) compiled a large proportion of the nutrient-solution formulations used during the first half of the twentieth century. For the present work, selected standard recipes from this compilation were considered, particularly those that remain relevant in contemporary research and horticultural practice. In addition, 38 crop-specific horticultural recipes from Sonneveld & Straver (1994) were included. These recipes were assigned to the three groups Olericulture, Floriculture, and Fruticulture. In the original source, Sonneveld & Straver (1994) also classified the formulations into the categories A, B, and C, reflecting their practical and experimental suitability for the respective crop. The resulting set of formulations is shown in Table 3.1.

Table 3.1: Target compositions of crop-specific nutrient solutions (Olericulture, Floriculture, Fruticulture) from Sonneveld & Straver (1994)

Category	Recipe	mmol L ⁻¹							μmol L ⁻¹						
		NO ₃ -N	NH ₄ -N	P	K	Ca	Mg	S	Fe	Mn	Zn	B	Cu	Mo	Si
Olericulture	Endive in Recirculating Water	19	1.25	2	9	5	1.5	1.125	40	5	4	30	0.75	0.5	0
	Eggplant in Rockwool	15.5	1.5	1.25	6.75	3.25	2.5	1.5	15	10	5	30	0.75	0.5	0
	Eggplant in Rockwool (RDW)	11.75	1	1	6.5	2.25	1.5	1.125	15	10	5	20	0.75	0.5	0
	Bean in Rockwool	12	1	1.25	5.5	3.25	1.25	1.125	10	10	4	20	0.5	0.5	0
	Courgette in Rockwool	16	1.25	1.25	7.25	3.625	2	1.25	10	10	5	30	0.75	0.5	0
	Cucumber in Rockwool	16	1.25	1.25	8	4	1.375	1.375	15	10	5	25	0.75	0.5	750
	Cucumber in Rockwool (RDW)	11.75	1	1.25	6.5	2.75	1	1	15	10	5	25	0.75	0.5	750
	Propagation vegetable plants in rockwool	16.75	1.25	1.25	6.75	4.5	3	2.5	25	10	5	35	1	0.5	0
	Sweet Pepper in Rockwool	15.5	1.25	1.25	6.5	4.75	1.5	1.75	15	10	5	30	0.75	0.5	0
	Sweet Pepper in Rockwool (RDW)	12.75	1.25	1	5.75	3.25	1.125	1	15	10	4	25	0.75	0.5	0
	Lettuce in Recirculating Water	19	1.25	2	11	4.5	1	1.125	40	5	4	30	0.75	0.5	500
	Tomato in Rockwool	13.75	1.25	1.25	8.75	4.25	2	3.75	15	10	5	30	0.75	0.5	0
	Tomato in Rockwool (RDW)	10.75	1	1.25	6.5	2.75	1	1.5	15	10	4	20	0.75	0.5	0
Floriculture	Alstroemeria in Rockwool	11.25	1.25	1.25	6	2.875	1	1.25	25	10	4	25	0.75	0.5	0
	Alstroemeria in Rockwool (RDW)	7.5	0.75	1	4.75	2	0.75	1.25	25	5	4	20	0.75	0.5	0
	Anemone in Rockwool	13	1	1.5	6.5	3.75	1	1.25	35	5	4	30	0.75	0.5	0
	Carnation in Rockwool or Peat	13	1	1.25	6.25	3.75	1	1.25	25	10	4	30	0.75	0.5	0
	Carnation in Rockwool (reuse drainage)	7	0.75	0.8	4	1.625	0.6	0.7	20	5	3	20	0.5	0.5	0
	Anthurium andreanum in Rockwool or Peat	4.5	0.8	0.7	3	1	0.7	1	15	3	3	20	0.5	0.5	0
	Aster in Rockwool or Peat	13	1	1.25	6.25	3.75	1	1.25	25	10	4	30	0.75	0.5	0
	Bouvardia in Rockwool	13	1.25	1.75	6	4.25	1	1.5	25	5	3.5	20	0.75	0.5	0
	Bouvardia in Rockwool (RDW)	8	1	1.5	4	2.5	0.5	0.75	25	5	3.5	20	0.75	0.5	0
	Chrysanthemum in Recirculating Water	12.75	1.25	1	7.5	2.5	1	1	60	20	3	20	0.5	0.5	0
	Cymbidium in phenol foam	4.5	0.5	0.8	3	1.2	0.75	1.05	8	10	4	20	0.4	0.4	0
	Cymbidium in rockwool/urethane foam	4	1	0.8	2.8	1	0.75	1.25	8	10	4	20	0.4	0.4	0
	Euphorbia fulgens in Rockwool	11.5	1	1.5	6	3.5	1	1.5	35	10	3	20	0.5	0.5	0
	Freesia in Rockwool or Sand	14.5	1.25	1.25	7.75	3.375	1.5	1.5	25	10	4	25	0.75	0.5	0
	Gerbera in Rockwool	11.25	1.5	1.25	5.5	3	1	1.25	35	5	4	30	0.75	0.5	0
	Gerbera in Rockwool (RDW)	7.25	0.75	0.6	4.5	1.6	0.4	0.7	25	5	3	20	0.5	0.5	0
	Gypsophila in Rockwool	17	1.25	1.25	4	6	1.7	1.2	25	10	4	30	0.8	0.5	0
	Hippeastrum in Pumice	13	1	1.25	7.5	3.125	1	1.25	10	10	5	30	0.5	0.5	0
	Pot Plants in Expanded Clay	10.6	1.1	1.5	5.5	3	0.75	1	20	10	3	20	0.5	0.5	0
	Rose in Rockwool	11	1.5	1.25	4.5	3.25	1.125	1.25	25	5	3.5	20	0.75	0.5	0
	Rose in Rockwool (RDW)	4.3	0.85	0.5	2.15	0.9	0.5	0.5	15	5	3	15	0.5	0.5	0
	Statice in Rockwool	12	1	1	6	3	1	1	15	10	5	2.5	0.75	0.5	0
Fruticulture	Strawberry in Peaty Substrates	11.5	1	1	5.5	3.25	1.25	1.5	20	10	7	25	0.75	0.5	0
	Strawberry in Recirculating Water	10	0.5	1.25	5.25	2.75	1.125	1.125	20	10	4	20	0.75	0.5	0
	Melon in Rockwool	16.25	1	1.25	7.5	4.75	1.25	1.5	10	10	4	20	0.5	0.5	750

3.3 Data Structures and Chemical Representation

3.3.1 Elementary Entities

After the 2019 revision of the International System of Units, the base units are defined by fixed numerical values of fundamental constants (BIPM, 2019). The mole is the SI unit for the amount of a substance; its symbol is mol. The defining constant is the Avogadro constant, N_A . Thus:

$$1 \text{ mol} = \frac{6.02214076 \times 10^{23}}{N_A} \quad (3.1)$$

An elementary entity may be an atom, a molecule, or any other particle. In this work, the elementary entities are ions as listed in Table 3.2. The molar masses are taken from the CRC Handbook of Chemistry and Physics (Haynes, 2017) and rounded to three decimal places. Not all elements reported are essential in plant nutrition. The list include also non-essential or even toxic elements, but elements that part of the plant ionome and potentially present in fertilizers or water even though not all elements reported here where integrated later.

Table 3.2: Molar masses of elementary entities used for nutrient formulation.

Symbol	Element name	Molar Mass [g mol ⁻¹]
C	Carbon	12.011
O	Oxygen	15.999
H	Hydrogen	1.008
N	Nitrogen	14.007
P	Phosphorus	30.974
K	Potassium	39.098
Mg	Magnesium	24.305
S	Sulfur	32.065
Ca	Calcium	40.078
B	Boron	10.811
Cu	Copper	63.546
Cl	Chlorine	35.453
Fe	Iron	55.845
Mn	Manganese	54.938
Ni	Nickel	58.693
Zn	Zinc	65.380
Mo	Molybdenum	95.950
Na	Sodium	22.990
Al	Aluminum	26.982
Co	Cobalt	58.933
Se	Selenium	78.971
Si	Silicon	28.085

3.3.2 Nutrient set

In Table 3.3 all implemented 40 fertilizers accessible by the solver are listed with formula, name and respective molar mass. The molar mass is calculated from the chemical formula using the molar masses of the constituent elements and their stoichiometric coefficients reported in 3.2.

Table 3.3: Fertilizer compounds and their molar masses.

Formula	Name	Molar mass [g mol ⁻¹]
NH ₄ Cl	Ammonium chloride	53.492
NH ₄ H ₂ PO ₄	Ammonium dihydrogen phosphate	115.025
((NH ₄) ₆)Mo ₇ O ₂₄ · 4 H ₂ O	Ammonium heptamolybdate tetrahydrate	1235.920
NH ₄ OH	Ammonium hydroxide	35.046
NH ₄ NO ₃	Ammonium nitrate	80.043
(NH ₄) ₂ SO ₄	Ammonium sulfate	132.139
H ₃ BO ₃	Boric acid	61.832
CaCl ₂ · 2 H ₂ O	Calcium chloride dihydrate	147.014
Ca(OH) ₂	Calcium hydroxide	74.092
Ca(NO ₃) ₂	Calcium nitrate	164.086
Ca(NO ₃) ₂ · 4 H ₂ O	Calcium nitrate tetrahydrate	236.146
5 Ca(NO ₃) ₂ · NH ₄ NO ₃ · 10 H ₂ O	Calcium nitrate–ammonium nitrate decahydrate (5:1)	1080.623
CaSO ₄ · 2 H ₂ O	Calcium sulfate dihydrate	172.169
CuCl ₂	Copper(II) chloride	134.452
CuSO ₄ · 5 H ₂ O	Copper(II) sulfate pentahydrate	249.682
(NH ₄) ₂ HPO ₄	Diammonium hydrogen phosphate	132.056
HCl	Hydrochloric acid	36.461
C ₁₈ H ₁₆ N ₂ O ₆ FeNa	Iron(III) sodium EDDHA	435.169
C ₁₀ H ₁₂ N ₂ O ₈ FeNa · 3 H ₂ O	Iron(III) sodium EDTA trihydrate	421.092
Mg(NO ₃) ₂ · 6 H ₂ O	Magnesium nitrate hexahydrate	256.403
MgSO ₄ · 7 H ₂ O	Magnesium sulfate heptahydrate	246.471
MnCl ₂ · 4 H ₂ O	Manganese(II) chloride tetrahydrate	197.904
MnSO ₄ · H ₂ O	Manganese(II) sulfate monohydrate	169.014
KH ₂ PO ₄	Monopotassium dihydrogen phosphate	136.084
HNO ₃	Nitric acid	63.012
H ₃ PO ₄	Phosphoric acid	97.994
KCl	Potassium chloride	74.551
KOH	Potassium hydroxide	56.105
KNO ₃	Potassium nitrate	101.102
K ₂ SiO ₃	Potassium silicate	154.278
K ₂ SO ₄	Potassium sulfate	174.257
NaCl	Sodium chloride	58.443
NaOH	Sodium hydroxide	39.997
Na ₂ MoO ₄ · 2 H ₂ O	Sodium molybdate dihydrate	241.956
NaNO ₃	Sodium nitrate	84.994
Na ₂ B ₄ O ₇ · 10 H ₂ O	Sodium tetraborate decahydrate (borax)	381.367
H ₂ SO ₄	Sulfuric acid	98.077
ZnCl ₂	Zinc chloride	136.286
ZnSO ₄ · 7 H ₂ O	Zinc sulfate heptahydrate	287.546
ZnSO ₄ · 6 H ₂ O	Zinc sulfate hexahydrate	269.531

3.3.3 Compound-by-nutrient matrix

Table 3.4 shows the compound-by-nutrient matrix, listing the amount of each nutrient in mmol provided per mmol of compound for all 40 fertilizers. The table is organised with the chemical formula of each compound in the first column, followed by a column indicating the solver stage (1 for salts, 2 for acids and bases) and then columns for each nutrient. Chelates are also part of the full matrix, but are excluded from the table shown here for better readability. Each compound is additionally linked to metadata such as compound name, category and acid/base status.

Table 3.4: Compound–nutrient stoichiometric matrix (mmol nutrient per mmol compound).
Stage 1 = salts, Stage 2 = acids/bases.

Formula	Solver	NO3_N	NH4_N	P	K	Ca	Mg	S	Na	Cl	Fe	Mn	Zn	B	Cu	Mo	Si
Ca(NO ₃) ₂	1	2	0	0	0	1	0	0	0	0	0	0	0	0	0	0	0
NH ₄ NO ₃	1	1	1	0	0	0	0	0	0	0	0	0	0	0	0	0	0
KH ₂ PO ₄	1	0	0	1	1	0	0	0	0	0	0	0	0	0	0	0	0
KNO ₃	1	1	0	0	1	0	0	0	0	0	0	0	0	0	0	0	0
K ₂ SO ₄	1	0	0	0	2	0	0	1	0	0	0	0	0	0	0	0	0
MgSO ₄ · 7 H ₂ O	1	0	0	0	0	0	1	1	0	0	0	0	0	0	0	0	0
CaCl ₂ · 2 H ₂ O	1	0	0	0	0	1	0	0	0	2	0	0	0	0	0	0	0
KCl	1	0	0	0	1	0	0	0	0	1	0	0	0	0	0	0	0
CaSO ₄ · 2 H ₂ O	1	0	0	0	0	1	0	1	0	0	0	0	0	0	0	0	0
CuSO ₄ · 5 H ₂ O	1	0	0	0	0	0	0	1	0	0	0	0	0	0	1	0	0
(NH ₄) ₂ SO ₄	1	0	2	0	0	0	0	1	0	0	0	0	0	0	0	0	0
H ₃ BO ₃	1	0	0	0	0	0	0	0	0	0	0	0	0	1	0	0	0
MnSO ₄ · H ₂ O	1	0	0	0	0	0	0	1	0	0	0	1	0	0	0	0	0
Na ₂ MoO ₄ · 2 H ₂ O	1	0	0	0	0	0	0	0	2	0	0	0	0	0	0	1	0
ZnSO ₄ · 7 H ₂ O	1	0	0	0	0	0	0	1	0	0	0	0	1	0	0	0	0
NH ₄ H ₂ PO ₄	1	0	1	1	0	0	0	0	0	0	0	0	0	0	0	0	0
Ca(NO ₃) ₂ · 4 H ₂ O	1	2	0	0	0	1	0	0	0	0	0	0	0	0	0	0	0
Mg(NO ₃) ₂ · 6 H ₂ O	1	2	0	0	0	0	1	0	0	0	0	0	0	0	0	0	0
C ₁₀ H ₁₂ N ₂ O ₈ FeNa · 3 H ₂ O	1	0	0	0	0	0	0	0	1	0	1	0	0	0	0	0	0
ZnSO ₄ · 6 H ₂ O	1	0	0	0	0	0	0	1	0	0	0	0	1	0	0	0	0
Na ₂ B ₄ O ₇ · 10 H ₂ O	1	0	0	0	0	0	0	0	2	0	0	0	0	4	0	0	0
C ₁₈ H ₁₆ N ₂ O ₆ FeNa	1	0	0	0	0	0	0	0	1	0	1	0	0	0	0	0	0
K ₂ SiO ₃	1	0	0	0	2	0	0	0	0	0	0	0	0	0	0	0	1
NaCl	1	0	0	0	0	0	0	0	1	1	0	0	0	0	0	0	0
(NH ₄) ₂ HPO ₄	1	0	2	1	0	0	0	0	0	0	0	0	0	0	0	0	0
CuCl ₂	1	0	0	0	0	0	0	0	0	2	0	0	0	0	1	0	0
((NH ₄) ₆)Mo ₇ O ₂₄ · 4 H ₂ O	1	0	6	0	0	0	0	0	0	0	0	0	0	0	0	7	0
ZnCl ₂	1	0	0	0	0	0	0	0	0	2	0	0	1	0	0	0	0
MnCl ₂ · 4 H ₂ O	1	0	0	0	0	0	0	0	0	2	0	1	0	0	0	0	0
NH ₄ Cl	1	0	1	0	0	0	0	0	0	1	0	0	0	0	0	0	0
NaNO ₃	1	1	0	0	0	0	0	0	1	0	0	0	0	0	0	0	0
5 Ca(NO ₃) ₂ · NH ₄ NO ₃ · 10 H ₂ O	1	11	1	0	0	5	0	0	0	0	0	0	0	0	0	0	0
HNO ₃	2	1	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0
KOH	2	0	0	0	1	0	0	0	0	0	0	0	0	0	0	0	0
H ₂ SO ₄	2	0	0	0	0	0	0	1	0	0	0	0	0	0	0	0	0
HCl	2	0	0	0	0	0	0	0	0	1	0	0	0	0	0	0	0
H ₃ PO ₄	2	0	0	1	0	0	0	0	0	0	0	0	0	0	0	0	0
NaOH	2	0	0	0	0	0	0	0	1	0	0	0	0	0	0	0	0
NH ₄ OH	2	0	1	0	0	0	0	0	0	0	0	0	0	0	0	0	0
Ca(OH) ₂	2	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0

3.4 Algorithmic Methods

3.4.1 Formula parsing and molar mass calculation

Describes rules and algorithms for parsing chemical formulas and calculating molar masses, including validation and error handling.

```
library(nutricalc)
compute_molar_mass("Ca(NO3)2·4H2O")
```

```
[1] 236.146
```

3.4.2 Stoichiometric conversions

Numerically, the official SI unit for the substance concentration is mol m^{-3} ; however, this unit is equivalent to mmol l^{-1} . Therefore, the following relationships for conversion between mmol l^{-1} and mg l^{-1} were applied, following Appelo and Postma (Appelo, 2005):

$$\begin{aligned}\text{mg l}^{-1} &= \text{mmol l}^{-1} \times M \\ \text{mmol l}^{-1} &= \frac{\text{mg l}^{-1}}{M}\end{aligned}$$

where

M is the molar mass of the corresponding element $[\text{g mol}^{-1}]$.

Within the library, conversion between mmol l^{-1} and mg l^{-1} as elements can be accomplished with the `convert_units()` function. When the input is provided as a named numeric vector, the names are interpreted as nutrient identifiers (e.g., Ca, P, NO3_N) and are internally mapped to their corresponding element symbols to select the appropriate molar mass (from `element_molar_mass_df`) for conversion. For example, converting a complete set of recipe values from mg l^{-1} to mmol l^{-1} :

```
convert_units(c(Ca = 211, P = 46), to = "mmol/L")
```

```
      Ca      P  
5.264734 1.485117
```

The conversion can also be applied to a single scalar concentration. In this case, the element must be specified explicitly, since a scalar input carries no nutrient name from which the element can be inferred. The element may be supplied either as a named argument or positionally in the order (x, element, to):

```
convert_units(x = 15, element = "N", to = "mg/L")
```

```
[1] 210.105
```

For reporting and verification a recipe can be summarized with the `recipe_summary()` function:

```
recipe_summary(c(  
  NO3_N = 15,  
  NH4_N = 1,  
  P      = 1.5,  
  K      = 6,
```

```
Ca    = 4,
Mg    = 2,
S     = 2),
unit = "mmol/L")
```

	Nutrient	Element	mmol/L	mg/L
1	NO3_N	N	15.0	210.105
2	NH4_N	N	1.0	14.007
3	P	P	1.5	46.461
4	K	K	6.0	234.588
5	Ca	Ca	4.0	160.312
6	Mg	Mg	2.0	48.610
7	S	S	2.0	64.130

When concentrations are declared in mg l^{-1} , masses are often reported in oxidized or compound forms (e.g. SO_4 or MgO). To convert between such compound-based representations and their elemental equivalents, the `convert()` function can be used:

$$x_{\text{to}} = x_{\text{from}} \times \frac{M_{\text{to}}}{M_{\text{from}}} \times \frac{\nu_{\text{from},E}}{\nu_{\text{to},E}}$$

where

x_{from} is the quantity expressed as the source formula (input value),

x_{to} is the equivalent quantity expressed as the target formula (output value),

M_{from} is the molar mass of the source formula $[\text{g mol}^{-1}]$,

M_{to} is the molar mass of the target formula $[\text{g mol}^{-1}]$,

$\nu_{\text{from},E}$ is the number of atoms of the anchor element E in the source formula [dimensionless],

$\nu_{\text{to},E}$ is the number of atoms of the anchor element E in the target formula [dimensionless].

```
convert("SO4", "S")
```

```
[1] 0.3337983
```

It is also possible to specify an explicit quantity to be converted, for example when converting a given mass concentration

```
convert("N", "NO3", 210)
```

```
[1] 929.5952
```

If multiple elements are shared between the source and target formulas, the anchor element E must be specified to avoid ambiguity.

```
convert("NH4)2HP04", "NH4H2P04", element = "P") # P as the anchor
```

[1] 0.871032

3.4.3 NNLS optimization with weighting

NNLS is a numerical optimization algorithm for solving linear least-squares problems under non-negativity constraints. It was written in 1973 in FORTRAN 77 by Lawson and Hanson at the Jet Propulsion Laboratory and published in their book *Solving Least Squares Problems* (Lawson & Hanson, 1974, 1995). In R, the Lawson–Hanson algorithm is implemented by the package `nnls` (Mullen & van Stokkum, 2024).

For nutrient solution formulation, the objective is to determine the amounts of salts that fit a specified target nutrient composition as closely as possible. Each salt contributes fixed stoichiometric amounts of ions, so the relationship between the amount of each salt and the resulting nutrient concentration in solution is linear. Because salts can only be added and not removed, their amounts are constrained to be non-negative. Nutrient solution design can therefore be formulated as a linear mixture fitting problem subject to non-negativity constraints, which can be solved using the NNLS algorithm.

Because many fertilizer salts supply multiple nutrients simultaneously exact matching of all nutrient targets is rarely possible. The optimization therefore identifies the best feasible compromise among competing nutrient constraints.

Assume that n fertilizer salts are available and m nutrients are considered.

In `NutriCalc`, fertilizer stoichiometry is stored in a matrix

$$S \in \mathbb{R}^{n \times m},$$

where rows correspond to salts ($j = 1, \dots, n$) and columns correspond to nutrients ($i = 1, \dots, m$). The entry S_{ji} gives the amount of nutrient i supplied by one unit of salt j (mmol nutrient per mmol salt).

Let

- $x \in \mathbb{R}_{\geq 0}^n$ denote the vector of salt amounts (mmol L^{-1}),
- $b \in \mathbb{R}^m$ denote the vector of target nutrient concentrations (mmol L^{-1}).

For fitting, the solver uses the transposed stoichiometry matrix

$$A = S^\top \in \mathbb{R}^{m \times n},$$

so that rows correspond to nutrients and columns correspond to salts. The system therefore consists of m nutrient equations and n non-negative decision variables. Predicted (achieved) nutrient concentrations are

$$\hat{b} = Ax.$$

Thus, each row of A represents the linear contribution of all salts to one nutrient, and each column represents the nutrient composition of one salt. The optimization therefore selects salt amounts so that the combined nutrient profile best approximates the desired composition.

3.4.4 Target scaling and importance weighting

NutriCalc does not solve the unweighted least-squares problem $\min_{x \geq 0} \|Ax - b\|_2$. Instead, residuals are rescaled relative to target magnitude and weighted according to user-defined nutrient importance.

Conceptually, the solver minimizes the total weighted squared deviation between achieved and target nutrient concentrations. For nutrients with nonzero targets, deviations are measured relative to the target (i.e. proportional error). For nutrients with zero targets, deviations are measured in absolute concentration units.

This relative scaling is necessary because nutrient concentrations often differ by several orders of magnitude (for example, macronutrients versus micronutrients). Without scaling, nutrients present at large absolute concentrations would dominate the optimization and prevent meaningful fitting of low-concentration nutrients.

Scaling and importance weighting operate on residuals rather than on concentrations themselves. Optimization decisions therefore depend on deviations from targets rather than absolute concentration levels.

3.4.4.1 Importance weighting of nutrient targets

In practical nutrient formulation, not all targets must be matched with the same precision. Some nutrients are critical for plant performance and must be maintained close to their target concentration, whereas others can vary within a broader range without adverse effects. The optimization therefore allows nutrient-specific importance levels that determine how strongly deviations from individual targets are penalized.

User-defined importance values I_i express the desired strictness of matching each nutrient. Higher importance means that deviations from the target are penalized more strongly, forcing the solver to prioritize matching that nutrient even if this leads to larger deviations in less important nutrients.

Importance values are restricted to the interval $[-2, 2]$ and converted into positive numerical weights using an exponential mapping:

$$\begin{aligned}\tilde{I}_i &= \min(2, \max(-2, I_i)), \\ w_i &= 0.1 \cdot 1000^{\tilde{I}_i}.\end{aligned}$$

Each unit increase in importance multiplies the penalty on deviations by a factor of 1000. This creates clear separation between importance levels and ensures that highly important nutrients strongly influence the optimization outcome. This mapping produces weights spanning approximately 10^{-7} to 10^5 .

The weights are incorporated into the least-squares objective using the diagonal matrix

$$W = \text{diag}(\sqrt{w_1}, \dots, \sqrt{w_m}),$$

which scales the residuals before minimization. The optimization therefore minimizes a weighted sum of squared deviations, where each nutrient contributes proportionally to its assigned importance.

3.4.4.2 Target-based scaling of residuals

Nutrient targets differ substantially in magnitude, with macronutrients typically present at concentrations several orders of magnitude larger than micronutrients. If deviations were measured only in absolute units (mmol L^{-1}), nutrients with large target concentrations would dominate the optimization, and small but biologically important deviations in low-concentration nutrients would have little influence on the solution.

To avoid this imbalance, residuals are scaled relative to target magnitude. For each nutrient i , define a scaling factor

$$s_i = \begin{cases} b_i, & b_i \neq 0, \\ 1, & b_i = 0. \end{cases}$$

For nutrients with nonzero targets, deviations are therefore measured relative to the target value. For nutrients with zero targets, deviations are measured in absolute concentration units.

To apply this scaling, each nutrient equation is divided by its corresponding factor s_i . In matrix form, the target-scaled design matrix is

$$A_{\text{sub}} = \text{diag}(1/s) A,$$

and the scaled target vector is b/s , where division is performed elementwise.

The NNLS problem is then

$$\min_{x \geq 0} \|W(A_{\text{sub}}x - b/s)\|_2^2.$$

This is equivalent to minimizing a weighted sum of squared scaled residuals. For nutrients with nonzero targets, the optimization minimizes weighted relative deviations; for nutrients with zero targets, it minimizes weighted absolute deviations. The scaling ensures that all nutrients contribute comparably to the optimization regardless of their concentration range, while importance weights control how strictly individual targets are enforced.

Operationally, the solver predicts nutrient concentrations from candidate salt amounts, computes deviations from the targets, rescales and weights these deviations according to target magnitude and nutrient importance and determines the non-negative salt combination that minimizes the resulting weighted least-squares objective.

3.4.5 Interpretation of the solution

The NNLS solver returns the optimal non-negative salt amounts x . Because the optimization is performed on scaled and weighted equations, achieved nutrient concentrations are recomputed using the original, unscaled stoichiometry:

$$\hat{b} = Ax.$$

The vector x therefore represents the physically meaningful fertilizer composition and is interpreted directly as mmol L^{-1} of each salt to add to the nutrient solution. These values can therefore be used directly to prepare the nutrient solution.

3.4.6 Error quantities reported by NutriCalc

After computing achieved nutrient concentrations, deviations from the target composition are quantified using several complementary error measures. These metrics describe both absolute and relative discrepancies between predicted and desired nutrient levels.

The absolute error for nutrient i is

$$e_i = \hat{b}_i - b_i,$$

expressed in mmol L^{-1} . This measures the direct concentration difference between achieved and target values.

Percent error expresses the deviation relative to the target concentration:

$$p_i = 100 \cdot \frac{e_i}{b_i}.$$

Percent error is computed elementwise. Non-finite values (e.g. division by zero) are recorded as missing. If $b_i = 0$ and $|\hat{b}_i| < 10^{-9}$, the percent error is set to 0.

To summarize overall deviation across all nutrients, sum of squared error is computed as

$$\text{SSE} = \sum_{i=1}^m e_i^2.$$

This metric reflects the total magnitude of absolute concentration differences.

Relative squared error evaluates deviations relative to target magnitude. Relative residuals are defined as

$$r_i = \frac{e_i}{b_i},$$

and non-finite values are omitted. The aggregate measure is reported as the sum of squared relative errors

$$\text{SSRE} = \sum_{i=1}^m r_i^2.$$

This metric summarizes proportional deviations across nutrients and provides a scale-independent measure of solution accuracy. Nutrients with zero target concentration where excluded from relative error metrics.

3.4.7 Two-stage NNLS solver: separation of neutral salts and acid–base reagents

In addition to the standard optimization, NutriCalc provides an optional two-stage solver that incorporates acid–base reagents. These compounds are included in the global stoichiometry matrix and distinguished from neutral fertilizer salts using the classification `salt_info$is_acid_base`. The purpose of this separation is to ensure that nutrient targets are satisfied using neutral salts first and whenever possible, while acids and bases are used only to correct remaining deficits that neutral salts cannot resolve. This prevents acid–base reagents from acting as unconstrained degrees of freedom in the primary optimization and yields chemically more interpretable fertilizer compositions.

The two-stage procedure is activated only when the acid/base option is enabled. When this option is disabled, acid/base compounds are excluded from the selected compound set and nutrient

optimization is performed using neutral salts only. Boric acid belongs to the neutral group in this classification, as it is treated as a micronutrient source rather than a pH-active reagent.

In the first stage, NNLS optimization is performed using only compounds classified as neutral salts. This stage uses the same weighted and target-scaled objective as the standard solver and produces achieved nutrient concentrations

$$\hat{b}^{(1)}.$$

Residual nutrient deviations are then computed as

$$r = b - \hat{b}^{(1)}.$$

Residual entries are filtered before any second-stage correction. Deviations within the relative tolerance are treated as negligible and set to zero. Residuals are also set to zero for nutrients that cannot be supplied by any acid or base (non-fixable nutrients). Negative residuals (oversupply) are set to zero so that only undersupplied nutrients remain eligible for correction.

If the two-stage mode is enabled and at least one fixable nutrient exceeds the tolerance criterion defined by

$$r_i > \max(\text{tol_abs}, \text{tol_rel } b_i),$$

a second NNLS optimization is performed using only acid/base compounds. The target for this stage is the filtered residual vector from stage 1. Scaling and importance weighting are applied using the same rules as in the standard solver, but computed from the residual target rather than the original nutrient targets. Stage 2 therefore supplies only remaining nutrient deficits that neutral salts could not provide.

Compound amounts from stage 1 and stage 2 are then inserted into their original compound indices to form the complete fertilizer vector (x). Final nutrient concentrations are recomputed from the full unscaled stoichiometry matrix,

$$\hat{b} = Ax,$$

and all error metrics are evaluated from these final concentrations.

Accordingly, the two-stage solver acts as a conditional refinement: neutral salts determine the primary nutrient composition, and acid–base reagents are introduced only when residual deficits exceeding the tolerance criterion remain after the neutral fit. Lowering the importance of a specific nutrient reduces the penalty for deviating from its target during stage 1, allowing a residual deficit to remain primarily in that nutrient while the remaining nutrients are matched more closely. Because

stage-2 optimization operates only on residual undersupply, this deficit can then be supplied during the second stage by the corresponding acid–base compounds.

3.5 Chemical Modeling

3.5.1 Water quality and input handling

Nutrient formulation was implemented using a target-based optimization workflow in NutriCalc. Users specify desired final nutrient concentrations as a nutrient target vector. The solver then computes the fertilizer salt additions needed to match this target. This approach is exact only when the initial water matrix is essentially deionized or demineralized water. In practical cultivation systems, however, source water already contains dissolved ions that contribute to the final solution and must therefore be accounted for before fertilizer optimization.

To represent this baseline contribution, water chemistry is modeled as a water vector containing pre-existing ion concentrations. The formulation step uses an adjusted target vector obtained by subtracting the baseline water vector from the desired nutrient target vector. For each nutrient ion i , the solver input is defined as:

$$T_i^{adj} = \max(T_i^{target} - W_i, 0)$$

where T_i^{target} is the desired final concentration and W_i is the concentration already present in the source water. Negative adjusted concentrations are not physically meaningful because fertilizer salts can only be added. Therefore, adjusted targets are bounded below by zero. The lower bound at zero reflects the additive-only fertilizer model used in the optimizer; excess baseline ions cannot be removed within this step. The resulting adjusted vector is then passed to the salt optimizer.

In addition to ionic concentrations, two buffering descriptors are important for water characterization: acid capacity (Säurekapazität, KS 4.3) and base capacity (Basenkapazität, KB 8.2). These are not treated as primary fertilizer targets, but are retained as water properties because they influence downstream interpretation of pH behavior.

Because users may operate with blended feed waters, NutriCalc supports mixing of multiple source waters prior to target subtraction. Each source is represented as a separate composition vector and the composite baseline is computed by mass balance. For two source waters with compositions W_1 and W_2 and volumes V_1 and V_2 , the mixed baseline composition is:

$$W^{mix} = \frac{V_1 W_1 + V_2 W_2}{V_1 + V_2}.$$

This expression is evaluated component-wise for all dissolved constituents of the water vector. The adjusted nutrient target is subsequently computed as:

$$T^{adj} = \max(T^{target} - W^{mix}, 0).$$

This enables realistic handling of scenarios such as blending municipal water with deionized water, rainwater, or other process waters before nutrient dosing.

Water quality inputs were derived from publicly available authority datasets where possible. For Berlin, NutriCalc retrieves supplier data and maps it to the internal water-vector representation. In practice, users can query location-specific baseline water data by postal code and optionally modify values to reflect local measurements or temporal variability.

Overall, this method ensures that fertilizer optimization is performed against the true chemical starting point rather than an idealized zero-background assumption. By explicitly modeling baseline composition, optional source-water mixing and buffering-related capacities, the resulting formulations are more representative of real cultivation solutions and suitable for subsequent pH and EC evaluation.

3.5.2 Estimation of pH via charge balance

Solution pH is calculated using `ph_from_achieved()` based on the achieved nutrient composition returned by the solver (mmol L^{-1}). The calculation is performed as a charge-balance problem in which pH is treated as an unknown variable. The hydrogen ion concentration is determined such that the aqueous solution satisfies electroneutrality, expressed in milliequivalents per litre (meq L^{-1}), such that total positive and negative charge are equal (Appelo, 2005).

The calculation is formulated as a root-finding problem in which the charge-balance residual is expressed as a function of pH. The root variable is defined as

$$\text{pH}_c = -\log_{10}([\text{H}^+]).$$

For each trial value of pH_c , equilibrium speciation is computed and the resulting charge imbalance is evaluated. The solution pH corresponds to the value of pH_c for which the charge residual equals zero within numerical tolerance.

Strong ions (e.g., K^+ , Na^+ , Ca^{2+} , Mg^{2+} , NO_3^- , Cl^-) contribute with fixed concentrations from the achieved nutrient composition, whereas variable ions (e.g., H^+ , OH^- , NH_4^+) are determined from equilibrium speciation. Weak acid–base systems (e.g., sulfate, phosphate, carbonate, borate, silicate) are included in the speciation model and contribute to the charge balance according to their equilibrium distributions.

Carbonate speciation is parameterized either from a carbonate-system input treated as total inorganic carbon when provided (e.g., `KS4_3`, interpreted approximately as HCO_3^-), or from fixed dissolved $\text{CO}_2(\text{aq})$ when provided (e.g., `KB8_2`, interpreted approximately as CO_2). Optional chelating ligands (e.g., for Fe) are included as fixed-charge anions when ligand totals are present (default net charges: EDTA, EDDHA, HBED $z = -4$ and DTPA $z = -5$).

Acid–base equilibria are evaluated using thermodynamic dissociation constants at 25 °C (Haynes,

2017). The speciation model includes the following equilibrium systems:

System	Equilibrium species
Water	H^+, OH^-
Ammonia	$\text{NH}_4^+, \text{NH}_3$
Sulfate	$\text{HSO}_4^-, \text{SO}_4^{2-}$
Phosphate	$\text{H}_3\text{PO}_4, \text{H}_2\text{PO}_4^-, \text{HPO}_4^{2-}, \text{PO}_4^{3-}$
Carbonate	$\text{CO}_2(\text{aq}), \text{HCO}_3^-, \text{CO}_3^{2-}$
Borate	$\text{B}(\text{OH})_3, \text{B}(\text{OH})_4^-$
Silicate	$\text{H}_4\text{SiO}_4, \text{H}_3\text{SiO}_4^-$

Non-ideal solution behavior is represented using Davies activity coefficients at 25 °C, using the Debye–Hückel parameter for water ($A = 0.5085$). Ionic strength is computed as

$$I = \frac{1}{2} \sum c_i z_i^2.$$

For each trial pH_c , ionic strength and activity coefficients are determined iteratively because they depend on the speciation they influence. Ionic strength is computed from species concentrations, activity coefficients are updated from ionic strength and speciation is recalculated. This fixed-point iteration is repeated until successive estimates converge. The resulting activity coefficients are then applied to all equilibrium expressions.

After convergence of the inner speciation loop, the charge residual is evaluated as total positive equivalents minus total negative equivalents. Positive charge includes fixed cations and variable H^+ and NH_4^+ . Negative charge includes fixed anions, OH^- and the equivalent contributions of sulfate, phosphate, carbonate, borate and silicate species, together with optional fixed chelate charge.

The bisection search is initialized over a bounded pH interval (default 3–9), reflecting the typical range of horticultural nutrient solutions and improving numerical stability. If the charge-balance function is not bracketed within this interval, the search range is expanded automatically.

The reported pH is activity-based,

$$\text{pH} = -\log_{10}(\gamma_H[\text{H}^+]),$$

and therefore may differ slightly from pH_c . All calculations assume 25 °C and exclude precipitation reactions and explicit metal–ligand complexation beyond optional fixed-charge chelate terms.

3.5.3 EC estimation using ionic diffusion coefficients

Electrical conductivity (EC) quantifies the ability of an aqueous solution to transport electrical charge and is therefore determined by the concentration and mobility of dissolved ions. Several

approaches exist for estimating EC from chemical composition, ranging from simple empirical relationships to ion-specific physicochemical models.

In practical greenhouse fertilization, EC is often estimated from the total ionic charge concentration expressed in milliequivalents per liter (meq L⁻¹). A widely used rule of thumb for greenhouse nutrient solutions is (van der Lugt et al., 2020):

$$\text{EC (mS cm}^{-1}\text{)} = \frac{\sum \text{meq cations} + \sum \text{meq anions}}{20}.$$

Because aqueous solutions satisfy electroneutrality,

$$\sum \text{meq cations} \approx \sum \text{meq anions},$$

this relationship can also be interpreted as a proportionality between EC and total ionic charge concentration. A closely related approximation for dilute aqueous solutions is described by Appelo (2005), where conductivity in $\mu\text{S cm}^{-1}$ is approximately proportional to total milliequivalents per liter (typically valid below about 1.5 mS cm⁻¹):

$$\sum \text{meq ions} \approx \frac{\text{EC } (\mu\text{S cm}^{-1})}{100}.$$

More refined empirical relationships have been proposed for greenhouse nutrient solutions with relatively constrained ionic composition. Based on experimental calibration, Sonneveld & Straver (1994) proposed the following regression relating EC to total cation concentration:

$$\text{EC} = 0.095 \cdot \sum \text{meq cations} + 0.19,$$

where EC is expressed in mS cm⁻¹.

Although such empirical formulations may improve predictive accuracy within their calibration range, they do not explicitly account for differences in ionic mobility among individual species. Because the ionic species present in solution are known in `NutriCalc`, EC can be estimated mechanistically from their concentration, charge and mobility. The limiting molar ionic conductivity can be related to the diffusion coefficient through the Nernst–Einstein relation (Kalka, 2020). In `NutriCalc`, this relation is extended by a mobility correction term accounting for non-ideal solution behavior. This method is used in hydrochemical programs such as PHREEQC (up to version 3.8) and AQION 8.7.15 (Appelo, 2010; Kalka, 2020).

$$\text{EC} = \left(\frac{F^2}{RT} \right) \sum_i D_i z_i^2 (\gamma_i)^{\alpha_i} c_i. \quad (6)$$

where

F is Faraday's constant [$9.6485 \times 10^4 \text{ C mol}^{-1}$],

R is the gas constant [$8.31446 \text{ J mol}^{-1} \text{ K}^{-1}$],

T is absolute temperature [$T = t_C + 273.15 \text{ K}$],

D_i is the diffusion coefficient of ion i [$\text{m}^2 \text{ s}^{-1}$],

z_i is the ionic charge number,

γ_i is the activity coefficient of ion i ,

α_i is an empirical mobility scaling exponent,

c_i is the molar concentration of ion i .

Concentrations used in the conductivity summation are expressed in mol m^{-3} . Concentrations obtained in mmol L^{-1} are numerically identical to mol m^{-3} and are therefore used directly without conversion. For conductivity, concentrations are expressed in mol m^{-3} whereas ionic strength for activity modeling is calculated from concentrations expressed in mol L^{-1} . Diffusion coefficients in the internal data tables are stored in units of $10^{-5} \text{ cm}^2 \text{ s}^{-1}$ at 25°C and converted to SI units prior to calculation:

$$1 \times 10^{-5} \text{ cm}^2 \text{ s}^{-1} = 1 \times 10^{-9} \text{ m}^2 \text{ s}^{-1}.$$

The activity of a dissolved species depends not only on its concentration but also on the ionic strength of the solution (Amelung et al., 2018). Ionic strength (I) is calculated for activity modeling from molar concentrations expressed in mol L^{-1} :

$$I = \frac{1}{2} \sum_i z_i^2 c_i.$$

Because the classical Nernst–Einstein relation applies strictly at infinite dilution, a mobility scaling factor is introduced to account empirically for inter-ionic interactions at finite ionic strength as described by Appelo (2010). The scaling exponent is defined as

$$\alpha_i = \begin{cases} \frac{0.6}{\sqrt{|z_i|}}, & I \leq 0.36|z_i| \\ \frac{\sqrt{I}}{|z_i|}, & I > 0.36|z_i| \end{cases}$$

providing an ionic-strength-dependent attenuation of individual ionic contributions.

Activity coefficients (γ_i) are obtained using the selected activity model implemented in NutriCalc. Two models are currently implemented: the limiting Debye–Hückel model and the Davies model (Kalka, 2023). In the limiting Debye–Hückel model, the activity coefficient is given by

$$-\log_{10}(\gamma_i) = Az_i^2\sqrt{I},$$

whereas in the Davies model it is given by

$$-\log_{10}(\gamma_i) = Az_i^2 \left(\frac{\sqrt{I}}{1 + \sqrt{I}} - 0.3I \right),$$

where $A = 0.5085$ at 25 °C, z_i is the ionic charge number and I is the ionic strength in mol L⁻¹.

The limiting Debye–Hückel formula is typically applied at ionic strengths up to 0.1 mol L⁻¹ whereas the Davies formula is generally considered applicable up to about 0.5 mol L⁻¹. Currently the Davies activity correction is used as the default in the `NutriCalc` Shiny app.

The calculation is applied to the fully speciated aqueous ion composition obtained after nutrient formulation and equilibrium pH calculation. All dissolved charged species, including hydrogen and hydroxide ions, contribute to the conductivity sum according to their diffusion properties and concentrations.

Diffusion coefficients of ionic species at 25 °C were retrieved from Haynes (2017). The approach assumes fully dissociated strong electrolytes and does not account for neutral species or metal–ligand complexes. Consequently, the model provides a physically based estimate of conductivity that resolves species-specific contributions but may deviate from measured EC where complexation, ion pairing or strong ion association is significant.

The osmotic potential is estimated from the calculated EC value using the empirical relationship $\Psi_{\text{osm}} = -36 \cdot \text{EC}$, where Ψ_{osm} is expressed in kPa and EC in mS cm⁻¹. This relationship is given by George et al. (2012) based on United States Salinity Laboratory Staff (1954), where it was originally formulated as a relation between osmotic pressure and the electrical conductivity of the saturation extract.

3.6 Shiny Application Workflow

3.6.1 Installation or launching the web application

To install and run the application locally, the package `nutricalc` can be installed from GitHub using the package `remotes` and the following code:

```
install.packages("remotes")
remotes::install_github("diolsa/nutricalc")
library(nutricalc)
run_app()
```

Alternatively, the deployed Shiny application can be accessed online without local installation at <https://nutricalc.shinyapps.io/nutricalc-app/>.

Once launched, the application opens with the title `NutriCalc` `Nutrient Solution Formulation in Horticultural Sciences`.

3.6.2 Input parameters

The left-hand panel of the Shiny graphical user interface (GUI), titled `Nutrient Targets & Fertilizers`, comprises five tabs: `Nutrient Targets`, `Water`, `Fertilizers`, `Recipes`, and `Tissue`.

3.6.2.1 Nutrient Targets

Within the `Nutrient Targets` tab, each nutrient is represented by three user-editable controls: a `Target` input field, a `Water` input field, and a `Priority` selector. The `Target` field defines the desired concentration in the final nutrient solution, whereas the `Water` field specifies the concentration contributed by the source water. Input values can be displayed and entered in mmol L^{-1} , mol L^{-1} , or mg L^{-1} . In addition, alkalinity-related water parameters can be entered using the `KS 4.3` and `KB 8.2` fields. The `Priority` control is implemented as a slider ranging from -2 to $+2$ and specifies the relative importance of each nutrient during optimization.

At the bottom of the panel, several global controls are available for workflow management. The `Set all to 0` button resets all target and water input values to zero, whereas `Reset to defaults` restores the predefined default settings. The `Use acids/bases` option activates the secondary optimization stage after completion of the initial solver step, thereby allowing the inclusion of acids and bases for further adjustment. Finally, the `Result` button initiates the solver and opens the calculated formulation output.

3.6.2.2 Water

The `Water` tab is used to define and modify the source-water composition. Mean water-analysis values for Berlin postal code areas (PLZ) from BWB (2026) can be retrieved by entering a valid postal code (in the range from 10115 to 14199). Using `Load Water`, the retrieved values are transferred directly to the `Water` input fields in the `Nutrient Targets` tab. Alternatively, `Mix Water` transfers the retrieved PLZ values to `Water 1` within the water-mixing interface.

Within the mixing interface, two separate source waters, `Water 1` and `Water 2`, can be defined. Concentrations for both sources can be entered manually. If one of the two sources is left empty, it is assumed to be deionized water. A slider located above the two input areas is used to specify the mixing ratio between `Water 1` and `Water 2`. In this way, blends of different water sources, such as tap water, drainage water, rainwater, aquaculture effluent, or other custom nutrient-containing solutions, can be simulated. By selecting `Load Mixed Water`, the composition of the resulting mixture is transferred to the `Water` input fields in the `Nutrient Targets` tab.

3.6.2.3 Fertilizers

The **Fertilizers** tab is used to define the fertilizer salts, acids and bases available for optimization. Compounds are displayed in a list together with their chemical formulas and names.

At the top of the tab, a **Salt search** text field is provided to filter the list. Both chemical formulas and compound names can be entered and the visibility of the listed compounds updates dynamically according to the search term. A **Clear search** button is located to the right of the search field and resets the search input, thereby restoring the full list.

Compounds can be selected or deselected manually using the checkbox list. In addition, the **Select all visible** and **Deselect all visible** buttons allow batch selection or deselection of compounds currently shown in the filtered list. Thus, these controls operate on the subset of compounds visible under the active search term rather than on the complete compound list.

Below the list, a status line provides summary information in the form **Using x selected of y visible (total z)**, where x denotes the number of currently selected compounds, y the number of currently visible compounds and z the total number of available compounds.

3.6.2.4 Recipes

The **Recipes** tab provides a collection of crop-specific and standard nutrient solutions used in horticulture. It is divided into four categories: **Olericulture**, **Fruticulture**, **Floriculture**, and **Standard Formulations**. Most recipes in the first three categories are based on Sonneveld & Straver (1994), whereas the fourth category contains formulations compiled from Hewitt (1966). The compilation by Hewitt (1966) represents one of the most comprehensive collections of hydroponic nutrient-solution formulations published in the twentieth century.

Once a **Category** is selected, the corresponding **Recipe** list is displayed. By selecting a recipe and using the **Apply recipe** button, the respective values are transferred to the **Target** input fields in the **Nutrient Targets** tab. For recipes reported together with specific fertilizer salts, selection of the recipe can also modify the set of compounds selected in the **Fertilizers** tab. In addition, recipes are stored in different concentration units, with some reported in mmol L^{-1} and others in mg L^{-1} .

3.6.2.5 Tissue

The **Tissue** tab allows formulation of a nutrient solution based on the mass-balance approach described by Langenfeld et al. (2022).

Tissue nutrient concentrations can be entered on a dry-mass basis and displayed in either % or ppm. In the central part of the tab, water-use efficiency (WUE) is calculated as dry mass divided by transpired water, with dry mass entered in g and transpired water in L. The resulting WUE value is displayed below the corresponding input fields.

Based on tissue nutrient concentration and WUE, nutrient demand per unit of transpired water is estimated and used to derive the corresponding nutrient-solution composition for analysis. In addition to the original approach of Langenfeld et al. (2022), total nitrogen is partitioned into nitrate and ammonium in accordance with the principle of electroneutrality. This extension is discussed in a later section.

3.6.3 Output parameters

The right-hand side of the Shiny graphical user interface (GUI) contains the output panel, which is displayed after the solver is initiated. The panel is divided into five tabs: Result, pH and EC, Solution, Anions and Cations.

3.6.3.1 Result

The Result tab summarizes the optimization output by comparing the calculated nutrient composition with the user-defined target composition. For each nutrient, the table reports the values specified in the Water vector, the Target vector, and the values calculated by the solver in Achieved. In addition, a Final value is reported, representing the sum of the Water contribution and the Achieved contribution. Concentrations are displayed in mol L^{-1} for values less than or equal to 0.1 mmol L^{-1} and in mmol L^{-1} for values above this threshold.

Deviation from the target composition is quantified using the error metrics Abs. Error and Pct. Error, representing the absolute deviation and the relative deviation in percent of Final from Target, respectively.

To evaluate the overall goodness of fit across all nutrients, deviations are additionally aggregated as squared error terms. Absolute deviations are summarized as the sum of squared errors (SSE), whereas relative deviations are summarized as the sum of squared relative errors (SSRE).

3.6.3.2 pH & EC

The pH & EC tab summarizes the calculated pH and EC values based on the Final composition. Calculations are performed for 25°C using Davies activity correction, while complex formation is not considered. On the left side of the tab, the calculated pH, ionic strength (mmol L^{-1}), and total charge equivalents (meq L^{-1}) are reported. By selecting Show pH charge breakdown, an expanded view of the pH charge balance can be displayed, showing the contributions of fixed cations, fixed anions, and variable components in meq L^{-1} .

On the right side of the tab, EC is reported in both mS cm^{-1} and $\mu\text{S cm}^{-1}$. From the calculated EC, osmotic potential is additionally derived and reported in kPa. By selecting Show EC contributions, the individual ionic contributions to the calculated EC can be displayed. For each ion, the table reports the concentration c (mM), charge number z , and the contribution to conductivity κ (mS cm^{-1}).

Above the pH and EC output sections, an Adjust concentration slider is provided with a range from -100% to +100%. This control scales the concentration of the reported Final composition and thereby allows interactive assessment of the resulting changes in pH, EC, and related derived parameters.

3.6.3.3 Solution

The Solution tab reports the compounds used to obtain the Achieved composition. For each salt, acid, or base, the table reports the applied amount in mmol L^{-1} , the calculated molar mass in g mol^{-1} , and the corresponding concentration in mg L^{-1} .

Above the Fertilizer amounts table, input fields are provided for working volume, stock factor, and stock volume. Based on these inputs, additional columns are computed for the required amount for the selected working-solution volume ($x \text{ L}$) and, when valid stock parameters are provided, for the selected stock preparation ($y \text{ L}$ of z -fold stock).

Compounds are grouped according to the BAK assignment logic into A-BAK, B-BAK, and Micro. Assignment is rule-based: compounds containing micronutrients are assigned to Micro; compounds containing Ca, but no micronutrients, are assigned to A-BAK; compounds containing P or S, but neither micronutrients nor Ca, are assigned to B-BAK. Remaining compounds (AB) are assigned dynamically to A-BAK or B-BAK in order to balance the overall nutrient load between both tanks.

3.6.3.4 Anions and Cations

The Anions and Cations tabs report the relative proportions of selected anionic or cationic nutrient components on a mmol L^{-1} basis using ternary diagrams. In the Anions tab, the default components are $\text{NO}_3\text{-N}$, P, and S, whereas in the Cations tab the default components are K, Ca, and Mg. Alternative ions can also be selected for comparison, including Cl for anions and $\text{NH}_4\text{-N}$ or Na for cations. The respective proportions of Achieved and Target are displayed directly in the ternary graph, and the corresponding percentage values are reported in the legend.

3.7 Case study selection

Three case studies were selected to validate the performance of the application across a range of use cases. For this purpose, three user types with different objectives were defined.

3.7.1 Case Study I: Alternative Hoagland II nutrient solution formulation

In this scenario, a plant scientist needs to prepare a classical Hoagland nutrient solution for an experiment using deionized water. The intended formulation is the macronutrient solution of Hoagland solution No. 2 according to Hoagland & Arnon (1938), consisting of 6 mM KNO_3 , 4 mM $\text{Ca}(\text{NO}_3)_2$, 1 mM $\text{NH}_4\text{H}_2\text{PO}_4$, and 2 mM $\text{MgSO}_4 \cdot 7\text{H}_2\text{O}$. The scientist requires the solution at several concentration levels corresponding to target EC values of 0.5, 1.0, 1.5, 2.0, 2.5, and 3.0 mS cm^{-1} . However, the standard calcium nitrate salt is unavailable and only calcium ammonium nitrate (CAN; $5\text{Ca}(\text{NO}_3)_2 \cdot \text{NH}_4\text{NO}_3 \cdot 10\text{H}_2\text{O}$) is available as a substitute.

3.7.2 Case Study II: Tomato nutrient solution using Berlin-Dahlem water with acid adjustment

In this scenario, a horticultural technician needs to prepare a nutrient solution for tomato cultivation in rockwool using technical-grade fertilizers and tap water from Berlin-Dahlem (postal code 14195) instead of deionized water. The reference formulation is the tomato recipe reported by Sonneveld & Straver (1994). Because the tap water contributes background ions and alkalinity, the two-stage solver is applied with nitric acid permitted in the second stage for acid adjustment. Calculations are performed for the preparation of 500 L nutrient solution and for 18 L of a 50-fold concentrated stock solution separated into A-BAK, B-BAK, and micronutrients.

3.7.3 Case Study III: Nutrient solution with increased sodium concentration for halophytes using mixed water sources

In this scenario, a horticultural scientist needs to formulate a nutrient solution for an experiment on sodium effects in *Salicornia europaea* using tap water from Berlin postal code 10997. Sodium treatments in the literature are frequently imposed using NaCl , but chloride has to be limited to 1.5 mmol L^{-1} . As the chloride concentration of the tap water exceeds this threshold, the tap water is mixed with demineralised water. The base recipe “Propagation vegetable plants in rockwool” reported by Sonneveld & Straver (1994) is used as the reference formulation. Ammonium nitrate (NH_4NO_3) is excluded from the set of permissible fertilizers. All calculations are performed for the preparation of 5 L stock solution.

4 Results

The following result section is divided in two parts. The first part a general evaluation of the solver for a set of crop specific recipes and the performance of pH and EC calculation. This is followed by three case studies dealing with specific solving problems and are intended to showcase different scenarios and the practical applicability of the shiny app.

4.1 General evaluation across crop specific recipes

To evaluate solver performance, a set of published crop specific nutrient targets from Sonneveld & Straver (1994) were used as model input. These recipes specify target nutrient concentrations for various crops in mmol L^{-1} . The solver was only applied allowing salts to each recipe (no acids or bases) to compute the optimal fertilizer composition and achieved nutrient concentrations and were compared to the specified targets. The following sections summarize the results of this evaluation, including achieved nutrient profiles, error metrics, pH and EC. In addition, because Sonneveld et al. (1999) and Sonneveld (2009) used one reference formulation to demonstrate a universal nutrient solution algorithm, the NutriCalc solution for this formulation was compared in detail with the published results.

4.1.1 Achieved Nutrient solution formulations

All reported recipes (list of recipes is found in Table 3.1) were solved and achieved results are summarized in Table 4.1. Squared absolute and relative errors are reported for each recipe, quantifying the deviation between achieved and target nutrient concentrations across all nutrients. All achieved results matched the target nutrient profiles with SSE and SSRE values close to zero. Recipes including Si as a target nutrient exhibited comparatively higher relative error values. In formulations using Fe, achieved Na concentrations were non-zero although Na was not specified in the literature target vectors and Na contributed to the absolute error. When Na was added to the target vector at the achieved concentration (Na target set equal to Fe), the corresponding SSE values decreased to zero. For recipes containing Si, applying the second solver approach with reduced weighting for $\text{NO}_3\text{-N}$ resulted in solutions with an SSRE of 0.

Table 4.1: Achieved Nutrient Solutions with SSE and SSRE. Asterisks denote two-stage optimization; parentheses indicate standard-solver errors.

Category	Recipe	mmol L ⁻¹							μmol L ⁻¹							Squared absolute error	Squared relative error	
		NO ₃ -N	NH ₄ -N	P	K	Ca	Mg	S	Na	Fe	Mn	Zn	B	Cu	Mo			Si
Olericulture	Endive in Recirculating Water	19	1.25	2	9	5	1.5	1.13	39.94	39.94	5	4	30	0.75	0.5	0	0.0016	0
	Eggplant in Rockwool	15.5	1.5	1.25	6.75	3.25	2.5	1.5	15	15	10	5	30	0.75	0.5	0	0.0002	0
	Eggplant in Rockwool (RDW)	11.75	1	1	6.5	2.25	1.5	1.12	15	15	10	5	20	0.75	0.5	0	0.0002	0
	Bean in Rockwool	12	1	1.25	5.5	3.25	1.25	1.12	10	10	10	4	20	0.5	0.5	0	0.0001	0
	Courgette in Rockwool	16	1.25	1.25	7.25	3.62	2	1.25	10	10	10	5	30	0.75	0.5	0	0.0001	0
	Cucumber in Rockwool*	16	1.25	1.25	8	4	1.38	1.37	15	15	10	5	25	0.75	0.5	750	0.0002 (0.9325)	0 (0.0053)
	Cucumber in Rockwool (RDW)*	11.75	1	1.25	6.5	2.75	1	1	15	15	10	5	25	0.75	0.5	750	0.0002 (0.9122)	0 (0.0097)
	Propagation vegetable plants in rockwool	16.75	1.25	1.25	6.75	4.5	3	2.5	24.98	24.98	10	5	35	1	0.5	0	0.0006	0
	Sweet Pepper in Rockwool	15.5	1.25	1.25	6.5	4.75	1.5	1.75	15	15	10	5	30	0.75	0.5	0	0.0002	0
	Sweet Pepper in Rockwool (RDW)	12.75	1.25	1	5.75	3.25	1.13	1	15	15	10	4	25	0.75	0.5	0	0.0002	0
	Lettuce in Recirculating Water*	19	1.25	2	11	4.5	1	1.12	39.94	39.94	5	4	30	0.75	0.5	500	0.0016 (0.423)	0 (0.0017)
	Tomato in Rockwool	13.75	1.25	1.25	8.75	4.25	2	3.75	15	15	10	5	30	0.75	0.5	0	0.0002	0
Tomato in Rockwool (RDW)	10.75	1	1.25	6.5	2.75	1	1.5	15	15	10	4	20	0.75	0.5	0	0.0002	0	
Floriculture	Alstroemeria in Rockwool	11.25	1.25	1.25	6	2.87	1	1.25	24.98	24.98	10	4	25	0.75	0.5	0	0.0006	0
	Alstroemeria in Rockwool (RDW)	7.5	0.75	1	4.75	2	0.75	1.25	24.98	24.98	5	4	20	0.75	0.5	0	0.0006	0
	Anemone in Rockwool	13	1	1.5	6.5	3.75	1	1.25	34.96	34.96	5	4	30	0.75	0.5	0	0.0012	0
	Carnation in Rockwool or Peat	13	1	1.25	6.25	3.75	1	1.25	24.98	24.98	10	4	30	0.75	0.5	0	0.0006	0
	Carnation in Rockwool (reuse drainage)	7	0.75	0.8	4	1.62	0.6	0.7	19.99	19.99	5	3	20	0.5	0.5	0	0.0004	0
	Anthurium andreanum in Rockwool or Peat	4.5	0.8	0.7	3	1	0.7	1	15	15	3	3	20	0.5	0.5	0	0.0002	0
	Aster in Rockwool or Peat	13	1	1.25	6.25	3.75	1	1.25	24.98	24.98	10	4	30	0.75	0.5	0	0.0006	0
	Bouvardia in Rockwool	13	1.25	1.75	6	4.25	1	1.5	24.98	24.98	5	3.5	20	0.75	0.5	0	0.0006	0
	Bouvardia in Rockwool (RDW)	8	1	1.5	4	2.5	0.5	0.75	24.98	24.98	5	3.5	20	0.75	0.5	0	0.0006	0
	Chrysanthemum in Recirculating Water	12.75	1.25	1	7.5	2.5	1	1	59.78	59.78	20	3	20	0.5	0.5	0	0.0036	0
	Cymbidium in phenol foam	4.5	0.5	0.8	3	1.2	0.75	1.05	8	8	10	4	20	0.4	0.4	0	0.0001	0
	Cymbidium in rockwool/urethane foam	4	1	0.8	2.8	1	0.75	1.25	8	8	10	4	20	0.4	0.4	0	0.0001	0
	Euphorbia fulgens in Rockwool	11.5	1	1.5	6	3.5	1	1.5	34.96	34.96	10	3	20	0.5	0.5	0	0.0012	0
	Freesia in Rockwool or Sand	14.5	1.25	1.25	7.75	3.37	1.5	1.5	24.98	24.98	10	4	25	0.75	0.5	0	0.0006	0
	Gerbera in Rockwool	11.25	1.5	1.25	5.5	3	1	1.25	34.96	34.96	5	4	30	0.75	0.5	0	0.0012	0
	Gerbera in Rockwool (RDW)	7.25	0.75	0.6	4.5	1.6	0.4	0.7	24.98	24.98	5	3	20	0.5	0.5	0	0.0006	0
	Gypsophila in Rockwool	17	1.25	1.25	4	6	1.7	1.2	24.98	24.98	10	4	30	0.8	0.5	0	0.0006	0
	Hippeastrum in Pumice	13	1	1.25	7.5	3.12	1	1.25	10	10	10	5	30	0.5	0.5	0	0.0001	0
	Pot Plants in Expanded Clay	10.6	1.1	1.5	5.5	3	0.75	1	19.99	19.99	10	3	20	0.5	0.5	0	0.0004	0
	Rose in Rockwool	11	1.5	1.25	4.5	3.25	1.12	1.25	24.98	24.98	5	3.5	20	0.75	0.5	0	0.0006	0
Rose in Rockwool (RDW)	4.3	0.85	0.5	2.15	0.9	0.5	0.5	15	15	5	3	15	0.5	0.5	0	0.0002	0	
Statice in Rockwool	12	1	1	6	3	1	1	15	15	10	5	2.5	0.75	0.5	0	0.0002	0	
Fruticulture	Strawberry in Peaty Substrates	11.5	1	1	5.5	3.25	1.25	1.5	19.99	19.99	10	7	25	0.75	0.5	0	0.0004	0
	Strawberry in Recirculating Water	10	0.5	1.25	5.25	2.75	1.12	1.13	19.99	19.99	10	4	20	0.75	0.5	0	0.0004	0
	Melon in Rockwool*	16.25	1	1.25	7.5	4.75	1.25	1.5	10	10	10	4	20	0.5	0.5	750	0.0001 (0.8753)	0 (0.005)

4.2 pH and EC Outputs

Table 4.2: Solver-derived pH, ionic strength, total charge equivalents (TCE), electrical conductivity (EC) and osmotic pressure for crop-specific formulations (salts-only; water = 0; KS 4.3 = 0; CO2(aq) = 0). EC sonneveld and reported EC for comparison.

Category	Recipe	pH	Ionic strength (mmol L ⁻¹)	TCE (meq L ⁻¹)	EC model (mS cm ⁻¹)	EC Sonneveld (mS cm ⁻¹)	EC reported (mS cm ⁻¹)	Osmotic pressure (kPa)
Olericulture	Endive in Recirculating Water	5.16	31.46	23.44	2.6	2.42	2.6	-93.45
	Eggplant in Rockwool	5.47	27.28	19.85	2.22	2.08	2.1	-79.75
	Eggplant in Rockwool (RDW)	5.57	20.15	15.09	1.76	1.62	1.7	-63.21
	Bean in Rockwool	5.45	21.34	15.57	1.77	1.67	1.7	-63.88
	Courgette in Rockwool	5.47	26.84	19.83	2.23	2.07	2.2	-80.31
	Cucumber in Rockwool	8.64	28.61	20.37	2.28	2.13	2.2	-81.97
	Cucumber in Rockwool (RDW)	8.72	21.61	15.4	1.76	1.65	1.7	-63.43
	Propagation vegetable plants in rockwool	5.46	33.41	23.14	2.49	2.39	2.4	-89.63
	Sweet Pepper in Rockwool	5.46	28.53	20.35	2.25	2.12	2.2	-81.15
	Sweet Pepper in Rockwool (RDW)	5.55	21.4	15.84	1.82	1.7	1.7	-65.64
	Lettuce in Recirculating Water	6.99	31.83	23.78	2.67	2.45	2.6	-96.04
	Tomato in Rockwool	5.46	32.78	22.6	2.48	2.34	2.3	-89.22
	Tomato in Rockwool (RDW)	5.46	20.53	15.09	1.75	1.62	1.6	-62.89
Floriculture	Alstroemeria in Rockwool	5.46	20.53	15.13	1.75	1.63	1.7	-62.85
	Alstroemeria in Rockwool (RDW)	5.43	15.38	11.12	1.31	1.25	1.1	-47.06
	Anemone in Rockwool	5.27	23.52	17.17	1.94	1.82	1.9	-69.94
	Carnation in Rockwool or Peat	5.46	23.16	16.88	1.92	1.79	1.8	-69.07
	Carnation in Rockwool (reuse drainage)	5.47	12.43	9.3	1.11	1.07	1.1	-40.12
	Anthurium andreanum in Rockwool or Peat	5.45	10.13	7.28	0.88	0.88	0.8	-31.61
	Aster in Rockwool or Peat	5.46	23.16	16.88	1.92	1.79	1.8	-69.07
	Bouvardia in Rockwool	5.2	24.89	17.88	2	1.89	1.9	-71.87
	Bouvardia in Rockwool (RDW)	5.27	15.13	11.12	1.29	1.25	1.2	-46.37
	Chrysanthemum in Recirculating Water	5.73	21.15	16.04	1.87	1.71	1.8	-67.47
	Cymbidium in phenol foam	5.67	10.58	7.46	0.89	0.9	0.8	-31.99
	Cymbidium in rockwool/urethane foam	5.67	10.48	7.36	0.88	0.89	0.8	-31.79
	Euphorbia fulgens in Rockwool	5.36	22.53	16.17	1.83	1.73	1.7	-65.8
	Freesia in Rockwool or Sand	5.45	25.53	18.88	2.15	1.98	2.1	-77.37
	Gerbera in Rockwool	5.33	20.76	15.16	1.74	1.63	1.7	-62.65
	Gerbera in Rockwool (RDW)	5.58	12.32	9.37	1.14	1.08	1.1	-41.04
	Gypsophila in Rockwool	5.44	29.96	20.78	2.24	2.16	2.2	-80.49
	Hippeastrum in Pumice	5.47	22.34	16.82	1.95	1.79	1.9	-70.16
	Pot Plants in Expanded Clay	5.37	19.19	14.21	1.64	1.54	1.6	-58.94
	Rose in Rockwool	5.32	20.76	14.87	1.69	1.6	1.6	-60.77
	Rose in Rockwool (RDW)	5.67	7.94	5.88	0.72	0.75	0.7	-25.89
	Statice in Rockwool	5.57	20.28	15.09	1.75	1.62	1.7	-63
Fruticulture	Strawberry in Peaty Substrates	5.62	21.85	15.62	1.78	1.67	1.7	-63.99
	Strawberry in Recirculating Water	5.47	18.84	13.61	1.56	1.48	1.5	-56.1
	Melon in Rockwool	8.73	29.87	20.89	2.3	2.17	2.2	-82.98

Table 4.3: pH and EC comparison across activity-coefficient models (salts-only; water = 0; KS 4.3 = 0; CO2(aq) = 0). EC Sonneveld is the linear estimate from TCE (meq/L).

Category	Recipe	pH Davies	pH DH	I Davies (mmol/L)	I DH (mmol/L)	TCE Davies (meq/L)	TCE DH (meq/L)	EC Davies (mS/cm)	EC DH (mS/cm)	EC Sonneveld (mS/cm)	EC reported (mS/cm)
Olericulture	Bean in Rockwool	5.45	5.42	21.34	21.34	15.57	15.57	1.82	1.77	1.67	1.7
	Courgette in Rockwool	5.47	5.43	26.84	26.84	19.83	19.83	2.31	2.23	2.07	2.2
	Cucumber in Rockwool	8.64	8.65	28.61	28.62	20.37	20.38	2.36	2.28	2.13	2.2
	Cucumber in Rockwool (RDW)	8.72	8.72	21.61	21.62	15.4	15.4	1.81	1.76	1.65	1.7
	Eggplant in Rockwool	5.47	5.43	27.28	27.28	19.85	19.85	2.29	2.22	2.08	2.1
	Eggplant in Rockwool (RDW)	5.57	5.54	20.15	20.15	15.09	15.09	1.8	1.76	1.62	1.7
	Endive in Recirculating Water	5.16	5.12	31.46	31.46	23.44	23.44	2.69	2.6	2.42	2.6
	Lettuce in Recirculating Water	6.99	6.93	31.83	31.83	23.78	23.78	2.77	2.67	2.45	2.6
	Propagation vegetable plants in rockwool	5.46	5.41	33.41	33.41	23.14	23.14	2.6	2.49	2.39	2.4
	Sweet Pepper in Rockwool	5.46	5.42	28.53	28.53	20.35	20.35	2.34	2.25	2.12	2.2
	Sweet Pepper in Rockwool (RDW)	5.55	5.51	21.4	21.4	15.84	15.84	1.87	1.82	1.7	1.7
	Tomato in Rockwool	5.46	5.41	32.78	32.78	22.6	22.6	2.59	2.48	2.34	2.3
	Tomato in Rockwool (RDW)	5.46	5.43	20.53	20.53	15.09	15.09	1.79	1.75	1.62	1.6
Floriculture	Alstroemeria in Rockwool	5.46	5.43	20.53	20.53	15.13	15.13	1.79	1.75	1.63	1.7
	Alstroemeria in Rockwool (RDW)	5.43	5.41	15.38	15.38	11.12	11.12	1.33	1.31	1.25	1.1
	Anemone in Rockwool	5.27	5.24	23.52	23.52	17.17	17.17	2	1.94	1.82	1.9
	Anthurium andreaeanum in Rockwool or Peat	5.45	5.44	10.13	10.14	7.28	7.28	0.89	0.88	0.88	0.8
	Aster in Rockwool or Peat	5.46	5.42	23.16	23.16	16.88	16.88	1.98	1.92	1.79	1.8
	Bouvardia in Rockwool	5.2	5.17	24.89	24.89	17.88	17.88	2.06	2	1.89	1.9
	Bouvardia in Rockwool (RDW)	5.27	5.25	15.13	15.13	11.12	11.12	1.31	1.29	1.25	1.2
	Carnation in Rockwool (reuse drainage)	5.47	5.46	12.43	12.43	9.3	9.3	1.13	1.11	1.07	1.1
	Carnation in Rockwool or Peat	5.46	5.42	23.16	23.16	16.88	16.88	1.98	1.92	1.79	1.8
	Chrysanthemum in Recirculating Water	5.73	5.7	21.15	21.15	16.04	16.04	1.92	1.87	1.71	1.8
	Cymbidium in phenol foam	5.67	5.65	10.58	10.58	7.46	7.46	0.9	0.89	0.9	0.8
	Cymbidium in rockwool/urethane foam	5.67	5.65	10.48	10.48	7.36	7.36	0.9	0.88	0.89	0.8
	Euphorbia fulgens in Rockwool	5.36	5.33	22.53	22.53	16.17	16.17	1.88	1.83	1.73	1.7
	Freesia in Rockwool or Sand	5.45	5.41	25.53	25.53	18.88	18.88	2.22	2.15	1.98	2.1
	Gerbera in Rockwool	5.33	5.31	20.76	20.76	15.16	15.17	1.79	1.74	1.63	1.7
	Gerbera in Rockwool (RDW)	5.58	5.56	12.32	12.32	9.37	9.37	1.16	1.14	1.08	1.1
	Gypsophila in Rockwool	5.44	5.4	29.96	29.96	20.78	20.78	2.33	2.24	2.16	2.2
	Hippeastrum in Pumice	5.47	5.44	22.34	22.34	16.82	16.83	2	1.95	1.79	1.9
	Pot Plants in Expanded Clay	5.37	5.34	19.19	19.19	14.21	14.21	1.68	1.64	1.54	1.6
	Rose in Rockwool	5.32	5.29	20.76	20.76	14.87	14.87	1.73	1.69	1.6	1.6
	Rose in Rockwool (RDW)	5.67	5.66	7.94	7.94	5.88	5.88	0.73	0.72	0.75	0.7
	Statice in Rockwool	5.57	5.54	20.28	20.28	15.09	15.09	1.79	1.75	1.62	1.7
Fruticulture	Melon in Rockwool	8.73	8.73	29.87	29.88	20.89	20.9	2.4	2.31	2.17	2.2
	Strawberry in Peaty Substrates	5.62	5.58	21.85	21.85	15.62	15.62	1.83	1.78	1.67	1.7
	Strawberry in Recirculating Water	5.47	5.44	18.84	18.84	13.61	13.61	1.6	1.56	1.48	1.5

4.2.0.1 Performance of pH Calculation

Both γ activity correction models (Davies and Debye–Hückel) were evaluated for pH prediction against the phosphate buffer system (50 mL of 0.1 mol L⁻¹ KH₂PO₄ + x mL of 0.1 mol L⁻¹ NaOH), which spans pH 5.8–8.0 (Bower & Bates, 1955; Haynes, 2017). In Figure 4.1, predicted pH values are plotted against the reference pH for increasing NaOH additions at 25 °C. The Davies model closely matched the reference values ($R^2 = 0.998$), whereas the Debye–Hückel model showed weaker agreement ($R^2 = 0.880$).

Diskussion: Consequently, γ Davies was selected as the default activity correction for pH prediction in NutriCalc.




Figure 4.1: Comparison of 50 mL of 0.1 mol L⁻¹ KH₂PO₄ + x mL of 0.1 mol L⁻¹ NaOH (25 °C).

4.2.0.2 Performance of EC Calculation

To evaluate EC calculation performance, models were compared against reference calibration EC values for KCl solutions at 25 °C. Electrical conductivity values for 0.01 and 0.1 mol L⁻¹ KCl were taken from Haynes (2017), which reports values based on primary conductivity standards originally published by Pratt et al. (2001). In Table 4.4, reference values are compared with NutriCalc EC predictions using both γ activity-correction models (Davies and Debye–Hückel). For comparison, the empirical approach by Sonneveld et al. (1999) and geochemical computations with PHREEQC (Parkhurst & Appelo, 2013) are also shown. PHREEQC and NutriCalc with γ Davies were very close to the values reported by Haynes (2017) at both concentrations. NutriCalc with γ Debye–Hückel slightly underestimated EC, while the underestimation was larger for the Sonneveld empirical approach.

Table 4.4: Electrical conductivity of KCl solutions at 25 °C.

KCl (mol L ⁻¹)	$\mu\text{S cm}^{-1}$				
	CRC	Sonneveld	NutriCalc Debye–Hückel	NutriCalc Davies	PHREEQC
0.01	1408	1140	1396	1408	1409
0.10	12825	9690	11996	12923	12988

4.2.1 Comparison with published formulation

The nutrient solution algorithm described by Sonneveld (2009) uses the cucumber recipe with reuse of drainage water published in Sonneveld & Straver (1994) as a demonstration of a universal nutrient-solution formulation method proposed in Sonneveld et al. (1999). The target macronutrient concentrations for this demonstration are 11.75 mmol L⁻¹ NO₃-N, 1.00 mmol L⁻¹ NH₄-N, 1.25 mmol L⁻¹ P, 6.50 mmol L⁻¹ K, 2.75 mmol L⁻¹ Ca, 1.00 mmol L⁻¹ Mg, and 1.00 mmol L⁻¹ S. Micronutrients are not included because Sonneveld (2009) reports fertilizer additions only for macronutrients in this example. For a direct comparison, NutriCalc was constrained to the same set of salts reported by Sonneveld (2009). The published Sonneveld formulation includes nitric acid. NutriCalc matched the targets exactly (SSE = 0; SSRE = 0) using the one-stage solver without requiring acid addition. The same error metrics (SSE and SSRE) were computed for both the published Sonneveld formulation and the NutriCalc solution and are shown in Table 4.5. For the composition of nutrient salts used for the two formulations, see Table 4.6.

Table 4.5: Comparison of Sonneveld formulation and NutriCalc solution.

Nutrient	Target	Sonneveld solution			NutriCalc solution		
		Achieved	SE	SRE	Achieved	SE	SRE
NO ₃ -N	11.75	14.96	10.3041	0.0746	11.75	0.0000	0.0000
NH ₄ -N	1.00	1.25	0.0625	0.0625	1.00	0.0000	0.0000
P	1.25	1.50	0.0625	0.0400	1.25	0.0000	0.0000
K	6.50	8.88	5.6644	0.1341	6.50	0.0000	0.0000
Ca	2.75	2.47	0.0784	0.0104	2.75	0.0000	0.0000
Mg	1.00	1.07	0.0049	0.0049	1.00	0.0000	0.0000
S	1.00	0.87	0.0169	0.0169	1.00	0.0000	0.0000
		SSE = 16.1937		SSRE = 0.3434	SSE = 0.0000		SSRE = 0.0000

tets testtest

Table 4.6: Comparison of fertilizer additions in the Sonneveld and NutriCalc formulations.

Name	Formula	mmol L ⁻¹	
		Sonneveld	NutriCalc
Calcium nitrate–ammonium nitrate decahydrate (5:1)	5 Ca(NO ₃) ₂ ·NH ₄ NO ₃ ·10 H ₂ O	0.49	0.55
Ammonium nitrate	NH ₄ NO ₃	0.76	0.45
Monopotassium phosphate	KH ₂ PO ₄	1.50	1.25
Magnesium nitrate hexahydrate	Mg(NO ₃) ₂ ·6 H ₂ O	1.07	1.00
Potassium sulfate	K ₂ SO ₄	0.87	1.00

Nitric acid	HNO_3	1.00	—
Potassium nitrate	KNO_3	5.64	3.25

4.3 Case Study Results

4.3.1 Case study 1 — Experimental scientist: classical solution reconstruction under restricted salt availability (deionized water) under different EC.

The macronutrient composition of Hoagland solution No. 2 consists of 14 mmol L⁻¹ NO₃-N, 1 mmol L⁻¹ NH₄-N, 1 mmol L⁻¹ P, 6 mmol L⁻¹ K, 4 mmol L⁻¹ Ca, 2 mmol L⁻¹ Mg and 2 mmol L⁻¹ S. In the absence of Ca(NO₃)₂ three exemplary alternative formulations using the technical salt CAN are shown in Table 4.7.

Table 4.7: Comparison of the original Hoagland solution No. 2 fertilizer combination and three alternative fertilizer combinations achieving the same target macronutrient composition. Values are given in mmol L⁻¹.

Salt	Original	Alternative 1	Alternative 2	Alternative 3
Ca(NO ₃) ₂	4.00	—	—	—
5Ca(NO ₃) ₂ ·NH ₄ NO ₃ ·10H ₂ O	—	0.80	0.80	0.80
KNO ₃	6.00	1.20	1.00	1.20
MgSO ₄ ·7H ₂ O	2.00	—	—	—
Mg(NO ₃) ₂ ·6H ₂ O	—	2.00	2.00	2.00
NH ₄ H ₂ PO ₄	1.00	—	—	0.20
KH ₂ PO ₄	—	1.00	1.00	0.80
K ₂ SO ₄	—	1.90	2.00	2.00
NH ₄ NO ₃	—	—	0.20	—
(NH ₄) ₂ SO ₄	—	0.10	—	—
Number of salts	4	6	6	6

4.3.1.1 Achieved composition

The target macronutrient composition of Hoagland solution No. 2 was achieved using 0.80 mmol L⁻¹ 5Ca(NO₃)₂·NH₄NO₃·10H₂O, 0.80 mmol L⁻¹ KH₂PO₄, 5.20 mmol L⁻¹ KNO₃, 2.00 mmol L⁻¹ MgSO₄·7H₂O and 0.20 mmol L⁻¹ NH₄H₂PO₄. This is shown as Alternative 3 in Table 4.7. For the different EC Designs values are shown in Table 4.8.

Table 4.8: Amounts of fertilizer salts required to prepare modified Hoagland solution No. 2 at target EC values from 0.5 to 3.0 mS cm⁻¹. Salt amounts are given in mmol L⁻¹.

Salt / parameter	Concentration in mmol L ⁻¹ for target EC (mS cm ⁻¹)					
	EC 0.5	EC 1.0	EC 1.5	EC 2.0	EC 2.5	EC 3.0
Strength (5Ca(NO ₃) ₂ ·NH ₄ NO ₃ ·10H ₂ O	0.17	0.34	0.54	0.73	0.92	1.12
KH ₂ PO ₄	0.17	0.34	0.54	0.73	0.92	1.12
KNO ₃	1.09	2.24	3.48	4.73	5.98	7.28
MgSO ₄ ·7H ₂ O	0.42	0.86	1.34	1.82	2.30	2.80
NH ₄ H ₂ PO ₄	0.04	0.09	0.13	0.18	0.23	0.28

4.3.1.2 Deviations from target

No deviation from the target composition of a Hoagland solution No. 2 was observed in the achieved solutions, resulting in an SSE and SSRE of 0.00 equally for the original formulation and for the alternative formulations with different fertilizer salts.

4.3.1.3 pH and EC

4.3.2 Case study 2 — Horticultural technician: tomato nutrient formulation using Berlin-Dahlem water with acid adjustment

4.3.2.1 Achieved composition

For a nutrient solution formulated for tomatoes cultivated in rockwool (Sonneveld & Straver, 1994) and prepared with tap water from postal code 14195 (BWB, 2026), the two-stage solver with nitric acid addition was used. The calculated fertilizer salts are shown in Figure 4.2, including the amounts required for the preparation of 500 L nutrient solution and the corresponding amounts for 18 L of a 50-fold concentrated stock solution separated into A-BAK, B-BAK and micronutrients.

4.3.2.2 Deviations from target

The tomato recipe specifies a target composition without Na, Cl, or Si. In the calculated solution, these elements were present due to their contribution from the tap water. Although the target composition was matched with an SSRE of 0, absolute deviations remained for Na, Cl, and Si, resulting in an SSE of 7.69. The absolute errors were 2.03 mmol L⁻¹ for Na, 1.86 mmol L⁻¹ for Cl, and 0.30 mmol L⁻¹ for Si. Of the Na concentration, 0.015 mmol L⁻¹ originated from Fe-EDTA.

4.3.2.3 pH and EC

The tap water from postal code 14195 had a calculated pH of 7.44 and an EC of 0.86 mS cm⁻¹. Besides the ions considered in the target nutrient composition, it supplied 2.02 mmol L⁻¹ Na⁺ and 1.86 mmol L⁻¹ Cl⁻, contributing 0.09 and 0.13 mS cm⁻¹, respectively, to the total EC. The electroneutral 0.3 mmol L⁻¹ Si present in the tap water did not contribute to the overall EC.

The calculated final nutrient solution had a pH of 5.82. Ionic strength amounted to 34.84 mmol L⁻¹, while the total concentration of charge equivalents was 24.61 meq L⁻¹. The calculated EC was 2.81 mS cm⁻¹, and the osmotic potential was -101.02 kPa. In the final solution, the largest ionic contribution to EC was from NO₃⁻ at 0.89 mS cm⁻¹, followed by K⁺ at 0.58 mS cm⁻¹ and SO₄²⁻ at 0.45 mS cm⁻¹.




Figure 4.2: Screenshot of the fertilizer salt composition calculated in the NutriCalc Shiny app for a tomato nutrient solution in rockwool using tap water from Berlin postal code 14195. The formulation is shown separately for A-BAK, B-BAK, and micronutrients stock solutions. Fertilizer requirements are reported for the preparation of 500 L nutrient solution and the amounts for 18 L of a 50-fold concentrated stock solution.

4.3.3 Case study 3 — Research scientist: nutrient formulation for a novel halophyte (*Salicornia*) using mixed water sources

4.3.3.1 Achieved composition

The calculated mixed water consists of 90 % tap water from Berlin postal code 10997 and 10 % demineralised water, thereby reducing the chloride concentration of the water source to 1.5 mmol L⁻¹. Based on this water composition and the reference recipe “Propagation vegetable plants in rockwool” reported by Sonneveld & Straver (1994), chloride is set to 1.5 mmol L⁻¹, yielding a calculated sodium concentration of 5.5 mmol L⁻¹ to maintain electroneutrality. Since ammonium nitrate (NH₄NO₃) is excluded from the permissible fertilizer set, the target composition is achieved using an alternative fertilizer combination. The final fertilizer amounts are expressed for the preparation of 5 L stock solution.




Figure 4.3: Screenshot of the fertilizer salt composition calculated in the NutriCalc Shiny app for a tomato nutrient solution in rockwool using tap water from Berlin postal code 14195. The formulation is shown separately for A-BAK, B-BAK, and micronutrients stock solutions. Fertilizer requirements are reported for the preparation of 500 L nutrient solution and the amounts for 18 L of a 50-fold concentrated stock solution.

4.3.3.2 Deviations from target

4.3.3.3 pH and EC

5 Discussion

5.1 Solver robustness and formulation performance

Across the evaluated formulations, the solver reproduced most published target compositions exactly or with only very small residual deviations. This indicates that the implemented fertilizer matrix is sufficiently expressive for a broad range of horticultural nutrient solutions. In particular, the crop specific targets from Sonneveld & Straver (1994) were matched with SSE and SSRE values close to zero in most cases, showing that the combination of stoichiometric matrix construction and normalised NNLS optimisation is capable of representing many practically relevant formulation problems with high fidelity.

Where deviations remained, they were chemically interpretable rather than arbitrary numerical failures. This is important because it means that residual error in the present framework should not immediately be interpreted as an algorithmic weakness. Instead, it often reflects specific properties of the fertilizer representation or of the target vector itself.

Salts used in the solver were selected on the basis of their solubility and their reported use in hydroponics. Salts that are not water soluble but may still be relevant in soil based systems were therefore not included in the implemented salt matrix.

The small SSE values reported for the crop set of Sonneveld & Straver (1994) are mainly caused by the use of chelated Fe sources in the salt matrix. In the implemented representation, Fe chelates introduce Na as a counter ion. Accordingly, a non zero Na contribution can occur in the achieved solution even when Na is absent from the published target vector. This should not be interpreted as a substantive mismatch in the nutritionally relevant targets, but rather as a consequence of using chemically realistic Fe fertilizers. Consistent with this interpretation, SSE drops to zero when the Na target is set equal to the Fe associated Na contribution.

A second recurring case involves recipes containing Si, commonly at around 0.75 mmol L^{-1} . In the present matrix, Si is supplied via K_2SiO_3 , which necessarily introduces additional K and also behaves as a strongly basic reagent. Consequently, exact matching depends on whether this coupled K contribution is represented in the target vector and or corrected by acid addition. When Si is omitted, or when the corresponding K contribution is accounted for explicitly, SSRE approaches zero.

This indicates that the residual mismatch in Si containing recipes is not primarily caused by numerical instability of the optimiser, but by chemically coupled fertilizer behaviour. In other words, the difficulty lies less in the NNLS procedure itself than in the fact that silicon supply through

potassium silicate cannot be treated independently of its accompanying alkalinity and counter ion contribution, because it behaves effectively as a base used in the role of a salt.

Although Si is generally considered a beneficial rather than an essential element, positive effects on yield and disease suppression have been reported (Rengel et al., 2023; Voogt & Sonneveld, 2001). Practical use of potassium silicate often requires stoichiometric acid compensation. In the formulations discussed by Voogt & Sonneveld (2001), this was addressed by combining silicate addition with nitric acid and with corresponding adjustment of nitrate salts.

The fact that Si containing cases improved when acid and base correction was allowed in NutriCalc therefore supports rather than undermines the chemical plausibility of the framework. Taken together, the results suggest that NutriCalc performs robustly within the tested horticultural domain and that the remaining deviations can usually be traced back to identifiable chemical causes.

5.2 Exact solvability, electroneutrality and non uniqueness

In this context, an electroneutral target is defined as a target vector for which the sum of cation and anion charge equivalents is balanced. A non electroneutral target is a target vector for which this condition is not fulfilled. Electroneutrality is a necessary but not sufficient condition for exact solvability when only electrically neutral salts are considered. Exact solvability also depends on the selected fertilizer set.

If the target is electroneutral and the available fertilizers span the required composition, the solver may find an exact solution. In such cases, both SSE and SSRE are zero, indicating that all targets have been matched exactly in both absolute and relative terms. However, this does not imply that the solution is unique. Different fertilizer combinations may produce the same final nutrient composition (Steiner, 1961). Some alternative solutions differ only in the hydration state of a salt, for example $\text{Ca}(\text{NO}_3)_2$ versus $\text{Ca}(\text{NO}_3)_2 \cdot 4\text{H}_2\text{O}$, whereas others may involve chemically distinct fertilizer combinations. In some cases, the number of exact solutions may be effectively infinite because different salts can compensate for one another while preserving the same final nutrient profile.

Accordingly, the NNLS solver returns one admissible non negative optimum, but it does not characterise the full set of possible exact solutions. Users can nevertheless influence the returned solution by restricting the fertilizer set through the selection and deselection of salts. In addition, small fertilizer amounts may appear in the solution if they provide even a minor reduction in the objective function. Thus, an exact or near exact solution is not necessarily the chemically simplest or most practical one.

If the target is non electroneutral, it cannot be matched exactly using only electrically neutral salts. In that case, the solver returns the non negative best fit solution that minimises the residual error. Because NutriCalc normalises residuals by target magnitude, the optimisation criterion primarily minimises the sum of squared relative errors for non zero targets and the sum of squared absolute errors for zero targets.

A useful implication of this formulation is that infeasible targets are not solved in an intuitively arithmetic sense, but according to the relative structure of the objective function. When a target cannot be matched exactly, the remaining mismatch is distributed according to the row scaling and the imposed weights. Thus, residual error should not be interpreted simply as failure. It may instead indicate that the target is not electroneutral, that the selected fertilizer set does not span the target composition, or that chemically realistic fertilizer sources introduce unavoidable side contributions.

5.3 Error metrics and interpretation of residual deviation

Reported errors quantify the deviation between target and achieved nutrient concentrations. For each nutrient, the formulation output reports the absolute error and, when the target is non zero, the relative error. Two aggregate metrics are derived: the sum of squared errors, SSE, and the sum of squared relative errors, SSRE. The per nutrient errors indicate which nutrients deviate from the target, by how much and in which direction. The squared error metrics summarise overall solution quality.

The optimisation itself is based on a row normalised non negative least squares, NNLS, formulation. Residuals are scaled by a nutrient specific factor: for nutrients with non zero targets, deviations are normalised by the corresponding target concentration. For nutrients with zero targets, a scaling factor of one is applied. As a result, all nutrients, including those with targets set to zero, are actively included in the objective function. Any non zero achieved concentration for a zero target nutrient contributes a quadratic penalty and is therefore discouraged during optimisation.

SSE reflects the total squared deviation across all nutrients and therefore captures leakage into zero target nutrients. SSRE, in contrast, summarises squared relative deviations only for nutrients with non zero targets and serves as the primary performance indicator for how closely specified targets were matched. However, both SSE and SSRE are post hoc reporting metrics. The solver decision is determined by the normalised and weighted NNLS objective. Consequently, a formulation with slightly lower relative deviation for non zero targets may still be rejected if it introduces substantial unintended supply of zero target nutrients, because this increases the overall objective value.

Without normalisation, an objective based solely on squared absolute errors would bias the optimisation toward macronutrients, since deviations in the $\mu\text{mol L}^{-1}$ range would be numerically negligible compared with deviations in the mmol L^{-1} range. Row wise normalisation therefore ensures that proportional deviations are penalised on a comparable scale, preventing magnitude driven bias and enabling balanced optimisation for both macro and micronutrients.

In this sense, SSE and SSRE should not be interpreted as direct optimisation targets, but as complementary descriptors of the achieved solution. SSE is particularly informative for identifying chemically relevant leakage into nutrients that were intended to remain absent, whereas SSRE is more informative for evaluating proportional agreement with specified non zero targets.

A useful illustration is the 1:1 salt coupling case, for example NaCl or NH_4NO_3 . If the targets are 1 mmol L^{-1} for one ion and 2 mmol L^{-1} for the coupled counter ion, relative error minimisation

yields an optimum at 1.2 mmol L^{-1} rather than the arithmetic midpoint of 1.5 mmol L^{-1} . This demonstrates a non intuitive but important property of the normalised objective: the optimum is shifted toward the smaller target because deviations are scaled relative to target magnitude.

5.4 Weighting and two stage optimisation as practical control tools

Weighting is optional, so the solver can also be used in an unweighted form, in which all nutrients contribute equally after normalisation. If weights are applied, the problem becomes a weighted non negative least squares problem, in which deviations in highly weighted nutrients contribute more strongly to the objective function than deviations in nutrients with lower weights. Lowering the weight of one nutrient shifts more of the residual error to that nutrient and allows the remaining nutrients to be matched more closely. In practical terms, this means that one nutrient can be deliberately sacrificed to improve the overall fit of the others when the target profile is difficult or impossible to achieve with the selected salts.

This feature is particularly useful for chemically constrained formulations. If one nutrient cannot be matched simultaneously with all others because of stoichiometric coupling within the available fertilizers, reducing its weight may substantially improve the fit for the remaining targets. When the second solver stage is enabled, additional adjustment with acids or bases may further improve the final composition and partially compensate for such trade offs.

It was possible to achieve an exact match between target and achieved nutrient concentrations for most of the crop set of Sonneveld & Straver (1994) using the one stage salts only solver. This demonstrates that the implemented salt matrix is sufficiently comprehensive to represent many standard horticultural target profiles.

The two stage solver nevertheless provides an important extension when an exact solution is not achievable with neutral salts alone. In this architecture, the first stage fits the composition using neutral fertilizer salts, whereas acids and bases are introduced only in a second corrective step. This separation is chemically meaningful because it prevents acids and bases from acting as unrestricted first stage degrees of freedom that could improve the numerical fit while reducing interpretability of the resulting fertilizer composition.

The results support this design particularly for pH active or stoichiometrically awkward cases, such as silicate containing formulations. The two stage framework should therefore not be understood merely as a numerical convenience, but as a practically motivated control mechanism that balances solvability, chemical interpretability and operational realism.

5.5 Electroneutrality and pH as coupled constraints

Electroneutrality is one of the most fundamental principles in nutrient solution formulation. In any aqueous electrolyte solution, the sum of positive charges must equal the sum of negative charges.

Thus, the total charge contributed by cations is balanced by the total charge contributed by anions (Kalka, 2024). While many dissolved compounds dissociate into well defined cations and anions and therefore contribute predictably to charge balance, some nutrients exist in different ionic species depending on pH. At the same time, pH should not be understood merely as a measure of hydrogen ion concentration, but more accurately as a measure of hydrogen ion activity. From this perspective, pH can be interpreted as a dependent variable constrained by the electroneutrality condition in aqueous solution (Stumm & Morgan, 1996). Accordingly, the activities of H_3O^+ and OH^- adjust in conjunction with pH dependent speciation systems such as phosphate, sulfate, carbonate, borate and silicate to maintain charge balance, rather than being determined solely by the stoichiometric addition and neutralisation of acid and base.

Electroneutrality ultimately determines whether a target nutrient composition is physically realisable. In the optimisation procedure, non negativity of salt addition was implemented explicitly as a mathematical constraint, whereas electroneutrality was not imposed as an additional constraint. This is because fertilizer salts contribute ions in stoichiometrically balanced combinations. Acid or base addition changes pH and ionic speciation without violating the overall charge balance of the solution. Consequently, if a target nutrient profile is itself not electroneutral, no exact solution exists and the solver can only minimise the deviation from the specified targets.

This is directly relevant for interpretation of pH within the framework. pH should not be treated as an independently selectable scalar, but as an emergent property of charge balance, acid base equilibria and activities. Integrating charge balance based pH estimation therefore extends formulation from simple nutrient matching toward chemically coherent solution evaluation.

5.6 Role of pH and EC modelling in the workflow

A notable strength of the framework is that formulation is not treated as complete once nutrient targets have been matched. Estimated pH and EC provide a second interpretive layer linking nutrient composition to broader physicochemical behaviour.

This is relevant because two solutions with similar nominal nutrient targets can differ in buffering, ionic strength, conductivity or pH response depending on fertilizer selection, water composition and corrective additions.

The pH of a nutrient solution is a critical parameter that determines the speciation of elements across their various physicochemical forms (free ions, complexes, chelates, ion pairs, solid and gaseous phases and different oxidation levels) and is therefore critical for nutrient bioavailability (Rijck & Schrevens (1998)).

The implemented pH estimation module provides a practical way to evaluate the expected pH of the formulated solution based on its composition and the applied acid or base correction. This allows users to anticipate whether the resulting pH will be within a desirable range for plant growth and to make informed decisions about formulation adjustments if necessary.

Within the tested ranges, the implemented pH and EC models provided useful chemically informed approximations. The Davies activity correction showed better pH agreement than the limiting Debye Hückel model in phosphate buffer comparisons. Likewise, EC estimation based on diffusion coefficients together with Davies activity correction agreed closely with reference KCl values and with PHREEQC based comparisons, whereas simpler empirical approaches tended to underestimate conductivity at higher concentrations.

These modules do not replace full geochemical simulation. Their value lies in practical workflow integration because users can evaluate not only whether a target vector can be approximated, but also what pH and EC consequences follow from the achieved formulation.

5.7 Relevance of the mass balance approach

“Judging from their publications and statements, it would seem that many investigators have grown crops in a solution chosen largely at random, and that, where the results have been successful, they have recommended that solution as being specific for the particular crop concerned.” (Steiner, 1961)

Optimal nutrient solutions can be formulated using the principle of mass balance (Langenfeld et al., 2022). In contrast to empirically derived nutrient recipes, this approach seeks to relate nutrient supply directly to plant demand. Plant nutrient demand can be approximated from tissue nutrient composition in combination with plant water consumption, thereby linking absolute nutrient uptake to water uptake. Langenfeld et al. (2022) distinguish between active uptake (N, P, K, Mn), intermediate uptake (Mg, S, Fe, Cu, Zn, Mo, Ni, Cl) and passive uptake (Ca, B). Passively taken up nutrients such as Ca are closely related to transpired water and are therefore well suited to a mass balance approach. Since tissue analysis represents a destructive end point measurement, non linearities in active nutrient uptake kinetics, including transitions between high affinity and low affinity transport systems, are of limited relevance in this context. With both WUE and tissue analysis, the exact temporal dynamics of uptake do not need to be resolved explicitly, as the temporal dimension is subsumed into the final nutrient solution composition. Thus, although active and passive uptake differ mechanistically and kinetically, these differences converge in the relationship between cumulative water use and the plant tissue nutrient content observed at harvest.

On this basis, crop specific target concentrations for the nutrient solution can be derived. Under ideal conditions, such a formulation would maintain the residual or recirculating solution in approximate balance over time, thereby minimising the accumulation or depletion of individual nutrients.

The present framework is particularly relevant here because a biologically plausible demand vector is not automatically a chemically realisable nutrient solution. Even if tissue composition and water use define a meaningful elemental target, this still has to be translated into an ionic target and then into a feasible fertilizer composition. NutriCalc provides a transparent way to make this translation explicit.

A remaining limitation of tissue based mass balance approaches is the translation of total plant N into a chemically feasible nutrient solution target. Specifically, total N must be partitioned into nitrate and ammonium. To address this, the present work introduces a refinement of the mass balance approach in which total N is partitioned into $\text{NO}_3\text{-N}$ and $\text{NH}_4\text{-N}$ by imposing electroneutrality as an explicit formulation constraint. This step enables the derivation of nutrient solutions that are chemically plausible and electrochemically balanced. In the example considered here, based on Langenfeld et al. (2022), a total N concentration of 90 mg L^{-1} yields a partitioning of 82 mg L^{-1} $\text{NO}_3\text{-N}$ and 8 mg L^{-1} $\text{NH}_4\text{-N}$.

This does not yet constitute a full physiological model of nutrient uptake, but it is a meaningful methodological step. It bridges biological demand estimation and chemically constrained fertilizer realisation within one transparent computational framework.

5.8 Comparison with existing formulation approaches

Existing nutrient formulation practice spans fixed recipes, manual iterative balancing, spreadsheet tools and dedicated software. Classical approaches such as the universal algorithm of Sonneveld et al. (1999) are chemically informed and practically valuable, but they are not necessarily expressed as explicit constrained optimisation frameworks. Spreadsheet based calculators can be useful in practice, but often remain difficult to extend reproducibly and may obscure the underlying calculation logic. Other software such as HydroBuddy offers flexible nutrient formulation functionality, but NutriCalc contributes a distinct combination of explicit stoichiometric representation, weighted row normalised NNLS fitting, charge balance oriented evaluation and modular open source implementation in R.

The main contribution of NutriCalc is therefore not merely that it calculates fertilizer quantities, but that it formalises nutrient formulation as a transparent, reproducible and modifiable constrained optimisation problem. The R implementation supports programmable workflows and integration with other analyses, whereas the Shiny interface broadens accessibility for users without programming experience.

5.9 Limitations and scope

Despite its strengths, the framework has clear limitations. First, precipitation reactions are not modeled beyond BAK separation. A target may therefore appear feasible in dissolved ion terms while being unstable in concentrated stock solutions.

Second, chemical speciation remains simplified. Major acid base systems and fixed charge chelates are represented, but a full complexation model is not implemented yet.

Third, the activity corrections used for pH and EC estimation remain approximate. Debye Hückel and Davies models are suitable for dilute to moderately concentrated systems, but they do not capture all non ideal effects at higher ionic strengths.

Fourth, non zero residuals may reflect limitations of the selected fertilizer matrix rather than universal chemical impossibility. Exact solvability is always conditional on the implemented fertilizer set, which is not yet complete.

Fifth, the optimisation currently targets nutrient fit rather than explicit minimisation of cost, number of salts or operational complexity. Although users can partly influence the returned solution through salt selection and weighting, these objectives are not yet formalised directly.

Finally, the framework is a formulation and evaluation model rather than a crop growth model. It can represent and assess nutrient solution chemistry, but it does not predict plant response, uptake kinetics or recirculating system dynamics over time.

These limitations define the scope of the framework rather than invalidating its usefulness. NutriCalc should therefore be understood as a transparent and extensible formulation tool that captures major stoichiometric and physicochemical constraints, while leaving more advanced geochemical and biological dynamics to future development.

6 Conclusions and Outlook

6.1 Scientific contribution

This thesis developed *NutriCalc*, an open-source computational R package with a Shiny interface for the formulation, analysis and optimisation of nutrient solutions in horticulture. The work formalised nutrient-solution design as a constrained linear optimisation problem and implemented this formulation using non-negative least squares. In doing so, it linked stoichiometric conversion, target-based optimisation, water-source correction and chemically informed evaluation of pH and electrical conductivity within one transparent workflow.

The results showed that a broad range of published horticultural target solutions can be reproduced exactly or with only minor residual deviations. Where deviations remained, these were generally chemically interpretable and could often be traced to fertilizer representation, electroneutrality constraints or chemically coupled sources such as potassium silicate. Beyond reproducing established recipes, the framework also demonstrated how nutrient formulation can be extended toward mass-balance-derived targets and charge-balance-based interpretation.

Taken together, the thesis shows that nutrient-solution formulation can be moved from recipe-based and partly opaque practice toward a reproducible, optimisation-based and chemically explicit computational workflow.

6.2 Practical relevance for horticulture

The practical relevance of the framework lies in its applicability to both research and horticultural practice. For researchers, *NutriCalc* provides a transparent tool for constructing and comparing nutrient treatments under explicitly defined chemical constraints. For practitioners, it offers a flexible way to formulate crop-specific nutrient solutions while accounting for local water quality, available fertilizers and physicochemical properties of the final solution.

By combining an R package with a Shiny-based graphical user interface, the framework is accessible both for programmable analytical workflows and for users without coding experience. In this sense, *NutriCalc* contributes not only methodologically but also operationally to digital nutrient management in horticulture.

6.3 Limitations

The scope of the present thesis is limited in several respects. Precipitation reactions and full metal-ligand complexation were not modeled explicitly, so the framework should not be interpreted as a complete geochemical simulation environment. Likewise, the pH and EC calculations remain approximations based on selected equilibrium and activity models. In addition, exact solvability always depends on the implemented fertilizer matrix, which is necessarily incomplete.

The framework is therefore best understood as a transparent formulation and evaluation tool that captures major stoichiometric and physicochemical constraints, but not the full complexity of nutrient chemistry or crop response.

6.4 Future extensions

Future development of NutriCalc could expand both the scientific scope of the framework and its practical applicability in horticultural research and nutrient-solution management. One important direction is the expansion of the fertilizer database. This includes adding further compounds commonly used in horticultural practice, enabling users to define custom salts directly within the application and extending the system with additional micronutrients such as Ni and Co together with appropriate fertilizer sources. Such developments would increase flexibility and allow the software to be adapted to different regional fertilizers and cultivation systems.

A second direction concerns the chemical model. In addition to the current treatment of nutrient composition, pH and EC, future versions could incorporate more detailed chemical speciation and explicit precipitation modelling. This would improve applicability particularly for the formulation of concentrated stock solutions and would further strengthen the chemical realism of the framework.

A third direction concerns the optimisation framework itself. At present, formulation is based on an NNLS approach that focuses on matching target nutrient concentrations as closely as possible. Future versions could include additional constraints such as fertilizer price, thereby allowing the formulation process to reflect not only chemical targets but also economic considerations.

A fourth direction involves the representation of nutrient targets and the expansion of built-in datasets. Because the internal formulation matrix operates on a stoichiometric basis, conversion between molar and mass-based units is straightforward. NutriCalc could therefore be extended to accept target nutrient profiles not only in mmol L^{-1} but also in mass-based units or nutrient percentages, which would improve usability for users more familiar with commercial fertilizer labels and analytical results reported in such forms. Likewise, the dataset of target nutrient profiles could be expanded to include a wider range of crops, production systems and developmental stages. For example, stage-specific target profiles could be provided for crops such as tomato, in which nutrient demand changes substantially between vegetative growth, flowering and fruit development. A simple external input format such as CSV could facilitate maintenance and extension of such datasets and would also allow users to adapt recipes to their own conditions.

A fifth direction is the further development of reporting and user interaction. From a practical perspective, a particularly useful next step would be the integration of export functions into the Shiny application. Because this thesis was developed in Quarto, PDF and CSV exports could be implemented to generate structured reports for individual nutrient formulations. These reports could include the selected fertilizer salts, achieved nutrient concentrations, deviations from target values and predicted pH and EC together with supporting tables and visualisations. This would strengthen the value of NutriCalc as both a scientific and an applied tool.

Overall, future extensions of NutriCalc could strengthen the package in three broad areas: broader input flexibility, more advanced chemical and optimisation routines and improved reporting and user interaction. The most important future extension, however, is the integration of experimental data, particularly for the mass-balance approach, in order to derive more precisely balanced and crop-specific nutrient recipes for closed hydroponic systems.

7 Summary (possibly in German and English)

8 References

- Amelung, W., Blume, H.-P., Fleige, H., Horn, R., Kandeler, E., Kögel-Knabner, I., Kretzschmar, R., Stahr, K., Wilke, B.-M., Scheffer, F., & Schachtschabel, P. (2018). *Scheffer/schachtschabel lehrbuch der bodenkunde* (17., überarbeitete und ergänzte Auflage). Berlin; Heidelberg: Springer Spektrum. <https://doi.org/10.1007/978-3-662-55871-3>
- Appelo, C. A. J. (2005). *Geochemistry, groundwater and pollution* (Second edition). Leiden: A.A. Balkema Publishers. <https://doi.org/10.1201/9781439833544>
- Appelo, C. A. J. (2010). *SPECIFIC CONDUCTANCE: How to calculate, to use, and the pitfalls*. Retrieved from <https://www.hydrochemistry.eu/exmpls/sc.html>
- BIPM. (2019). *The international system of units (SI)*. Bureau International des Poids et Mesures. <https://doi.org/10.59161/AUEZ1291>
- Bower, V. E., & Bates, R. G. (1955). pH values of the clark and lubs buffer solutions at 25 c. *Journal of Research of the National Bureau of Standards*, 55, 197. Retrieved from <https://api.semanticscholar.org/CorpusID:101259939>
- BWB. (2026). *Analysedaten nach postleitzahlen: Bitte geben sie ihre postleitzahl in das suchfenster ein, um die werte für ihren wohnbezirk zu erhalten*. Retrieved from <https://www.bwb.de/de/analysedaten-nach-postleitzahlen.php?PLZ=x>
- Chang, W., Cheng, J., Allaire, J. J., Sievert, C., Schloerke, B., Xie, Y., Allen, J., McPherson, J., Dipert, A., & Borges, B. (2025). *Shiny: Web application framework for r*. Comprehensive R Archive Network (CRAN). <https://doi.org/10.32614/CRAN.package.shiny>
- Fernández Pinto, D. (2022). *HydroBuddy: An open source nutrient calculator for hydroponics and general agriculture*. Retrieved from <https://github.com/danielfppps/hydrobuddy>
- George, E., Horst, W. J., & Neumann, E. (2012). Adaptation of plants to adverse chemical soil conditions. In *Marschner's mineral nutrition of higher plants* (pp. 409–472). Elsevier. <https://doi.org/10.1016/b978-0-12-384905-2.00017-0>
- Haynes, W. M. (Ed.). (2017). *CRC handbook of chemistry and physics* (97th edition). Boca Raton; London; New York: CRC Press.
- Hewitt, E. J. (1966). *Sand and water culture methods used in the study of plant nutrition* (2. rev. edition). Farnham Royal: Commonwealth Agricultural Bureaux.
- Hoagland, D. R. (1920). Optimum nutrient solutions for plants. *American Association for the Advancement of Science*. Retrieved from <https://www.science.org/doi/10.1126/science.52.1354.562>
- Hoagland, D. R., & Arnon, D. I. (1938). *The water-culture method for growing plants without soil*.
- Incrocci, L. (2007). *Nutrient solution calculator: Retrieved from the internet archive (wayback machine)* (Dipartimento di Biologia delle Piante Agrarie, University of Pisa, Ed.). Retrieved from <https://web.archive.org/web/20240813175407/https://www.wur.nl/en/research-results/projects-and-programmes/euphoros/calculation-tools/nutrient-solution-calculator.htm>

- Incrocci, L., M., Giménez Moolhuyzen, M., & Guzmán Palomino, J. M. (2020). *Calculadora de soluciones nutritivas (SN)*. Retrieved from <https://repositorio.ual.es/handle/10835/7670?show=full>
- Jones, J. B. (2014). *Complete guide for growing plants hydroponically*. Boca Raton: CRC Press.
- Jones, J. J., Shaw, C., Chen, T., Staß, C. M., Ulrichs, C., Riewe, D., Kloas, W., & Geilfus, C. (2024). Plant nutritional value of aquaculture water produced by feeding Nile tilapia (*Oreochromis niloticus*) alternative protein diets: A lettuce and basil case study. *Plants, People, Planet*, 6(2), 362–380. <https://doi.org/10.1002/ppp3.10457>
- Kalka, H. (2020). *Electrical conductivity based on diffusion coefficients*. Retrieved from <https://www.aqion.de/site/77>
- Kalka, H. (2023). *Aktivitätskoeffizient und Aktivitätsmodelle*. Retrieved from <https://www.aqion.de/site/41>
- Kalka, H. (2024). *Charge-balance error (CBE)*. Retrieved from <https://www.aqion.de/site/92>
- Knop, W. (1860). Ueber die Ernährung der Pflanzen durch wässrige Lösungen bei Ausschluss des Bodens. *Die Landwirtschaftlichen Versuchs-Stationen*, (2).
- Langenfeld, N. J., Pinto, D. F., Faust, J. E., Heins, R., & Bugbee, B. (2022). Principles of nutrient and water management for indoor agriculture. *Sustainability*, 14(16), 10204. <https://doi.org/10.3390/su141610204>
- Lawson, C. L., & Hanson, R. J. (1974). *Solving least squares problems*. Englewood Cliffs, NJ: Prentice-Hall.
- Lawson, C. L., & Hanson, R. J. (1995). *Solving least squares problems* ([10. Dr.]). Philadelphia: SIAM.
- McCleskey, R. B., Nordstrom, D. K., & Ryan, J. N. (2012). Comparison of electrical conductivity calculation methods for natural waters. *Limnology and Oceanography: Methods*, 10(11), 952–967. <https://doi.org/10.4319/lom.2012.10.952>
- Mullen, K. M., & van Stokkum, I. H. M. (2024). *The lawson-hanson algorithm for non-negative least squares (NNLS): An R interface to the lawson-hanson implementation of an algorithm for non-negative least squares (NNLS). Also allows the combination of non-negative and non-positive constraints*. Comprehensive R Archive Network (CRAN). <https://doi.org/10.32614/CRAN.package.nnls>
- Pardossi, Carmassi G., Diara C., I. L., Maggini R., & Massa D. (2011). *Fertigation and substrate management in closed soilless culture*. Dipartimento di Biologia delle Piante Agrarie, University of Pisa.
- Parkhurst, D. L., & Appelo, C. A. J. (2013). *Description of input and examples for PHREEQC version 3: A computer program for speciation, batch-reaction, one-dimensional transport, and inverse geochemical calculations*. Reston, Virginia: U.S. Department of the Interior U.S. Geological Survey. Retrieved from <http://purl.fdlp.gov/GPO/gpo37078>
- Posit team. (2025). *RStudio: Integrated development environment for R*. Posit Software, PBC. Retrieved from <http://www.posit.co/>
- Pratt, K. W., Koch, W. F., Wu, Y. C., & Berezansky, P. A. (2001). Molality-based primary standards of electrolytic conductivity (IUPAC technical report). *Pure and Applied Chemistry*, 73(11), 1783–1793. <https://doi.org/10.1351/pac200173111783>
- R Core Team. (2025). *R: A language and environment for statistical computing*.

- Rengel, Z., Cakmak, I., White, P. J., Marschner, H., Prof. Dr. agr., Dres. h.c, Landwirtschaft, Agrarwissenschaftler, Zuckmantel, Stuttgart, Hohenheim, Marschner, P., 1929-1996, 30.10.1929-21.09.1996, & University of Western Australia School of Agriculture. (2023). *Marschner's mineral nutrition of plants* (4th ed.). London; San Diego, CA: Academic Press, an imprint of Elsevier.
- Resh, H. M. (2013). *Hydroponic food production: A definitive guidebook for the advanced home gardener and the commercial hydroponic grower* (7th ed.). Boca Raton [u.a.]: CRC Press.
- Rijk, G. de, & Schrevens, E. (1998). Comparison of the mineral composition of twelve standard nutrient solutions. *Journal of Plant Nutrition*, 21(10), 2115–2125. <https://doi.org/10.1080/01904169809365548>
- Shive, J. W. (1915). A three-salt nutrient solution for plants. *American Journal of Botany*, 2(4), 157. <https://doi.org/10.2307/2435048>
- Sievert, C., Cheng, J., Aden-Buie, G., Aguilar, J., & Park, T. (2025). *Bslib: Custom 'bootstrap' 'sass' themes for 'shiny' and 'rmarkdown'*. Comprehensive R Archive Network (CRAN). <https://doi.org/10.32614/CRAN.package.bslib>
- Smith, M. R., & Sanselme, L. (2025). *Ternary: Create ternary and holdridge plots*. Comprehensive R Archive Network (CRAN). <https://doi.org/10.32614/CRAN.package.Ternary>
- Sonneveld, C. (2009). *Plant nutrition of greenhouse crops* (1st ed.). Dordrecht: Springer.
- Sonneveld, C., & Straver, N. (1994). *Nutrient solutions for vegetables and flowers grown in water or substrates*. Retrieved from <https://edepot.wur.nl/237302>
- Sonneveld, C., Voogt, W., & Spaans, L. (1999). A universal algorithm for calculation of nutrient solutions. *Acta Horticulturae*, (481), 331–340. <https://doi.org/10.17660/actahortic.1999.481.38>
- Steiner, A. A. (1961). A universal method for preparing nutrient solutions of a certain desired composition. *Plant and Soil*, 15(2), 134–154. <https://doi.org/10.1007/BF01347224>
- Stumm, W., & Morgan, J. J. (1996). *Aquatic chemistry: Chemical equilibria and rates in natural waters* (3. ed.). New York, NY: Wiley.
- United States Salinity Laboratory Staff. (1954). *Diagnosis and improvement of saline and alkali soils*. Washington, DC: United States Department of Agriculture.
- van der Lugt, G., Holwerda, H. T., Hora, K., Bugter, M., Hardeman, J., & Vries, P. de. (2020). *Nutrient solutions for greenhouse crops: Made available by: Eurofins agro, geerten, van der lugt, nouryon, SQM, yara* (4th ed.).
- Voogt, W., & Sonneveld, C. (2001). Silicon in horticultural crops grown in soilless culture. In L. E. Datnoff, G. H. Snyder, & G. H. Korndörfer (Eds.), *Silicon in agriculture* (pp. 115–131). Elsevier. [https://doi.org/10.1016/S0928-3420\(01\)80010-0](https://doi.org/10.1016/S0928-3420(01)80010-0)
- Wickham, H. (2025). *Stringr: Simple, consistent wrappers for common string operations*. Comprehensive R Archive Network (CRAN). <https://doi.org/10.32614/CRAN.package.stringr>
- Wickham, H., Hester, J., Chang, W., & Bryan, J. (2025). *Devtools: Tools to make developing r packages easier*. Comprehensive R Archive Network (CRAN). <https://doi.org/10.32614/CRAN.package.devtools>
- Wickham, H., Hester, J., & Ooms, J. (2026). *xml2: Parse XML*. Comprehensive R Archive Network (CRAN). <https://doi.org/10.32614/CRAN.package.xml2>
- YARA. (2026). *YaraTera KRISTA MgS*. Retrieved from <https://www.yara.co.uk/crop-nutrition/>

fertiliser/soluble/yaratera-krista-mgs/

- [1] ".19.05.1995"
- [3] "Riedel.2015"
- [5] "Rakshit.2015"
- [7] "Naeem.2017"
- [9] "Ryan.2014"
- [11] "Sheykhani.2024"
- [13] "Stout.1939"
- [15] "Datnoff.2001"
- [17] "Beran.2014"
- [19] "Asao.2012"
- [21] ".2025"
- [23] "Einstein.1905"
- [25] "Jenne.1979"
- [27] "Gruda.2021"
- "Rijck.1998b"
- "Resh.2015"
- "Pratt.2001"
- "Mohmed.2025"
- "Savvas.1999"
- "Turan.2022"
- "ScienceDirectOnlineservice.2001"
- "Bugbee.2024"
- "Benavente.2025"
- "AliAlMeselmani.2022"
- ".2012"
- "Evangelou.1998"
- "I..2012"

[1] 27

Appendix

Declaration of Independence