

# On the minimal number of generators of a finite group

Diogo Santos

November 1, 2024

- Finding the minimal number of generators of a finite group  $H$

- Finding the minimal number of generators of a finite group  $H$

Can be reduced to:

- Finding the minimal number of generators of a finite group  $H$  such that  $d(H/N) \leq m$  for every non-trivial normal subgroup  $N$ , but  $d(H) > m$

# The case $m = 1$

## Theorem

*Let  $H$  be a finite nilpotent group such that  $d(H/N) \leq 1$  for every non-trivial normal subgroup  $N$ , but  $d(H) > 1$ . Then  $H \cong \mathbb{Z}_p \times \mathbb{Z}_p$  for some prime  $p$ .*

# The case $m = 1$

## Theorem

*Let  $H$  be a finite nilpotent group such that  $d(H/N) \leq 1$  for every non-trivial normal subgroup  $N$ , but  $d(H) > 1$ . Then  $H \cong \mathbb{Z}_p \times \mathbb{Z}_p$  for some prime  $p$ .*

## Proof.

- $H = P_1 \times \dots \times P_n$  where  $P_i$  is a Sylow  $p_i$ -subgroup for  $1 \leq i \leq n$  and  $p_1, \dots, p_n$  are distinct primes.

# The case $m = 1$

## Theorem

*Let  $H$  be a finite nilpotent group such that  $d(H/N) \leq 1$  for every non-trivial normal subgroup  $N$ , but  $d(H) > 1$ . Then  $H \cong \mathbb{Z}_p \times \mathbb{Z}_p$  for some prime  $p$ .*

## Proof.

- $H = P_1 \times \dots \times P_n$  where  $P_i$  is a Sylow  $p_i$ -subgroup for  $1 \leq i \leq n$  and  $p_1, \dots, p_n$  are distinct primes.
- If  $P_1, \dots, P_n$  are cyclic, we obtain  $H \cong \mathbb{Z}_{p_1 \dots p_n}$  which contradicts  $d(H) > 1$ . Without loss of generality we can thus assume that  $P_1$  is not cyclic.

# The case $m = 1$

## Theorem

*Let  $H$  be a finite nilpotent group such that  $d(H/N) \leq 1$  for every non-trivial normal subgroup  $N$ , but  $d(H) > 1$ . Then  $H \cong \mathbb{Z}_p \times \mathbb{Z}_p$  for some prime  $p$ .*

## Proof.

- $H = P_1 \times \dots \times P_n$  where  $P_i$  is a Sylow  $p_i$ -subgroup for  $1 \leq i \leq n$  and  $p_1, \dots, p_n$  are distinct primes.
- If  $P_1, \dots, P_n$  are cyclic, we obtain  $H \cong \mathbb{Z}_{p_1 \dots p_n}$  which contradicts  $d(H) > 1$ . Without loss of generality we can thus assume that  $P_1$  is not cyclic.
- $n \geq 2 \implies P_1 \cong H/(1 \times P_2 \dots \times P_n)$  and thus  $d(P_1) = d(H/(1 \times P_2 \dots \times P_n)) = 1$ , contradiction.

# The case $m = 1$

## Theorem

*Let  $H$  be a finite nilpotent group such that  $d(H/N) \leq 1$  for every non-trivial normal subgroup  $N$ , but  $d(H) > 1$ . Then  $H \cong \mathbb{Z}_p \times \mathbb{Z}_p$  for some prime  $p$ .*

## Proof.

- $H = P_1 \times \dots \times P_n$  where  $P_i$  is a Sylow  $p_i$ -subgroup for  $1 \leq i \leq n$  and  $p_1, \dots, p_n$  are distinct primes.
- If  $P_1, \dots, P_n$  are cyclic, we obtain  $H \cong \mathbb{Z}_{p_1 \dots p_n}$  which contradicts  $d(H) > 1$ . Without loss of generality we can thus assume that  $P_1$  is not cyclic.
- $n \geq 2 \implies P_1 \cong H/(1 \times P_2 \dots \times P_n)$  and thus  $d(P_1) = d(H/(1 \times P_2 \dots \times P_n)) = 1$ , contradiction.
- By Theorem ??,  $\Phi(H) = 1$  hence  $H = (\mathbb{Z}_{p_1})^q$  by Theorem ??.



# The case $m = 1$

Proof.

- $q = 2$  since

$$q - 1 = d((\mathbb{Z}_{p_1})^{q-1}) = d(H/(\mathbb{Z}_{p_1} \times 1 \times \dots \times 1)) = 1.$$

