NUMERICAL EVIDENCE FOR CONJECTURES ON NEGATIVE DISCRETE SECOND MOMENTS OF DIRICHLET L-FUNCTIONS

DIONATAN RESENDE

ABSTRACT. Pearce-Crump recently conjectured that, for a primitive Dirichlet character χ modulo q, $\sum_{0<\gamma\le T}|L'(\rho,\chi)|^{-2}\sim\frac{A_q}{2\pi}T$ with $A_q=\frac{6}{\pi^2}\prod_{p\mid q}\frac{p}{p+1}$, and $\sum_{0<\gamma\le T}|L(2\rho,\chi^2)/L'(\rho,\chi)|^2\sim\frac{\varphi(q)}{q}\frac{T}{2\pi}$. These conjectures come from a long-mollifier heuristic and generalise conjectures of Gonek and Ng for the Riemann zeta-function. We test them numerically for the real primitive characters of conductor 4, 5, 8 and 12, using between 2000 and 3000 zeros of each L-function, computed and verified for completeness, together with the first 27000 zeros of $\zeta(s)$, comparing each L-function with $\zeta(s)$ at the same local density of zeros. In all eight tests the predicted arithmetic factor lies inside the 95% bootstrap interval and the value without arithmetic factor is excluded. For conductor 12, where the two conjectures predict $1/2$ and $1/3$, we measure $0.501\,[0.445,0.555]$ and $0.340\,[0.301,0.381]$. All predictions and analysis choices were fixed before the corresponding computations. The arithmetic diagonals of regularised shifted means have residues matching the two conjectured constants. A uniform passage from these means to the discrete moments remains unresolved; we do not claim an error term or the absence of lower-order terms. Rare close pairs of zeros limit the interpretation of the finite-height numerical comparisons.

1. INTRODUCTION

Let $\rho=\frac{1}{2}+i\gamma$ denote the non-trivial zeros of $\zeta(s)$. Assuming the Riemann Hypothesis and that all zeros are simple, Gonek [4] conjectured

$$\sum_{0<\gamma\le T}\frac{1}{|\zeta'(\rho)|^2}\sim\frac{3}{\pi^3}\,T, \tag{1}$$

and Milinovich and Ng [7] proved a lower bound of half this size. Ng [8] conjectured the companion asymptotic $\sum_{0<\gamma\le T}|\zeta(2\rho)/\zeta'(\rho)|^2\sim T/2\pi$, with a matching lower bound of half the size due to Sinha [10]. Negative discrete moments of this kind enter explicit formulae for the summatory functions of the Möbius and Liouville functions, and their sizes were modelled with random matrix theory by Hughes, Keating and O'Connell [6]. Numerically, Hiary and Odlyzko [5] found agreement with (1) using large sets of zeros near the 10^{16}th, 10^{20}th and 10^{23}rd zeros.

For a primitive Dirichlet character χ modulo $q>1$, Pearce-Crump [9] proved lower bounds for

$$J_\chi(T)=\sum_{0<\gamma\le T}\frac{1}{|L'(\rho,\chi)|^2},\qquad K_\chi(T)=\sum_{0<\gamma\le T}\left|\frac{L(2\rho,\chi^2)}{L'(\rho,\chi)}\right|^2,$$

uniformly in the conductor, and formulated the following conjectures [9, Conjectures 7 and 8]:

$$J_\chi(T)\sim\frac{A_q}{2\pi}\,T,\qquad A_q=\frac{6}{\pi^2}\prod_{p\mid q}\frac{p}{p+1}, \tag{2}$$

$$K_\chi(T)\sim\frac{\varphi(q)}{q}\,\frac{T}{2\pi}. \tag{3}$$

Both reduce to the conjectures of Gonek and Ng when $q=1$. They were obtained from Farmer's long-mollifier heuristic [3], which, as Pearce-Crump points out, is not a proof and is applied there to

Date: Internal draft v3, 13 October 2026.

discrete sums rather than to the continuous mean values for which it was designed. Their derivation from the ratios conjecture [1, 2] is left open in [9].

The content of (2) and (3) beyond the case $q = 1$ lies in the arithmetic factors $\prod_{p|q} p/(p+1)$ and $\varphi(q)/q$. The aim of this note is to test these factors numerically. We are not aware of previous numerical tests of (2) or (3); our literature search is described in Section 8. We record a heuristic consistency check from the random matrix model of Hughes, Keating and O'Connell (Section 5) and the corresponding arithmetic diagonal residues (Section 6). The latter calculation does not establish a uniform ratios prediction for the discrete moments.

2. Method

2.1. **Normalisation.** Since the zeros of $L(s,\chi)$ at height t have density $\frac{1}{2\pi}\log\frac{qt}{2\pi}$, (2) and (3) say, to first order, that the zero-wise averages of

$$v_n = \frac{\log(q\gamma_n/2\pi)}{|L'(\rho_n,\chi)|^2}, \qquad w_n = \frac{\log(q\gamma_n/2\pi)\,|L(2\rho_n,\chi^2)|^2}{|L'(\rho_n,\chi)|^2}$$

tend to A_q and $\varphi(q)/q$ respectively (to $6/\pi^2$ and 1 for ζ).

2.2. **A differential statistic.** At the heights accessible to us, sums over a few thousand zeros fluctuate by several per cent around their expected values. For $\zeta(s)$, the ratio of $\sum 1/|\zeta'(\rho)|^2$ over consecutive windows of 3000 zeros to $\frac{3}{\pi^3}\Delta T$ ranges from 0.907 to 1.088 in the first 27000 zeros, with no trend in the height; most windows fall below 1. These descriptive values alone do not distinguish lower-order effects from rare-pair fluctuations. Rather than compare absolute averages with the conjectured constants, we compare each L-function with $\zeta(s)$ at the same local density $D = \log(q\gamma/2\pi)$. We split the zeros of $L(s,\chi)$ into ten bins at the deciles of D and define

$$R = \frac{\operatorname{mean}_\chi(v)}{\sum_b \frac{N_b}{N}\,\operatorname{mean}_{\zeta,\,b}(v)},$$

where N_b is the number of zeros of $L(s,\chi)$ in bin b and $\operatorname{mean}_{\zeta,b}(v)$ is the average of v over the zeros of ζ whose density $\log(\gamma/2\pi)$ lies in the same bin; R_8 is defined in the same way with w. Conjectures (2) and (3) predict

$$R = \prod_{p|q}\frac{p}{p+1}, \qquad R_8 = \frac{\varphi(q)}{q}.$$

We designed this statistic in the expectation that density matching would reduce common finite-height effects. This remains a modelling assumption; lower-order corrections have not been derived, and density matching does not guarantee cancellation of rare-pair effects. The shared ζ reference adds sampling fluctuations. The block bootstrap measures variability in the observed samples and may underestimate uncertainty from unobserved rare close pairs. The absolute ratios are reported in Section 4 for comparison.

2.3. **Alternatives.** For each test we also record two alternatives: H_0, no arithmetic factor ($R = 1$); and, for (2), the factor $\prod_{p|q} p^2/(p^2-1)$ obtained by simply removing the Euler factors at $p \mid q$ from $1/\zeta(2)$; for (3), the factor of (2), which tests whether the two moments require different factors.

2.4. **Uncertainty.** The terms $1/|L'(\rho,\chi)|^2$ have a heavy tail produced by close pairs of zeros: the largest 1% of the terms carry 12–23% of the sums. We use 95% percentile intervals from a block bootstrap (blocks of 100 consecutive zeros, resampled independently for $L(s,\chi)$ and $\zeta(s)$, 2000 replicates). Because the bootstrap may under-cover with heavy tails, we also recompute R after removing the five largest terms of $L(s,\chi)$.

2.5. **Pre-registration.** For each group of tests, the predictions, alternatives, statistics, bootstrap settings and decision rules were written down and fixed (with SHA-256 hashes) before the derivatives were computed. A test is counted as favourable if the predicted value lies in the interval and H_0 does not.

3. Data

3.1. **Zeros of $\zeta(s)$.** We use the first 27000 zeros ($\gamma \le 23476.2$) from Odlyzko's table `zeros1` (declared accuracy $3 \cdot 10^{-9}$). The tests for $q = 4, 5$ use the first 20000 zeros, and those for $q = 8, 12$ use 27000, so that the densities of the reference cover those of the L-functions.

3.2. **Zeros of $L(s, \chi)$.** We use the real primitive characters $\chi_{-4} = (\frac{-4}{\cdot})$ (odd), $\chi_5 = (\frac{\cdot}{5})$, $\chi_8 = (\frac{8}{\cdot})$ and $\chi_{12} = (\frac{12}{\cdot})$ (even). Zeros were located by sign changes of the real function $Z_\chi(t) = e^{i\theta_\chi(t)} L(\frac{1}{2} + it, \chi)$, with L evaluated by Hurwitz zeta sums in `mpmath` at 30 digits, a scan step of at most $1/8$ of the mean spacing, and refinement to 10^{-10}. Each list was checked for (V1) agreement of the count with $\theta_\chi(T)/\pi$ at the end points, (V2) agreement with an independent rescan at half the step, (V3) a second evaluation at random zeros, and (V4) reality of Z_χ.

Sign-change scans lose pairs of very close zeros. This happened: the original lists for χ_{-4} and χ_5 failed V1/V2 (deficits of 14 and 6 zeros in 12000), and the list for χ_8 failed V2 (2000 found against 2002 in the rescan). In every case the missing zeros were recovered by rescanning all gaps larger than 1.6 mean spacings with a step of $1/32$ of the mean spacing; they were always close pairs (for χ_8, the pair 1470.2269, 1470.2791). The corrected lists pass all checks, and the corrected χ_8 list has exactly the 2002 zeros found by the rescan. We use the zeros 1–3000 for $q = 4, 5$ ($\gamma \le 2891.3$ and 2808.1) and 1–2000 for $q = 8, 12$ ($\gamma \le 1856.6$ and 1764.1).

3.3. **Derivatives and $L(2\rho, \chi^2)$.** Since $|Z'_\chi(\gamma)| = |L'(\rho, \chi)|$ at a zero, we computed $|\zeta'(\rho)|$ with `mpmath.siegelz` and $|L'(\rho, \chi)|$ by a central difference of Z_χ with step 10^{-6} at 20 digits (relative error below $4 \cdot 10^{-10}$ against step 10^{-5} at 30 digits). For real χ, χ^2 is the principal character, so $L(2\rho, \chi^2) = \zeta(1 + 2i\gamma) \prod_{p|q} (1 - p^{-1-2i\gamma})$.

4. Results

q	moment	R measured	95% interval	predicted	alternatives excluded
4	J	0.655	$[0.592, 0.715]$	2/3	1, 4/3
5	J	0.827	$[0.753, 0.901]$	5/6	1, 25/24
8	J	0.687	$[0.605, 0.784]$	2/3	1, 4/3
12	J	0.501	$[0.445, 0.555]$	1/2	1, 3/2
4	K	0.490	$[0.430, 0.549]$	1/2	1, 2/3
5	K	0.753	$[0.673, 0.835]$	4/5	1 (not 5/6)
8	K	0.558	$[0.450, 0.684]$	1/2	1 (not 2/3)
12	K	0.340	$[0.301, 0.381]$	1/3	1, 1/2

Table 1. Differential statistics R (for J_χ) and R_8 (for K_χ). Removing the five largest terms gives 0.633 ($q = 4$) and 0.794 ($q = 5$) for J, without changing any conclusion.

In all eight tests the predicted factor lies inside the interval and H_0 is excluded (Table 1). In six of them the other alternative is excluded as well. In particular, for $q = 4$ and $q = 12$ the two moments require their own factors: exchanging the factors of (2) and (3) is excluded. The

characters χ_8 and χ_{-4} have the same predicted factors but opposite parity, and their measured values are compatible.

For comparison, the absolute ratios $J_\chi(T)/(\frac{A_q}{2\pi}T)$ are 0.937, 0.937, 0.970 and 0.966 for $q = 4, 5, 8, 12$, and $K_\chi(T)/(\frac{\varphi(q)}{q}\frac{T}{2\pi})$ are 0.921, 0.871, 1.023 and 0.949. The corresponding ratios for $\zeta(s)$ over the same numbers of zeros are 0.920 and 0.932 (3000 zeros) and 0.949 and 0.959 (2000 zeros). All these values lie within a few per cent of 1, whereas the alternative without arithmetic factor would give absolute ratios between 1.2 and 3.

5. A heuristic check of the arithmetic factors

The constants in (2) and (3) can also be recovered from the random matrix model of Hughes, Keating and O'Connell [6], independently of the mollifier heuristic. For $\zeta(s)$ that model predicts $\sum_{0<\gamma\le T}|\zeta'(\rho)|^{2k} \sim \frac{T}{2\pi}\frac{G^2(k+2)}{G(2k+3)}a(k)(\log T)^{k(k+2)+1}$, where G is Barnes' function and $a(k) = \prod_p(1-\frac{1}{p})^{k^2}\,\mathbb{E}_\theta|1-p^{-1/2}e^{i\theta}|^{-2k}$; for $k=-1$ this gives (1). For a primitive character χ modulo q we keep the same random matrix factor and replace the local average by $\mathbb{E}_\theta|1-\chi(p)p^{-1/2}e^{i\theta}|^{-2k}$, which equals 1 when $p \mid q$, while the factor $(1-\frac{1}{p})^{k^2}$ is kept for every prime. For $k=1$ this reproduces the factor $\varphi(q)/q$ in the mean square $\int_0^T |L(\frac12+it,\chi)|^2\,dt$, as follows from the approximate functional equation and the mean value theorem. For $k=-1$ it gives

$$\prod_{p\nmid q}\Big(1-\frac{1}{p^2}\Big)\prod_{p\mid q}\Big(1-\frac{1}{p}\Big) = \frac{6}{\pi^2}\prod_{p\mid q}\frac{p}{p+1} = A_q.$$

In the same way, the local average of $|L_p(2\rho,\chi^2)|^2|L_p(\rho,\chi)|^{-2}$ equals $1/(1-\frac1p)$ for $p\nmid q$ and 1 for $p\mid q$, which gives the factor $\varphi(q)/q$ of (3). Thus two independent heuristics lead to the same constants. The alternatives excluded by the data also have a simple meaning in this picture: the factor $\prod_{p\mid q}p^2/(p^2-1)$ corresponds to dropping $(1-\frac1p)$ at $p\mid q$, and exchanging the factors of (2) and (3) corresponds to using the local average of one moment for the other. This is a heuristic consistency check and says nothing about lower-order terms; we do not claim it is new.

6. Regularisation and the ratios calculation

An earlier draft inferred an error term and the absence of terms of size $T/\log T$ from a diagonal-only ratios calculation. That inference is withdrawn: uniformity as the denominator shifts tend to zero was not established. The restrictions in [2, (2.11) and Remark 2.2] do not supply it for a bare reciprocal mean.

There is an exact local starting point. Let $0<a<b$ be heights that are not ordinates of zeros, and assume GRH and simplicity of the zeros of $L(s,\chi)$ in this interval. Then

$$\sum_{a<\gamma<b}\frac{1}{|L'(\rho,\chi)|^2} = \lim_{\epsilon\downarrow0}\frac{\epsilon}{\pi}\int_a^b \frac{dt}{|L(\frac12+\epsilon+it,\chi)|^2},$$

$$\sum_{a<\gamma<b}\left|\frac{L(2\rho,\chi^2)}{L'(\rho,\chi)}\right|^2 = \lim_{\epsilon\downarrow0}\frac{\epsilon}{\pi}\int_a^b \left|\frac{L(1+2\epsilon+2it,\chi^2)}{L(\frac12+\epsilon+it,\chi)}\right|^2 dt.$$

Each identity follows by isolating a simple zero and using the Cauchy kernel $\epsilon/(\epsilon^2+(t-\gamma)^2)$; its integral is π. This is a fixed-interval statement, not a uniform large-height asymptotic.

If the squared character is principal and the central value is nonzero, the numerator of the second integral has a pole at height zero. Starting the regularisation at zero adds an auxiliary-pole boundary contribution, which must be subtracted. This correction to the regularisation identity is not a complete lower-order prediction for the discrete moment; the calculation is recorded in the accompanying analytic working note.

For a formal diagonal calculation, put $\alpha = 2\epsilon$ and let χ_0 be the principal character modulo q. The two Dirichlet-series diagonals are

$$D_7(\alpha) = \frac{L(1+\alpha,\chi_0)}{L(2+2\alpha,\chi_0)}, \qquad D_8(\alpha) = L(1+\alpha,\chi_0).$$

Their residues at $\alpha = 0$ are A_q and $\varphi(q)/q$, respectively. Thus the diagonal pole is consistent with the leading constants in (2) and (3). This algebraic agreement alone does not derive the full discrete asymptotic. In particular, the finite parts of D_7, D_8 are not automatically lower-order terms of the discrete sums: the fixed-interval limit removes them, and the singular part of the remaining terms must also be controlled. Recovering the conjectures and their lower-order corrections from an appropriate ratios formula requires a joint analysis of the height and displacement limits.

7. Discussion and limitations

The data support the arithmetic factors in (2) and (3) for four real characters, including a composite conductor. The evidence is limited in several ways: the heights are low ($\gamma < 3000$); only real characters of small conductor were used; the reference set of ζ-zeros is shared between tests; the bootstrap intervals may be optimistic because of heavy tails; and the differential statistic compares two heavy-tailed sums, whose typical behaviour at the same density of zeros we assume to be comparable.

The next analytic task is a ratios calculation with the displacement dependence retained, followed by a justified passage to the discrete moments. Neither an error term nor the absence of $T/\log T$ corrections is established here. Numerical comparison should follow a frozen prediction and account for rare close pairs; additional data volume alone does not resolve this analytic issue. Complex characters and larger conductors remain possible tests of the arithmetic factors.

8. Literature search

We searched zbMATH Open, the arXiv, and the works citing Gonek's paper on negative moments (via OpenAlex) for numerical computations of $L'(\rho,\chi)$ at zeros of Dirichlet L-functions or of the sums J_χ, K_χ, and found none; the only numerical study of the corresponding moments of ζ that we found is [5]. MathSciNet was not available to us.

Data and code

The zeros, derivatives, pre-registration documents and analysis code, including a script that reproduces Table 1, are available at `https://doi.org/10.5281/zenodo.23020957`. All computations were run on a single laptop with two cores; the computation and verification of the zeros of the L-functions dominated the cost.

Acknowledgements

The original computations, analysis and drafting used substantial assistance from Claude (Anthropic). The present analytic revision also used Codex (OpenAI) and remains subject to the author's mathematical review before circulation.

References

[1] J. B. Conrey, D. W. Farmer, M. R. Zirnbauer, Autocorrelation of ratios of L-functions, Comm. Number Theory Phys. 2 (2008), no. 3, 593–636.

[2] J. B. Conrey, N. C. Snaith, Applications of the L-functions ratios conjectures, Proc. London Math. Soc. (3) 94 (2007), no. 3, 594–646.

[3] D. W. Farmer, Long mollifiers of the Riemann zeta-function, Mathematika 40 (1993), no. 1, 71–87.

[4] S. M. Gonek, The second moment of the reciprocal of the Riemann zeta-function and its derivative, talk at the Mathematical Sciences Research Institute, Berkeley, June 1999.

[5] G. A. Hiary, A. M. Odlyzko, Numerical study of the derivative of the Riemann zeta function at zeros, Comment. Math. Univ. St. Pauli 60 (2011), 47–60.
[6] C. P. Hughes, J. P. Keating, N. O'Connell, Random matrix theory and the derivative of the Riemann zeta function, Proc. R. Soc. Lond. A 456 (2000), 2611–2627.
[7] M. B. Milinovich, N. Ng, A note on a conjecture of Gonek, Funct. Approx. Comment. Math. 46.2 (2012), 177–187.
[8] N. Ng, Limiting distributions and zeros of Artin L-functions, PhD thesis, University of British Columbia, 2000.
[9] A. Pearce-Crump, Negative discrete second moments of Dirichlet L-functions, arXiv:2606.25094 (2026).
[10] S. Sinha, Some Conditional Results involving Arithmetic Functions, PhD thesis, University of Missouri, 2025.

Independent researcher. ORCID: 0009-0009-8217-1623
Email address: dionatandieresende@gmail.com