

Fuzzy Logic : Introduction




Lecture 2


1

Lotfi Zadeh

*As the complexity of a system increases,
our ability to make precise and yet significant
statements about its behaviour diminishes until a
threshold is reached beyond which precision and
significance (or relevance) become almost
mutually exclusive characteristics*



2



3

Fuzzy Logic

Consider the following rule:

If the patient has *high* fever
and he is shivering *a lot*
then influenza is *highly likely*

4


Fuzzy Rule

The words shown in *italics* are imprecise terms. These are called *fuzzy predicates*.


Such a rule is termed as a *fuzzy rule*.

5

Conversion of conventional method into fuzzy method



6



Outline

- Crisp Set and Fuzzy Set
- Membership Function
- Fuzzy Set Operation
- Terminologies

8

Ordinary (Crisp) Sets to Fuzzy Sets



9

World of Difference




- **Crisp Set:** To dichotomize the individuals in some given universe of discourse into two groups: members (those that certainly belong in the set) and non members (those that certainly do not).
- **Fuzzy Set:** mathematically by assigning to each possible individual in the universe of discourse a value representing its grade of membership in the fuzzy set


10

Examples

Crisp Set Theory:



Fuzzy Set Theory:



11

Crisp Sets: An Overview

- $Z = \{ \dots, -2, -1, 0, 1, 2, \dots \}$ – The set of all integers
- $N = \{ 1, 2, 3, \dots \}$ – The set of all positive integers or natural numbers
- $N_0 = \{ 0, 1, 2, \dots \}$ – The set of all non-negative integers
- $N_n = \{ 1, 2, \dots, n \}$
- $N_{0,n} = \{ 0, 1, 2, \dots, n \}$
- R : The set of all real numbers
- R^+ : The set of all nonnegative real numbers

Cont. 12

Crisp Sets: An Overview

- $[a, b]$, $(a, b]$, $[a, b)$, (a, b) : Closed, left-open, right open, open interval or real numbers between a and b , respectively.
- $\langle x_1, x_2, \dots, x_n \rangle$: ordered n -tuple of elements x_1, x_2, \dots, x_n

Cont...

13

Crisp Sets: An Overview

- Three basic Methods to define set within a given universal set X
- **List Method**: for finite sets
 $A = \{a_1, a_2, \dots, a_n\}$
- **Rule Method**: for property based sets
 $A = \{x \mid P(x)\}$
 $P(x)$ is true or false and $x \in X$

Cont...

14

Crisp Sets: An Overview

- **Characteristic Function** : declares which elements of X are members of the set and which are not.
- Set A is defined by its characteristic function χ_A , as follows:

$$\chi_A(X) = \begin{cases} 1 & \text{for } x \in A \\ 0 & \text{for } x \notin A \end{cases} \quad \chi_A : X \rightarrow \{0,1\}$$

15

Fundamental Properties of Crisp set Operation

- Involution $\overline{\overline{A}} = A$
- Commutative $A \cup B = B \cup A, A \cap B = B \cap A$
- Associativity $(A \cup B) \cup C = A \cup (B \cup C)$
 $(A \cap B) \cap C = A \cap (B \cap C)$
- Distributivity $A \cap (B \cup C) = (A \cap B) \cup (A \cap C)$
 $A \cup (B \cap C) = (A \cup B) \cap (A \cup C)$
-

16

Crisp Sets: An Overview

Homework : General background from recommended text book of *Klir and Yuan*

17

Fuzzy Sets : Basic Types

- The characteristic function of crisp set assigns value of either 1 or 0 to each individual in the universal set.
- Two distinct notations are most commonly used to denote membership functions

18

Fuzzy Sets : Basic Types

Membership function of a fuzzy set A is denoted by μ_A ; that is


$$\mu_A : X \rightarrow [0,1]$$

The function is denoted by A and has the same form:

$$A : X \rightarrow [0,1]$$


19

Membership Function




20

Membership Function



21

Membership Function




22

Membership Function

$$A_4(x) = \begin{cases} (1 + \cos(p_4\pi(x-r)))/2 & \text{when } x \in [r-1/p_4, r+1/p_4] \\ 0 & \text{otherwise} \end{cases}$$

$$p_4=2, r=2$$



23


Simple Example

- Fuzzy sets
(Three Types)
 - Little educated
 - Highly educated
 - Very highly educated
- Finite Universal set consists of seven levels of education
 - 0 – No education
 - 1 – Elementary school
 - 2 – High school
 - 3 – Two-year college degree
 - 4 – Bachelor's degree
 - 5 – Master's degree
 - 6 – Doctoral Degree

24



Little educated
Highly educated
Very highly educated

Fuzzy Set Expression




25

Temperature in the range (a) Fuzzy (b) a traditional variables



26

Example Fuzzy Variable



- Each function tells us how much we consider a character in the set if it has a particular aggressiveness value
- Or, how much truth to attribute to the statement: “The character is nasty (or meek, or neither)?”


27

Basic Operations on Fuzzy Sets

- Complement
- Intersection
- Union

28


Fuzzy Set Operations: Complement



- The degree to which you believe something is **not** in the set is 1.0 minus the degree to which you believe it is in the set

29

Fuzzy Set Operations: Intersection (AND)




- If you have x degree of faith in statement A, and y degree of faith in statement B, how much faith do you have in the statement A and B?
 - Eg: How much faith in “that person is about 6' high and tall”
- Does it make sense to attribute more truth than you have in one of A or B?

$$(A \cap B)(x) = \min[A(x), B(x)]$$

30

Fuzzy Set Operations: Union (OR)

Membership



- If you have x degree of faith in statement A, and y degree of faith in statement B, how much faith do you have in the statement A or B?
 - Eg: How much faith in “that person is about 6' high or tall”
- Does it make sense to attribute **less truth** than you have in one of A or B?

$$(A \cup B)(x) = \max[A(x), B(x)]$$

31

CAR EXAMPLE

32

Car Example

- “If our *distance to the car in front* is *small*, and the *distance is decreasing slowly*, then *decelerate quite hard*”
 - Fuzzy variables in blue
 - Fuzzy sets in red
 - Conditions are on membership in fuzzy sets
 - Actions place an output variable (decelerate) in a fuzzy set (the quite hard deceleration set)
- We have a certain belief in the truth of the condition, and hence a certain strength of desire for the outcome
- Multiple rules may match to some degree, so we require a means to arbitrate and choose a particular goal - *defuzzification*

33

Car Examples

- Rules for controlling a car:
 - Variables are *distance* to car in front and how fast it is changing, *delta*, and *acceleration* to apply
 - Sets are:
 - Very small, small, perfect, big, very big - for distance
 - Shrinking fast, shrinking, stable, growing, growing fast for delta
 - Brake hard, slow down, none, speed up, floor it for acceleration
 - Rules for every combination of distance and delta sets, defining an acceleration set


34

Car Example

- Assume we have a particular numerical value for distance and delta, and we need to set a numerical value for acceleration
 - Extension: Allow fuzzy values for input variables (degree to which we believe the value is correct)


35

Set Definitions for Example




36

Instance for Example



37

Instance for Example




38

Matching for Example

- Relevant rules are:
 - If distance is small and delta is growing, maintain speed
 - If distance is small and delta is stable, slow down
 - If distance is perfect and delta is growing, speed up
 - If distance is perfect and delta is stable, maintain speed
- For first rule, distance is small has 0.75 truth, and delta is growing has 0.3 truth
 - So the truth of the **and** is 0.3
- Other rule strengths are 0.6, 0.1 and 0.1

39



40