

Learn DAA : From B K Sharma

TCS-503: Design and Analysis of Algorithms

Unit V: Selected Topics: NP
Completeness


Unit V

- Selected Topics:
 - NP Completeness
 - Approximation Algorithms
 - Randomized Algorithms
 - String Matching

Learn DAA : From B K Sharma

NP-completeness

Travelling Salesman Problem, Clique Problem.....



I can't find an efficient algorithm for,

But neither can these famous people

Learn DAA : From B K Sharma

NP-completeness

Three key concepts for showing a problem to be NP-C are

Decision Problem vs Optimization Problem

Reductions $A \leq_p B$

A first NP- Complete Problem

Learn DAA : From B K Sharma

NP-completeness

A First NP-Complete Problem


The technique for showing that a problem is in the class NP-Complete requires that we have one NP-Complete problem to begin with.

Cook First time showed that Circuit Satisfiability problem is in class NPC.

All proofs ultimately follow by reduction from NP-Completeness of Circuit-Satisfiability.

Learn DAA : From B K Sharma



The Structure of *NP-completeness*



Learn DAA : From B K Sharma

What is Circuit-Sat Problem?

We are given a Boolean combinational circuit composed of AND, OR, and NOT gates and we wish to know whether there is any set of Boolean inputs to this circuit that causes its output to be 1.



The assignment $x_1=1$, $x_2=1$ and $x_3=0$ to the inputs of this circuit causes the output of the circuit to be 1.

Learn DAA : From B K Sharma

What is Formula Satisfiability(SAT) Problem?

A Boolean Formula Φ given, find an assignment of values (0,1) to the variables of Φ that causes it to evaluate to 1.

For example:

$$\Phi = (x_1 \vee x_2) \wedge (x_1 \vee \neg x_2) \wedge (\neg x_1 \vee \neg x_2)$$

Let Certificate be :

$$x_1 = 1, x_2 = 0,$$

(a. k. a. assignment)

$$\Phi = 1 \wedge 1 \wedge 1 = 1.$$

Formula Satisfiability is first to be proven NP-Complete.

Learn DAA : From B K Sharma

Prove that SAT is NPC

To prove that SAT is NPC:

1. First we prove that $SAT \in NP$ and then
2. SAT is NP-hard by $CIRCUIT-SAT \leq_p SAT$

To show that SAT belongs to NP, we show that a certificate consisting of a satisfying assignment for an input formula φ can be verified in polynomial time.

Prove that SAT is NPC

(1) SAT \in NP


Let the formula be

$\Phi = (x_1 \vee x_2) \wedge (x_1 \vee \neg x_2) \wedge (\neg x_1 \vee \neg x_2)$ and **Certificate:** $x_1 = 1, x_2 = 0$


Then $\Phi = 1 \wedge 1 \wedge 1 = 1$

This can be easily verified that Φ is satisfiable in polynomial time $O(n)$, where n is the number of clauses (steps) in the formula.

(2) SAT is NPH ,i.e. CIRCUIT-SAT \leq_p SAT



$$\begin{aligned}\Phi = & x_6 \wedge (x_4 \leftrightarrow x_1 \vee x_2) \\ & \wedge (x_5 \leftrightarrow \neg x_3) \\ & \wedge x_6 \leftrightarrow (x_4 \wedge x_5)\end{aligned}$$




$$\begin{aligned}\Phi = & 1 \wedge (1 \leftrightarrow 1 \vee 1) \\ & \wedge (1 \leftrightarrow 1) \\ & \wedge (1 \leftrightarrow (1 \wedge 1)) \\ = & 1 \wedge 1 \wedge 1 \wedge 1 \\ = & 1\end{aligned}$$

Given the Circuit, the formula Φ can be produced in Polynomial Time $O(n + m)$, where n is number of wires (6) and m is number of gates(3).

Learn DAA : From B K Sharma

Looking Forward


From now on, in order to show some NP problem is NP-C, we merely need to reduce SAT to it.



Learn DAA : From B K Sharma

... and Beyond!

Moreover, every NP-problem we discover, provides us with a new way for showing problems to be NP-C.



Learn DAA : From B K Sharma

Prove that 3-CNF-SAT is NPC

(1) 3-CNF-SAT \in NP

Let $\Phi = (x_1 \vee \neg x_1 \vee \neg x_2) \wedge (x_3 \vee x_2 \vee x_4) \wedge (\neg x_1 \vee \neg x_3 \vee \neg x_4)$ and

Certificate: $x_1=1, x_2=1, x_3=0, x_4=1$

Then $\Phi = (1 \vee 0 \vee 0) \wedge (0 \vee 1 \vee 1) \wedge (0 \vee 1 \vee 0)$
 $= 1 \wedge 1 \wedge 1 = 1$

Thus, the formula Φ is satisfiable.

This can be easily verified that Φ is satisfiable in polynomial time $O(n)$, where n is the number of clauses (steps) in the formula.

Thus, 3-CNF-SAT \in NP.

Learn DAA : From B K Sharma

Prove that 3-CNF-SAT is NPC

(2) 3-CNF-SAT is NPH, i.e., $SAT \leq_p 3\text{-CNF-SAT}$.

The reduction algorithm consists of three steps:

- 2(a). The first step converts given formula into CNF.
- 2(b) The second step transforms the formula so that each clause has exactly three distinct literals., that is in 3-CNF.
- 2(c) We must also show that the reduction can be computed in polynomial time.

Learn DAA : From B K Sharma

Prove that 3-CNF-SAT is NPC

(2) 3-CNF-SAT is NPH, i.e., $SAT \leq_p 3\text{-CNF-SAT}$.

2(a). The first step converts given formula into CNF.

Let $\Phi = x_1 \leftrightarrow (x_2 \wedge x_3)$

$$= (x_1 \rightarrow (x_2 \wedge x_3)) \wedge ((x_2 \wedge x_3) \rightarrow x_1)$$

Since, $[P \leftrightarrow Q \equiv (P \rightarrow Q) \wedge (Q \rightarrow P)]$

$$= (\neg x_1 \vee (x_2 \wedge x_3)) \wedge (\neg(x_2 \wedge x_3) \vee x_1)$$

[Since, $P \rightarrow Q \equiv \neg P \vee Q$]

$$\Phi_1 = (\neg x_1 \vee x_2) \wedge (\neg x_1 \vee x_3) \wedge (\neg x_2 \vee \neg x_3 \vee x_1)$$

[Distributive law $P \vee (Q \wedge R) \equiv (P \vee Q) \wedge (P \vee R)$ and De

Morgan's Laws $\neg(P \wedge Q) \equiv \neg P \vee \neg Q$]

which is in CNF.

Prove that 3-CNF-SAT is NPC

(2) 3-CNF-SAT is NPH, i.e., $SAT \leq_p 3\text{-CNF-SAT}$.

2(b) The second step transforms the formula so that each clause has exactly three distinct literals., that is in 3-CNF.

$$\begin{aligned}\Phi_1 &= (\neg x_1 \vee x_2) \wedge (\neg x_1 \vee x_3) \wedge (\neg x_2 \vee \neg x_3 \vee x_1) \\ &= C_1 \quad \wedge \quad C_2 \quad \wedge \quad C_3\end{aligned}$$

Since, C_1 and C_2 have 2 distinct literals,

$$\Phi_2 = (\neg x_1 \vee x_2 \vee y_1) \wedge (\neg x_1 \vee x_2 \vee \neg y_1) \wedge$$

Since, if $y_1=0$, $\neg y_1=1$ or if $y_1=1$, $\neg y_1=0$

$$(\neg x_1 \vee x_3 \vee y_2) \wedge (\neg x_1 \vee x_3 \vee \neg y_2)$$

Since, if $y_2=0$, $\neg y_2=1$ or if $y_2=1$, $\neg y_2=0$

$$\wedge (\neg x_2 \vee \neg x_3 \vee x_1)$$

Prove that 3-CNF-SAT is NPC

(2) 3-CNF-SAT is NPH, i.e., $SAT \leq_p 3\text{-CNF-SAT}$.

2(c) We must also show that the reduction can be computed in polynomial time.

- ✓ The Construction of Φ_1 from Φ has introduced 1 more clause.

[Φ Contains 2 clauses, Φ_1 contains 3 clauses.]


- ✓ The construction of Φ_2 from Φ_1 has introduced 2 more clauses.

[Φ_1 Contains 3 clauses, Φ_2 contains 5 clauses.]

Thus, the size of the resulting formula Φ_2 is polynomial in the length of the original formula Φ .
 Φ has two clauses($n=2$). Φ_2 has 5 clauses ($n^2 + 1$).


What is Complete Sub-graph?

Complete Sub-graph: a subset of vertices of a graph fully connected to each other. One where there is an edge between every two vertices.




What is Clique?

Clique is a complete sub-graph of a graph.



Learn DAA : From B K Sharma

What is Clique?




k -clique = complete subgraph of k nodes

Learn DAA : From B K Sharma

What is Clique Problem?

Find the clique with the **largest number of vertices** in a graph.


This graph has Clique of size 2



$\text{Clique}(G, 2) = \text{YES}$

$\text{Clique}(G, 3) = \text{NO}$

This graph has Clique of size 3



$\text{Clique}(G, 3) = \text{YES}$

$\text{Clique}(G, 4) = \text{NO}$

Optimization problem

Find a clique of maximum size.

Restated as Decision problem

Does Graph have a **clique of size k**?


Learn DAA : From B K Sharma

Prove that Clique Problem is NPC?

1) Clique \in NP

1(a) To show that CLIQUE \in NP, for a given graph $G = (V, E)$, we use the set $V' \subseteq V$ of vertices in the clique as a certificate for G .

$V = \{A, B, C, D\}$



Certificate: $V' = \{A, B\}$


1(b) Checking whether V' is a clique can be accomplished in polynomial time by checking whether, for each pair $u, v \in V'$, the edge (u, v) belongs to E .

Learn DAA : From B K Sharma

Prove that Clique Problem is NPC?

1) Clique \in NP

1(b) *Checking whether V' is a clique can be accomplished in polynomial time by checking whether, for each pair $u, v \in V'$, the edge (u, v) belongs to E .*



Check, is there an edge (A, B) and B, A ? Answer is Yes.
Similarly we can check $V' = \{A, C\}$, $\{AD\}$, $\{BC\}$, $\{C, D\}$ etc.

Learn DAA : From B K Sharma

Prove that Clique Problem is NPC?

1) Clique \in NP

The length of the certificate: $O(n)$ ($n=|V|$)

Time complexity: $O(n^2)$

Learn DAA : From B K Sharma

Prove that Clique Problem is NPC?

1) Clique \in NP

Given a subset V' of V

Verify $|V'|=k$

$O(k)$ time.

Verify every pair of vertices in $|V'|$ have an edge in E .

$O(k^2 |E|)$ time.

Learn DAA : From B K Sharma

Prove that Clique Problem is NPC?

2) Clique is NPH

3-CNF-SAT \leq_p CLIQUE

Let $\Phi = C_1 \wedge C_2 \wedge \dots \wedge C_k$ be a 3-CNF formula with k clauses, each of which has 3 distinct literals.

For example,

$$\begin{aligned}\Phi &= (x_1 \vee x_1 \vee x_2) \wedge (\neg x_1 \vee \neg x_2 \vee \neg x_2) \wedge (\neg x_1 \vee x_2 \vee x_2) \\ &= C_1 \quad \quad \quad \wedge \quad \quad \quad C_2 \quad \quad \quad \wedge \quad \quad \quad C_3\end{aligned}$$








For each clause put a triple of vertices in the graph,
one for each literal.

Learn DAA : From B K Sharma

Prove that Clique Problem is NPC?

2) Clique is NPH

3-CNF-SAT \leq_p CLIQUE




Learn DAA : From B K Sharma

Prove that Clique Problem is NPC?

2) Clique is NPH

$3\text{-CNF-SAT} \leq_p \text{CLIQUE}$

Put an edge between two vertices if they are in different triples and their literals are *consistent*, meaning not each other's negation.



#nodes = 3(# clauses)

$k = \text{\#clauses}$

Learn DAA : From B K Sharma

Prove that Clique Problem is NPC?

2) Clique is NPH

$$\begin{aligned}\Phi &= (x_1 \vee x_1 \vee x_2) \wedge (\neg x_1 \vee \neg x_2 \vee \neg x_2) \wedge (\neg x_1 \vee x_2 \vee x_2) \\ &= (0 \vee 0 \vee 1) \wedge (1 \vee 0 \vee 0) \wedge (1 \vee 1 \vee 1) \\ &= 1 \wedge 1 \wedge 1 = 1\end{aligned}$$

Now Pose Question:

Question : Is ϕ satisfiable? **Answer:** Yes: $\neg x_1 = 1$ (true) and $x_2 = 1$ (true)

assign the value **true** to every variable occurring in the clique)

Learn DAA : From B K Sharma

Prove that Clique Problem is NPC?

2) Clique is NPH

$3\text{-CNF-SAT} \leq_p \text{CLIQUE}$

The 3-cnf formula is satisfiable iff the graph has a k-clique.

The 3-CNF formula ϕ has $m=3$ -clauses and is satisfiable then the Clique must have the size $k=3$.

Learn DAA : From B K Sharma

Prove that Clique Problem is NPC?


2) Clique is NPH

What's left?

Demonstrate the reduction is polynomial

$$|E| \leq (\# \text{ of literals in } \varphi)^2 = O(n^2)$$


$$\begin{aligned}
 & (x_1 \vee x_1 \vee x_1) \wedge (\neg x_1 \vee \neg x_1 \vee x_2) \wedge \\
 & (x_2 \vee x_2 \vee x_2) \wedge (\neg x_2 \vee \neg x_2 \vee x_1)
 \end{aligned}$$



Learn DAA : From B K Sharma

Independent Set

Given a graph $G(V,E)$, an **independent set** V' is a collection of nodes in G such that no two nodes in V' have an edge connecting them.




Learn DAA : From B K Sharma

Independent Set

An **independent set** is a set of nodes with no edges between them.

independent sets

\emptyset , {1}, {2}, {3}, {4}, {1, 2}, {1, 3}



Learn DAA : From B K Sharma

Complement (G^c) of a graph (G)



The complement G^c of a graph G contains exactly those edges not in G .

$$G = (V, E) \Rightarrow G^c = (V, E^c)$$

$$E^c = \{(u, v) : u, v \in V, \text{ and } (u, v) \notin E\}$$




Complement (G^c) of a graph (G)

The complement G^c of a graph G contains exactly those edges not in G .



Vertex Cover


A **vertex cover** for a graph G is a subset of vertices incident to every edge in G .



Size of a vertex cover: the number of vertices in it.

Learn DAA : From B K Sharma

Vertex Cover



vertex cover = set of vertices that cover all edges

Learn DAA : From B K Sharma

Vertex Cover Problem

Optimization Problem

What is the minimum size vertex cover in G ?

Restated as a decision problem:

Does a vertex cover of size k exist in G ?

Learn DAA : From B K Sharma

Prove that Vertex Cover Problem is NPC

$\text{CLIQUE} \leq_p \text{VERTEX-COVER}$

Reduce k -clique to vertex cover


Learn DAA : From B K Sharma

Prove that Vertex Cover Problem is NPC


Compute G_c in polynomial time

G has a clique of size k iff G^c has a vertex cover of size $|V| - k$


$\langle G, k \rangle$ (clique) $\rightarrow \langle G^c, |V| - k \rangle$ (vertex cover)




Clique = 2



$V - C = 2$



Clique = 2




$V - C = 3$

G has a clique of size $k \Leftrightarrow G^c$ has a vertex cover of size $n - k$


Learn DAA : From B K Sharma

Prove that Vertex Cover Problem is NPC



(a)

A graph $G(V, E)$ with CLIQUE $V' = \{u, v, x, y\}$



(b)

The graph G^C produced by the reduction algorithm that has vertex cover $V - V' = \{w, z\}$