

1) Using a goodness of fit, we can assess whether a set of obtained frequencies differ from a set of frequencies.

a) Mean b) Actual c) Predicted d) Expected

Answer: d) Expected

2) Chi-square is used to analyse

a) Score b) Rank c) Frequencies d) All of these

Answer: c) Frequency

3) What is the mean of a Chi Square distribution with 6 degrees of freedom?

a) 4 b) 12 c) 6 d) 8

Answer: c) 6

4) Which of these distributions is used for a goodness of fit testing?

a) Normal distribution b) Chi-squared distribution c) Gamma distribution d) Poisson distribution

Answer: b) Chi-squared distribution

5) Which of the following distributions is Continuous

a) Binomial Distribution b) Hypergeometric Distribution c) F Distribution d) Poisson Distribution

Answer: c) F Distribution

6) A statement made about a population for testing purpose is called?

a) Statistic b) Hypothesis c) Level of Significance d) Test Statistic

Answer: b) Hypothesis

7) If the assumed hypothesis is tested for rejection considering it to be true is called?

a) Null Hypothesis b) Statistical Hypothesis c) Simple Hypothesis d) Composite Hypothesis

Answer: a) Null Hypothesis

8) If the Critical region is evenly distributed then the test is referred as?

a) Two tailed b) One tailed c) Three tailed d) Zero tailed

Answer: a) Two Tailed

9) Alternative Hypothesis is also called as?

a) Composite hypothesis b) Research Hypothesis c) Simple Hypothesis d) Null Hypothesis

Answer: b) Research Hypothesis

10) In a Binomial Distribution, if 'n' is the number of trials and 'p' is the probability of success, then the mean value is given by

a) np b) n

Answer: a) np