

Galois Connections

A Complete Introduction

David A. Smith, Direct Knowledge, USA ^{*†‡}

Thursday, February 9, 2023

Table of contents

1	Introduction	1
2	What are Galois Connections?	1

Example 0.1. Show that the statement

$$(p \wedge \neg q) \rightarrow [(\neg p \vee \neg q) \rightarrow (p \wedge \neg q)] \quad (1)$$

is a tautology.

1 Introduction

2 What are Galois Connections?

Let (X, \preceq) and (Y, \leq) be partially ordered sets.

Definition 2.1. If $f_* : X \rightarrow Y$ and $f^* : Y \rightarrow X$ are functions such that

$$f_*(x) \leq y \iff x \preceq f^*(y) \quad (2)$$

^{*}email: david@directknowledge.com

[†]© 2023 David A. Smith

[‡]With authorization, you can freely share and reproduce portions of this work for educational or personal use. Please note that distributing any portion of it in print form requires further permission from its original authors, as does posting online to public servers or mailing lists without prior consent.

for all $x \in X$ and all $y \in Y$, then (f_*, f^*) is called a **Galois connection** between (X, \preceq) and (Y, \leq) .

There are several definitions of Galois connections in the literature; however they are all order-isomorphic to the definition above.

Proposition 2.1. *Let $f_* : X \rightarrow Y$ and $f^* : Y \rightarrow X$ be functions. Then (f_*, f^*) is a Galois connection if and only if*

1. both f_* and f^* are monotone,
2. $x \preceq (f^* \circ f_*)(x)$ for all $x \in X$, and
3. $(f_* \circ f^*)(y) \leq y$ for all $y \in Y$.

Proof. Suppose (f_*, f^*) is a Galois connection. By (2) it follows

$$f_*(x) \leq f_*(x) \iff x \preceq (f^* \circ f_*)(x).$$

Since \leq is reflexive, $x \preceq (f^* \circ f_*)(x)$ follows immediately. Similarly, by (2) it follows

$$(f_* \circ f^*)(y) \leq y \iff f^*(y) \preceq f^*(y)$$

proving that (??) also holds. Assume $x_1 \preceq x_2$. By (??) we have $x_2 \preceq (f^* \circ f_*)(x_2)$. By (2) it follows $f_*(x_1) \leq f_*(x_2)$ and so f_* is monotone. Assume $y_1 \leq y_2$. By (??) we have $(f_* \circ f^*)(y_1) \leq y_1$. By (2) it follows $f^*(y_1) \preceq f^*(y_2)$ and so f^* is also monotone.

Conversely, assume (??), (??), and (??) all hold. Assume $f_*(x) \leq y$. By (??) and (??), it follows $(f^* \circ f_*)(x) \preceq f^*(y)$ and $x \preceq (f^* \circ f_*)(x)$, respectively. By transitivity of \preceq , we have $x \preceq f^*(y)$ as needed. Assume $x \preceq f^*(y)$. By (??) and (??), it follows $f_*(x) \leq (f_* \circ f^*)(y)$ and $(f_* \circ f^*)(y) \leq y$. By transitivity of \leq , we have $f_*(x) \leq y$ as needed. Therefore, (2) holds and so (f_*, f^*) is a Galois connection. \square

Proposition 2.2. *If (f_*, g) and (f_*, h) are Galois connections between (X, \preceq) and (Y, \leq) , then $g = h$. Likewise, if (g, f^*) and (h, f^*) are Galois connections between (X, \preceq) and (Y, \leq) , then $g = h$.*

Proof. Let $f_* : X \rightarrow Y$ and $g : Y \rightarrow X$ be a Galois connection. Also let f_* and $h : Y \rightarrow X$ be a Galois connection. By (2) we have the following

$$f_*(x) \leq y \iff x \preceq g(y) \tag{3}$$

$$f_*(x) \leq y \iff x \preceq h(y) \quad (4)$$

By (3) we have $(f_* \circ h)(y) \leq y \iff h(y) \preceq g(y)$. Notice $(f_* \circ h)(y) \leq y$ holds by (??).(??); and thus $h(y) \preceq g(y)$. By (4) we have $(f_* \circ g)(y) \leq y \iff g(y) \preceq h(y)$. Notice $(f_* \circ g)(y) \leq y$ holds by (??).(??); and thus $h(y) \preceq g(y)$. Since \preceq is antisymmetric, it follows $g(y) = h(y)$ for arbitrary y . The second statement is the dual of the first and follows just as easily using (??).(??). \square

If (f, g) and (f, h) are Galois connections between (P, \leq_P) and (Q, \leq_Q) , then $g = h$. To see this, observe that $p \leq_P g(q)$ iff $f(p) \leq_Q q$ iff $p \leq_P h(q)$, for any $p \in P$ and $q \in Q$. In particular, setting $p = g(q)$, we get $g(q) \leq_P h(q)$ since $g(q) \leq_P g(q)$. Similarly, $h(q) \leq_P g(q)$, and therefore $g = h$. By a similar argument, if (g, f) and (h, f) are Galois connections between (P, \leq_P) and (Q, \leq_Q) , then $g = h$. Because of this uniqueness property, in a Galois connection $f = (f^*, f_*)$, f^* is called **the upper adjoint** of f_* and f_* **the lower adjoint** of f^* .

Proposition 2.3. *If (f_*, f^*) is a Galois connection between (X, \preceq) and (Y, \leq) , then*

1. $f^*(y)$ is the maximum of $\{x \in X : f_*(x) \leq y\}$ and
2. $f_*(x)$ is the minimum of $\{y \in Y : x \preceq f^*(y)\}$.

Proof. Let $M = \{x \in X : f_*(x) \leq y\}$. By (??).(??) we have $f^*(y) \in M$. Let $x \in M$. Then $f_*(x) \leq y$ and since f^* is monotone, it follows $(f^* \circ f_*)(x) \preceq f^*(y)$. By (??).(??), we have $x \preceq (f^* \circ f_*)(x)$. By transitivity of \preceq , we have $x \preceq f^*(y)$ and thus $f^*(y)$ is the maximum of M . For the second statement, let $N = \{y \in Y : x \preceq f^*(y)\}$. By (??).(??) we have $f_*(x) \in N$. Let $y \in N$. Then $x \preceq f^*(y)$ and since f_* is monotone, it follows $f_*(x) \leq (f_* \circ f^*)(y)$. By (??).(??), we have $(f_* \circ f^*)(y) \leq y$ and so by transitivity, it follows $f_*(x) \leq y$. Thus $f_*(x)$ is the minimum of N . \square

Proposition 2.4. *If (f_*, f^*) is a Galois connection between (X, \preceq) and (Y, \leq) , then*

1. $f_* \circ f^* \circ f_* = f_*$, $f^* \circ f_* \circ f^* = f^*$,
2. $x \in f^*(Y)$ if and only if x is a fixed point of $f^* \circ f_*$,
3. $y \in f_*(X)$ if and only if y is a fixed point of $f_* \circ f^*$,
4. $f^*(Y) = (f^* \circ f_*)(X)$, and $f_*(X) = (f_* \circ f^*)(Y)$.

Proof. (??): Using (??), we have $f_*(x) \leq (f_* \circ f^* \circ f_*)(x)$. By (2) with $x := (f^* \circ f_*)(x)$ and $y := f_*(x)$ it follows $(f_* \circ f^* \circ f_*)(x) \leq f_*(x)$ using that \preceq is reflexive.

Since \leq is antisymmetric, it follows $f_*(x) = (f_* \circ f^* \circ f_*)(x)$ for arbitrary x , thus proving (??) holds. (??): By definition, $x \in f^*(Y)$ is equivalent to $f^*(y) = x$ for some $y \in Y$. Then

$$(f^* \circ f_*)(x) = (f^* \circ f_* \circ f^*)(y) = f^*(y) = x$$

follows by (??). (??): It follows by (??) that $f^*(Y) \subseteq (f^* \circ f_*)(X)$. Conversely, let $x \in (f^* \circ f_*)(X)$. Then $x = f^*(y)$ for some $y \in f_*(X) \subseteq Y$. By definition, $x \in f^*(Y)$ and so $(f^* \circ f_*)(X) \subseteq f^*(Y)$. \square

Proposition 2.5. *If (f_*, f^*) is a Galois connection between (X, \preceq) and (Y, \leq) , then*

1. $x \preceq f^*(y) \iff f_*(x) \leq (f_* \circ f^*)(y) \iff f_*(x) \leq y \iff (f^* \circ f_*)(x) \preceq f^*(y)$,
2. $f^*(x) \preceq f^*(y) \iff (f_* \circ f^*)(x) \leq (f_* \circ f^*)(y) \iff (f_* \circ f^*)(x) \leq y$, and
3. $f_*(x) \leq f_*(y) \iff (f^* \circ f_*)(x) \preceq (f^* \circ f_*)(y) \iff x \preceq (f^* \circ f_*)(x)$,
4. $f^*(x) = f^*(y) \iff (f_* \circ f^*)(x) = (f_* \circ f^*)(y)$, and
5. $f_*(x) = f_*(y) \iff (f^* \circ f_*)(x) = (f^* \circ f_*)(y)$.

Proof. For the first statement we have

$$\begin{aligned} x \preceq f^*(y) &\implies f_*(x) \leq (f_* \circ f^*)(y) \implies f_*(x) \leq y \\ &\implies (f^* \circ f_*)(x) \preceq f^*(y) \implies x \preceq f^*(y) \end{aligned}$$

For the third statement we have

$$f^*(x) \preceq f^*(y) \implies (f_* \circ f^*)(x) \leq (f_* \circ f^*)(y) \implies (f_* \circ f^*)(x) \leq y$$

For the fourth statement we have

$$\begin{aligned} f^*(x) = f^*(y) &\iff f^*(x) \preceq f^*(y) \wedge f^*(y) \preceq f^*(x) \\ &\iff (f_* \circ f^*)(x) \leq (f_* \circ f^*)(y) \wedge (f_* \circ f^*)(y) \leq (f_* \circ f^*)(x) \\ &\iff (f_* \circ f^*)(x) = (f_* \circ f^*)(y) \end{aligned}$$

The remaining statements are the dual and easily proved. \square

By an **order isomorphism** from an partially ordered set X to another partially ordered set Y we shall mean an isotone bijection $f : X \rightarrow Y$ whose inverse $f^{-1} : Y \rightarrow X$ is also an isotone.

Proposition 2.6. *Partially ordered sets (X, \preceq) and (Y, \leq) are isomorphic if and only if there is a surjective mapping $f : X \rightarrow Y$ such that*

$$x \preceq y \iff f(x) \leq f(y).$$

Proof. The necessity is clear. Suppose conversely that such a surjective mapping f exists. Then f is also injective; for if $f(x) = f(y)$ then from $f(x) \leq f(y)$ we obtain $x \preceq y$ and from $f(y) \leq f(x)$ we obtain $y \preceq x$, so that $x = y$. Hence f is a bijection. Clearly, f is isotone; and so also is f^{-1} , since $x \preceq y$ can be written $f(f^{-1}(x)) \leq f(f^{-1}(y))$ which gives $f^{-1}(x) \preceq f^{-1}(y)$. \square

Proposition 2.7. *If (f_*, f^*) is a Galois connection between (X, \preceq) and (Y, \leq) , then $f^*(Y)$ and $f_*(X)$ are order-isomorphic.*

Proof. This follows immediately from (??) and (??). \square

Definition 2.2. A function f on X is called a **(co-)closure function** if

1. f is extensive, $\forall x \in X, x \preceq f(x)$, ($\forall x \in X, f(x) \preceq x$),
2. f is monotone, $x_1 \preceq x_2 \implies f(x_1) \preceq f(x_2)$, for all $x_1, x_2 \in X$, and
3. f is idempotent, $f(f(x)) = f(x)$ for all $x \in X$.

Proposition 2.8. *If (f_*, f^*) is a Galois connection between (X, \preceq) and (Y, \leq) , then $f^* \circ f_*$ is a closure function for X and $f_* \circ f^*$ is a co-closure function for Y .*

Proof. Let $x \in X$. Then $x \preceq (f^* \circ f_*)(x)$ follows by (??).(??). Since both f^* and f_* are monotone we have, $x_1 \preceq x_2 \implies (f^* \circ f_*)(x_1) \preceq (f^* \circ f_*)(x_2)$. Thus $f^* \circ f_*$ is also monotone. By associativity of functions and (??).(??) we have

$$(f^* \circ f_*) \circ (f^* \circ f_*) = f^* \circ (f_* \circ f^* \circ f_*) = f^* \circ f_*$$

as needed. The dual statement is proved just as easily. \square

By (??), the closed elements of $f^* \circ f_*$ and $f_* \circ f^*$ are precisely the elements that are an image of some element under f^* , respectively f_* .

There is a one-to-one correspondence between Galois connections and (co-)closure functions.

Proposition 2.9. *If f is a closure (respectively co-closure) function, then there is a Galois connection (f_*, f^*) such that $f = f^* \circ f_*$ (respectively $f = f_* \circ f^*$).*

Proof. Let $f : X \rightarrow X$ be a closure over (X, \preceq) . Let \overline{X} be the set of closed elements of f that is $f(X) = \overline{X}$. We will construct a Galois connection between \overline{X} and X using (??). Let $f_* = f$, that is $f_* : X \rightarrow \overline{X}$ defined by $f_*(x) = f(x)$ for all $x \in X$. Let $f^* : \overline{X} \rightarrow X$ be the inclusion mapping, that is $f^*(x) = x$ for all $x \in \overline{X}$. Notice $f_* \circ f^*$ is the identity on \overline{X} and $f^* \circ f_* = f$.

1. Notice f_* is monotone since f is monotone and that f^* is monotone since the identity is monotone.
2. Let $x \in X$. Since f is extensive, we have $x \preceq f(x)$. Thus it follows,

$$x \preceq f(x) = (f^* \circ f_*)(x) = (f^* \circ f_*)(x).$$

3. Let $y \in \overline{X}$. There exists $x \in X$ such that $y = f(x)$. Since f is idempotent we have

$$f(y) = (f \circ f)(y) \preceq y$$

for all $y \in \overline{X}$ as needed.

Therefore, $(f_*, f^*) = (f, f^*)$ where $f^* : \overline{X} \rightarrow X$ is the inclusion mapping is a Galois connection between $(X, \prec 0$ and (\overline{X}, \preceq) . □

Proposition 2.10. *Let R be a relation between X and Y . Let*

$$f_R(A) = \{b \in Y : \forall a(a \in A \implies (a, b) \in R)\} \text{ and} \tag{5}$$

$$f^R(B) = \{a \in X : \forall b(b \in B \implies (a, b) \in R)\}. \tag{6}$$

Then (f_R, f^R) is a Galois connection between $(P(X), \subseteq)$ and $(P(Y), \supseteq)$

Proof. Clearly, $f_R : P(X) \rightarrow P(Y)$ and $f^R : P(Y) \rightarrow P(X)$ are functions. By (2) we must show

$$f_R(A) \supseteq B \iff A \subseteq f^R(B) \tag{7}$$

for all $A \in P(X)$ and all $B \in P(Y)$. Assume $B \subseteq f_R(A)$. We will show $A \subseteq f^R(B)$. Let $x \in A$. If $y \in B$, then $y \in f_R(A)$. Then, by (5), it follows $(x, y) \in R$. So we have shown, $y \in B \implies (x, y) \in R$ as needed to show $x \in f^R(B)$. Conversely, assume $A \subseteq f^R(B)$. We will show $B \subseteq f_R(A)$. Let $y \in B$. If $x \in A$, then $x \in f^R(B)$. Then, by (6), it follows $(x, y) \in R$. So we have shown, $x \in A \implies (x, y) \in R$ as needed to show $y \in f_R(A)$. Therefore, (7) holds. \square