

Bradley Johnson

ITAI 4374

Professor Patricia McManus

5 February 2026

Part 1: From Biological to Artificial Neural Networks

Biological neurons inspired the foundational design of artificial neural networks by demonstrating how complex behavior can emerge from simple processing units connected in large networks. In a biological neuron, dendrites receive incoming electrical signals, the cell body (soma) integrates those signals, and the axon transmits the output to other neurons through synapses. Artificial neurons mirror this structure by using input connections, weighted summation, an activation function, and an output signal.

Despite this inspiration, significant differences exist. Biological neurons operate using electrochemical signals, adapt continuously through synaptic plasticity, and function asynchronously. Artificial neurons rely on mathematical operations, predefined activation functions, and discrete training phases. Biological systems are extremely energy-efficient and adaptable but difficult to model precisely. Artificial neural networks are faster, scalable, and easier to optimize but lack the flexibility, robustness, and contextual understanding of biological intelligence. Each approach has strengths: biological neurons excel in adaptability and generalization, while artificial neurons excel in speed, precision, and large-scale computation.

FROM THE MAKERS OF WOLFRAM LANGUAGE AND MATHEMATICA

WolframAlpha

neuron anatomy

NATURAL LANGUAGE **MATH INPUT** **EXTENDED KEYBOARD** **EXAMPLES** **UPLOAD** **RANDOM**

Input interpretation:

neural cell (human)

Alternate names:

nerve cell | neuron | neurone

Functional description:

consists of a cell body, an axon, and dendrites and serves to receive, conduct, and transmit impulses in the nervous system

Hierarchy relationships: [More](#)

anatomical groups	neural cell somatic cell
structural groups	portion of neural tissue
regional parts	axon dendrite cell body of neuron
members	motor neuron set of neuroglial cells afferent neuron Purkinje neuron vestibular sensory cell ... (total: 24)
bilateral components	neural cell
developmental origin	ectodermal cell

[Definitions >](#)

Typical number in the human body:

100 billion

Typical physical characteristics:

shape | stellate

Constitutional parts:

protoplasm of neuron | plasma membrane of neuron | plasma membrane of neural cell

Spatial characteristics:

external structure | neurilemma

UMLS concept ID:

C0027882

Wikipedia page hits history: [Log scale](#)

(in hits per day)
(based on monthly averages of daily hits to the English-language "neural cell" page, including mobile traffic starting in 2016)

[Sources](#) [Download Page](#) **POWERED BY THE WOLFRAM LANGUAGE**

Biological Component / Characteristic	Biological Function	Artificial Neural Network Equivalent	Example in AI Architectures (CNNs, RNNs, etc.)
Cell body (Soma)	Integrates incoming signals and determines neuron activation	Processing unit / activation function	Neuron activation in dense layers of MLPs and CNNs
Dendrites	Receive signals from other neurons	Input connections / input nodes	Input feature maps in CNN convolution layers
Axon	Transmits output signal to other neurons	Output connection	Output passed to next layer in CNNs and RNNs
Synapses	Adjust signal strength between neurons	Weights	Learnable weights updated via backpropagation
Synaptic plasticity	Strengthening or weakening of connections over time	Weight updates during training	Gradient descent in CNNs and RNNs
Hierarchical organization	Information flows from simple to complex processing	Layered network structure	CNN feature hierarchy (edges → shapes → objects)
Temporal signal transmission	Processes information over time	Recurrent connections	RNNs and LSTMs for sequence modeling
Parallel processing	Multiple neurons fire simultaneously	Parallel computation	GPU-based training of deep networks
Specialized brain regions	Dedicated functional areas	Task-specific architectures	CNNs (vision), LSTMs (memory), Transformers (attention)

Part 2: Understanding the Brain's Architecture

The human brain is hierarchically organized, with lower-level regions handling basic sensory input and higher-level regions managing reasoning, memory, and decision-making. Specialized areas perform distinct tasks: the visual cortex processes visual information, the auditory cortex handles sound, and the hippocampus manages memory formation. This organization closely resembles modern AI architectures.

Convolutional Neural Networks (CNNs) parallel the visual cortex by processing information in layers, extracting edges, shapes, and increasingly complex features. Long Short-Term Memory (LSTM) networks mirror aspects of hippocampal function by maintaining and updating information over time, enabling learning from sequences. Attention mechanisms in transformer models resemble biological attention systems by prioritizing relevant information while filtering out noise.

The amygdala plays a key role in emotion processing, especially fear, risk evaluation, and threat detection. Insights from biological emotion processing can inform AI systems by improving decision-making under uncertainty, enhancing risk assessment, and strengthening pattern recognition in high-stakes environments. Emotion-inspired mechanisms may also enable adaptive learning by assigning priority to critical experiences, improving responsiveness and resilience in AI models.

FROM THE MAKERS OF WOLFRAM LANGUAGE AND MATHEMATICA

WolframAlpha

human brain anatomy

NATURAL LANGUAGE MATH INPUT

EXTENDED KEYBOARD EXAMPLES UPLOAD RANDOM

Examples for Brain Anatomy

Brain Anatomy

Get information about brain anatomy:

hypothalamus =

amygdala =

cerebellum =

Compare parts of brain anatomy:

lateral geniculate body and medial geniculate body =

primary motor cortex, primary somatosensory cortex =

Properties of Brain Anatomy

Find information about properties of brain anatomy:

signal of motor cortex goes to =

Brodmann area 4 associated cortical area =

neurons in hippocampus CA2 =

Amygdala function =

Brain area associated with mood =

Part 3: Integration Analysis and Future Implications

Understanding biological neural networks provides valuable guidance for improving artificial intelligence design. Biological systems demonstrate how intelligence can

emerge from massively parallel, adaptive, and energy-efficient architectures. These insights may inspire new AI models that move beyond rigid layer-based designs toward more dynamic, self-organizing systems. For example, biologically inspired learning rules such as Hebbian learning could complement or replace backpropagation, addressing limitations in scalability and biological plausibility.

A critical comparison reveals key trade-offs. Biological systems are remarkably energy-efficient, operating on approximately 20 watts, while modern AI models require vast computational resources. Biological learning is continuous, adaptive, and robust to noise, whereas artificial systems often require large labeled datasets and retraining. However, artificial systems excel in processing speed, precision, and the ability to scale across massive datasets. In terms of generalization, biological intelligence remains superior, adapting to novel situations with minimal prior exposure.

Future directions in AI research may involve neuromorphic computing, spiking neural networks, and hybrid systems that integrate biological principles with digital computation. Insights from emotion processing, attention, and memory could help overcome current AI limitations in reasoning, adaptability, and context awareness. Ultimately, deeper integration between biological understanding and artificial design may lead to more efficient, resilient, and human-like intelligent systems capable of learning and adapting in complex real-world environments.

References

Kandel, E. R., Koester, J. D., Mack, S. H., & Siegelbaum, S. A. (2021). *Principles of neural science* (6th ed.). McGraw-Hill.

Wolfram Alpha. Neuron anatomy. <https://www.wolframalpha.com>

Wolfram Alpha. Human brain anatomy. <https://www.wolframalpha.com>