

Comparative Evidence for the Correlated Evolution of Polyploidy and Self-compatibility in Solanaceae

Kelly Robertson¹, Emma E. Goldberg², and Boris Igić³

Dept. of Biological Sciences, 840 West Taylor St., M/C 067, University of Illinois at Chicago, Chicago, IL 60607.

¹ Email: <krober6@uic.edu>

² Email: <eeg@uic.edu>

³ Email: <boris@uic.edu>

Abstract

2 Breakdown of self-incompatibility occurs repeatedly in flowering plants with important
evolutionary consequences. In plant families where self-incompatibility is mediated by
4 S-RNases, previous evidence suggests that polyploidy may often directly cause
self-compatibility through the formation of diploid pollen grains. We use three approaches
6 to examine relationships between self-incompatibility and ploidy. First, we test whether
evolution of self-compatibility and polyploidy is correlated in the nightshade family
8 (Solanaceae), and find the expected close association between polyploidy and
self-compatibility. Second, we compare the rate of breakdown of self-incompatibility in the
10 absence of polyploidy against the rate of breakdown that arises as a byproduct of
polyploidization, and we find the former to be greater. Third, we apply a novel extension
12 to these methods to show that the relative magnitudes of the macroevolutionary pathways
leading to self-compatible polyploids are time-dependent. Over small time intervals, the
14 direct pathway from self-incompatible diploids is dominant, whereas the pathway through
self-compatible diploids prevails over longer time scales. This pathway analysis is broadly
16 applicable to models of character evolution in which sequential combinations of rates are
compared. Finally, given the strong evidence for both irreversibility of the loss of
18 self-incompatibility in the family and the significant association between self-compatibility
and polyploidy, we argue that ancient polyploidy is highly unlikely to have occurred within
20 the Solanaceae, contrary to previous claims based on genomic analyses.

Keywords: self-incompatibility, polyploidy, breeding systems, statistical phylogenetics,
22 comparative methods, angiosperms

INTRODUCTION

Self-incompatibility (SI) is the common ability of hermaphrodite plants to recognize and reject their own pollen with a genetically-based mechanism. Approximately one half of all extant angiosperm species prevent self-fertilization by deploying SI (Brewbaker 1959; Igić and Kohn 2006). Although many different mechanisms of SI evolved in flowering plants, a particular system, which uses RNases in the female component of self-rejection, appears to be ancient. It likely originated at least 90 million years ago, and it may cause SI in dozens of extant eudicot families, including some of the most diverse (Igić et al. 2008). A notable property of SI systems is the high rate at which they transition to self-compatibility (SC), considered one of the most common evolutionary transitions in plants (Stebbins 1974). Consequently, many individuals, populations, and species do not express SI. Empirical data on the mechanisms that underlie losses of SI are sparse. Although the genetic causes of transitions from SI to SC vary widely (Stone 2002), a significant proportion of such transitions may involve polyploidization (Livermore and Johnstone 1940; Crane and Lewis 1942; Stout and Chandler 1942; Lewis 1947; Brewbaker 1954; Pandey 1968; Charlesworth 1985; Chawla et al. 1997; Entani et al. 1999; Miller and Venable 2000).

Polyploidy is the duplication of an entire genome, resulting in three or more chromosome sets (Grant 1971). Polyploidy is thought to be common within angiosperms (20-89%; Stebbins 1950; Masterson 1994), although estimates vary widely depending on classification criteria and ability to separate recent from ancient duplication events.

Traditionally, estimates of polyploid frequency were calculated using the inferred chromosome number for a particular group (Stebbins 1938) or angiosperms as a whole (Grant 1963; Goldblatt 1980; Masterson 1994). However, polyploids may undergo genetic rearrangements, selective gene loss, and re-diploidization over time (Wolfe 2001), precluding cytological diagnoses of polyploid events over deeper evolutionary time scales (Byrne and Blanc 2006; Otto and Whitton 2000). The advent of genomic sequence data has enabled detection of multiple likely ancient polyploidization events in flowering plants (Ku et al. 2000; Blanc et al. 2000, 2003; Paterson et al. 2000; Vision et al. 2000; Simillion

et al. 2002; Bowers et al. 2003; Blanc and Wolfe 2004; Schlueter et al. 2004; Barker et al.

2008). In part as a result of these studies, it is widely held that nearly all angiosperms may have undergone a polyploid event in their evolutionary history (Soltis et al. 2009).

Sequence-based methods to detect ancient polyploidy generally estimate the ages of paralogous sequences, measured as the number of synonymous substitutions per site. A whole genome duplication event is expected to produce a detectable peak of sequence similarity approximating the time of the duplication event (Lynch and Conery 2000, 2003).

Other methods infer genome duplications from comparative study of microsynteny. Such studies use long sequence reads to seek out duplicated regions with identical or similar gene order (e.g., Vision et al. 2000). The association between polyploidy and the radiation of early angiosperm lineages may suggest a causal positive association between polyploidy and diversification rate (reviewed in Soltis et al. 2009; but see Meyers and Levin 2006).

The causes may be related to the number of genotypic and phenotypic changes associated with polyploidy, including changes in morphology, phenology and life-history characteristics (Levin 1983; Ramsey and Schemske 2002), wider ecological tolerances—perhaps leading to an increase in range size (Grant 1971; Levin 1983; Hijmans et al. 2007)—as well as the formation of new gene complexes (Wendel 2000). Because of their far-reaching effects, we are most interested in alterations to the breeding and mating system, specifically, the increased propensity of polyploids to self-fertilize (Lewis 1943, 1947; Brewbaker 1953, 1954, 1958; Pandey 1968; de Nettancourt et al. 1974; Miller and Venable 2000; Barringer 2007; Husband et al. 2008).

The evolutionary link between mating system and ploidy was proposed and empirically investigated at least since Stebbins (1950, 1957) and Grant (1956), who found a strong association between polyploidy and self-fertilization. Phylogenetic comparative studies also find some evidence for the correlated evolution of selfing and ploidy (Barringer 2007; Husband et al. 2008). Theoretical models predict that this association principally depends on the ability of polyploidy to ameliorate the expression of inbreeding depression, as well as the effect of mating system on the rate of polyploid formation and establishment (reviewed

in Ramsey and Schemske 1998; Barringer 2007; Husband et al. 2008). A positive
80 correlation may, however, result directly from the genome duplication process itself, at
least in families with gametophytic SI (Livermore and Johnstone 1940; Crane and Lewis
82 1942; Stout and Chandler 1942; Lewis 1947; Brewbaker 1954; Pandey 1968). Evidence
from both natural and experimentally induced tetraploids from several families, including
84 Rosaceae, Fabaceae, Onagraceae, and Solanaceae, suggests that polyploidy almost
invariably disrupts GSI (reviewed in Ramsey and Schemske 1998; Stone 2002). Lewis
86 (1947) observed that the diploid pollen heterozygous at the S-locus, often termed
“heteroallelic,” was not degraded, whereas the diploid pollen homozygous at the S-locus
88 was rejected via a normal SI response. A suite of observational and experimental data
subsequently led to the modification and generation of several models, which aim to
90 explain the molecular physiological mechanism of RNase-based SI (RSI; reviewed in Hua
et al. 2008; McClure 2009). The tendency of polyploidization to break down SI is so well
92 established that the models are constructed specifically around the ability to explain it as a
byproduct of the proposed mechanism McClure (2009).

84 However, a broad comparative study did not find evidence for a correlation between
ploidy and SI among all SI systems (Mable 2004). The relationship was significant within
96 groups with gametophytic SI, in which the pollen phenotype is determined by its own
haplotype (RSI is one subtype of gametophytic SI systems). It appears that groups with
98 sporophytic SI, in which the pollen phenotype is determined by the maternal genotype,
show no such pattern and reduce the power of the combined analysis. In a separate study,
100 Miller and Venable (2000) found strong support for a similar pattern of association within
gametophytic SI in a study of the evolution of gender dimorphism. Nevertheless, although
102 they are rare, exceptions exist. In the Rosaceae family, where RSI is common and generally
appears associated with diploid species, at least three tetraploid species, *Prunus cerasus*,
104 *P. spinosa* and *P. fruticosa*, express a functional SI response in heteroallelic pollen (Hauck
et al. 2002, 2006; Nunes et al. 2006; Huang et al. 2008; Vieira et al. 2008).

106 Here, we conduct combined analyses of ploidy and breeding system character

evolution, aimed at detecting whether the two characters evolve in a correlated manner.

108 Our study relies on a large comparative dataset to detect the strength of this association
within the Solanaceae. We infer the relative rates of breakdown of SI due to
110 polyploidization and other mechanisms, which do not involve genome duplication. Because
polyploids can arise on the background of SC or SI, we also compare the magnitudes of
112 transitions leading to polyploidy. In the process, we propose and use a novel general
method to quantify the contributions of pathways of sequential character state changes
114 involved in generating a strong association between polyploidy and SC.

METHODS

Study System

116 Solanaceae, the nightshade family, contains approximately 2,700 species (Olmstead and
Bohs 2007), many of which are of considerable agricultural importance (e.g., tomato,
potato, eggplant, tobacco, petunia). Partly due to its economic value, the group enjoys a
120 rich tradition in cytogenetic and breeding system studies. There is a wealth of available
karyotype data, and the family has long been a model system for study of the genetic basis
122 of SI (East and Mangelsdorf 1925). This group is presently the subject of a large
collaborative effort aimed at gaining detailed understanding of phylogenetic relationships
124 among its species (Olmstead and Bohs 2007). In all of the analyses presented below, we
rely extensively on such sources of previously generated data.

Data Collection and Character Coding

Self-incompatibility

Taxa were scored as SC or SI based on papers found through extensive searches of the
primary literature, and as reported in Igić et al. (2006). We searched the online databases
*ISI Web of Knowledge Science Citation Index*TM and the *Google Scholar*
(<http://www.google.com/schhp>) using dozens of search terms related to plant breeding

systems and names of taxa. Our search included papers published through 2009. In addition, we used a personal library of reprints (B.I.) with books, monographs, and manuscripts about Solanaceae species, dating from about 1850 through 1995. A vast proportion of designations was made by the authors of the original studies. Occasionally, the authors did not designate a taxon as SC or SI but provided sufficient information for the calculation of the index of self-incompatibility (ISI), a measure of the relative success of manually performed self- and cross-pollinations (Lloyd 1965; Bawa 1974). We employ the relative number of fruits per flower after each pollination treatment as a metric of fertilization success. The ratio of fruit set after self- versus cross-fertilization is subtracted from unity, resulting in a continuous index of values that encompasses all possible strengths of SI. Thus, the metric of ISI is calculated as

$$ISI = 1 - \frac{\text{relative selfed success}}{\text{relative outcrossed success}}.$$

The upper bound is unity (complete SI), but when self-pollinations result in higher fruit set than cross-pollinations, it is possible to obtain negative ISI values. Historically, species with ISI values above 0.8 have been classified as SI (Bawa 1974). Because of the relative dearth of species with intermediate values of ISI, classification is largely insensitive to the exact cut-off value in most angiosperm families studied to date (A. Raduski and B. I., unpublished data).

Ploidy

Taxa were scored as diploid (D) or polyploid (P) based on reports from both haploid and somatic chromosome counts in the Index to Plant Chromosome Number (<http://mobot.mobot.org/W3T/Search/ipcn.html>) and the Royal Botanical Gardens, Kew (<http://www.kew.org>) online databases as well as the primary literature (see above). The scoring criterion was based on the inferred basal chromosome number for the taxonomic group in question.

Members of the subfamily Solanoideae, the tribe Anthocercideae, and the genus

Nicotiana form a monophyletic group with the base chromosome number $x=12$ (Olmstead and Palmer 1992; Olmstead and Sweere 1994; Olmstead et al. 2008). Within the Solanoideae, species in *Capsicum*, *Chamaesaracha*, *Datura*, *Dunalia*, *Dyssochroma*, *Grabowskia*, *Iochroma*, *Jaltomata*, *Lycianthes*, *Lycium*, *Mandragora*, *Nicandra*, *Nolana*, *Nothocestrum*, *Phrodus*, *Physalis*, *Salpichroa*, *Solandra*, *Solanum*, *Vassobia*, *Withania*, and *Witheringia* were scored as diploid when the somatic chromosome number ranged from $2n=16-24$, and polyploid when the somatic chromosome numbers ranged from $2n=34-96$. Also part of the Solanoideae subfamily is the tribe Hyoscyameae, containing the genera *Anisodus*, *Atropa*, *Atropanthe*, *Hyoscyamus*, *Physochlaina*, *Przewalskia*, and *Scopolia*. The genera *Anisodus* and *Atropa* are reported as having a base chromosome number of $x=6$ (Tu et al. 2005) or $x=12$ (Olmstead and Palmer 1992; Badr et al. 1997; Olmstead et al. 1999). We encoded all *Anisodus* and *Atropa* species as polyploid because they had somatic chromosome numbers in excess of $2n=48$. The genus *Hyoscyamus* has a base chromosome number $x=14,17$ (Tu et al. 2005). Species with chromosome counts $2n=28, 34$ were scored as diploid, while the species *H. albus* and *H. pusilus* with somatic chromosome counts equal to $2n=68$ (Goldblatt and Johnson 1979), were scored as polyploid. The remaining genera in the Hyoscyameae, *Atropanthe*, *Physochlaina*, *Przewalskia*, and *Scopolia* have somatic chromosome counts ranging from $2n=42-48$ (Tu et al. 2005), and were therefore encoded as polyploid. Within the tribe Anthocercideae, species in the genera *Crenidum*, *Duboisia*, *Grammosolen*, and *Symonanthus* have somatic chromosome numbers ranging from $2n=60-102$ (Goldblatt and Johnson 1979; Darlington and Wylie 1956), and were encoded as polyploid. The genus *Nicotiana* forms a dysploid series with basal chromosome numbers ranging from $x=7-12$. Species in this genus were encoded as diploid or polyploid based on Goodspeed (1954).

In the tribe Petunieae, *Hunzikera* and *Nierembergia* have somatic chromosome numbers $2n=32$ and $2n=36$, respectively (Acosta et al. 2006). We encoded these species as polyploid because other genera in this group have low haploid chromosome numbers; $n=7-11$ in *Petunia*, *Bouchetia*, *Calibrachoa*, *Fabiana*, *Leptoglossis*, and *Brunfelsia* (Goldblatt and

Johnson 1979).

In the subfamily Cestroideae, species in the genera *Browalia*, *Cestrum*, *Salpiglossis*, *Sesseeae*, *Streptosolen*, and *Vestia* were scored as diploid when somatic chromosome numbers ranged from $2n=16-22$ (Olmstead et al. 2008). The only species scored as a polyploid in this group, *Browalia speciosa*, has a somatic chromosome number $2n=44$ (Welsh and Sink 1981). In tribe Benthamiellieae, which is likely sister to the Cestroideae (Olmstead et al. 2008), species in the genera *Benthamiella*, *Combera*, and *Pantacantha* have somatic chromosome counts $2n=22$ (Goldblatt and Johnson 1979) and were scored as diploids. No polyploids were found within Benthamiellieae.

The remaining genera with available ploidy information are *Schizanthus* and *Schwenckia*. *Schwenckia* is part of a polytomy with the “ $x=12$ ” clade (containing the subfamily Solanoideae and the tribe Anthocercideae), the subfamily Cestroideae, and the Petunieae tribe (Olmstead et al. 2008). *Schwenckia browallioides* has a somatic chromosome number $2n=24$ (Goldblatt and Johnson 1979), and was scored as a diploid. *Schizanthus* forms a sister group to the rest of Solanaceae. *Schizanthus grahamii* and *Schizanthus pinnatus* have somatic chromosome counts of $2n=20$ (Gaiser 1926; Goldblatt and Johnson 1979), and were both scored as diploids.

Conflicting Report

We closely tracked potential errors in reports for the breeding system and ploidy character states. Out of hundreds of studies, we found that the information from a single study is consistently in conflict with others. Marks (1965) lists several species as SI in *Solanum* section *Petota*. Specifically, the polyploid *S. agrimoniifolium* Rybd., *S. colombianum* Dun. (syn. *S. moscopanum* Hawkes), *S. oxycarpum* Schiede, *S. stoloniferum* Schltdl. (syn. *S. polytrichon* Rybd), *S. iopetalum* (Bitt.) Hawkes, *S. guerreroense* Corr., and *S. hougasii* Corr. are listed as SI. Instead, these are listed as SC in a minimum of two independent sources (see Table S1 for complete listing of SI-SC and D-P character state data for taxa used in this study). We omitted all data from Marks (1965) because the criterion used for

SI determination was failure of fruit set after hand self-pollination, yet these plants
198 autonomously set fruit in the greenhouse with high seed set. While it is possible, and
sometimes observed, that SI species set selfed fruit, it is typically seedless (parthenocarpic),
200 which is inconsistent with the data obtained by Marks. Other reports not included in this
study are from the sterile triploids *Solanum commersonii*, *S. cardiophyllum*, and *S. maglia*
202 (Hawkes and Hjerting 1969; Correll 1962).

Polymorphisms

204 It is widely held that ploidy and breeding system frequently transition between states,
mostly unidirectionally (Stebbins 1974; Meyers and Levin 2006). This is presumably
206 because the mutation rates that generate polyploids and SC, as well as the conditions for
fixation of these mutations in plant lineages (evolutionary transitions), are not limiting.
208 The high magnitude and directionality of the two processes are thought to result in the
commonly observed pattern of occurrence of rare SC individuals in otherwise SI
210 populations (Rick and Tanksley 1981; Rick 1986; Ando et al. 1998; Bohs 2000), and
polyploid individuals in otherwise diploid populations (Lewis and Suda 1976; Halverson
212 et al. 2008). Thus, a major difficulty in determining the correlation of ploidy and SI/SC
state is that these traits are rarely measured in the same individual or population. As our
214 analysis is designed to determine the correlation of character state transitions at the
species level, not the frequency (and correlation) of the initial mutations, we systematically
216 eliminated polymorphisms for data with phylogenetic information. SI species in which rare
SC individuals are found, without strong evidence for local fixation, were encoded as SI
218 (*Capsicum pubescens*, *Nicotiana glauca*, *N. langsdorfii*, *Petunia axillaris*, *P. reitzii*,
Solanum arcanum, *S. chilense*, *S. habrochaites*, *S. pennellii*, *S. peruvianum*,
220 *S. sisymbriifolium*, and *Witheringia solanacea*). Diploid species in which rare polyploid
individuals are found were encoded as diploid (*Solanum campanulatum*, *S. cinereum*,
222 *S. torvum*, *S. verrucosum* and *S. polyadenium*). *Solanum tuberosum*, *S. bulbocastanum* and
L. californicum are polymorphic for both compatibility and ploidy. When the traits are

measured in the same individual, SI segregates with the diploid cytotype and SC with the polyploid cytotype [e.g., *Solanum tuberosum*: SI-D (Kirch et al. 1989) and SC-P (Lewis 1943), *Solanum bulbocastanum*: SI-D and SC-P (Hermesen and Boer 1971), *Lycium californicum*: SI-D and SC-P (Yeung et al. 2005)]. We encoded these species as SI-D in our analysis, although the association of SC with the polyploid cytotype is itself suggestive of a causal relationship between SC and polyploidy.

Non-phylogenetic Analyses of Correlated Character Evolution

We used a χ^2 -test of association to investigate whether there is a relationship between ploidy and compatibility. We conducted association tests in both a 2×2 table, for which polymorphic species were collapsed into the SI and diploid category as described above, and in a 3×3 table, where they were encoded as a separate category. In the cases where the expected value for any cell was less than five, we applied the Yates' correction for continuity (Yates 1934). An assumption of the χ^2 test is that the data are independent, which is violated when species share ancestry (Felsenstein 1985; Pagel 1994). We use phylogenetic tests (described below) to allow for species relationships.

Reconstruction of Phylogenetic Relationships

We constructed a composite phylogenetic hypothesis from a skeletal family phylogeny (Olmstead et al. 2008) augmented by insertion of clades obtained from eighteen fine-scale molecular phylogenetic studies (Levin et al. 2006; Weese and Bohs 2007; Bohs 2007; Prohens et al. 2006; Spooner et al. 1993; Peralta and Spooner 2001; Spooner et al. 1992; Mione et al. 1994; Walsh and Hoot 2001; Smith and Baum 2006; Whitson and Manos 2005; Mace et al. 1999; Levin and Miller 2005; Miller et al. 2008; Clarkson et al. 2004; Ando et al. 2005; Montero-Castro et al. 2006; Perez et al. 2006). We imposed the branching order found in strict consensus trees of the individual data sets listed above. Polytomies were randomly resolved to allow analyses that require bifurcating branching events.

Branch lengths proportional to time are not currently available from systematic studies

of the family, and we consequently relied on two standard methods for assigning them. First, we set all branch lengths to 1, except for those introduced by resolving polytomies, which were set to 10^{-6} ; we refer to this as the “unit” branch length assumption. Second, we used branch lengths computed from node depth with the algorithm described by Grafen (1989), with a root-to-tip time of one and a scaling power of $\rho = 1$. Neither of these methods is an entirely valid substitute for true branch lengths but, when paired, they stand in for a wide range of possibilities. Comparing results from the unit and Grafen assumptions is likely to give a strong indication of how robust our conclusions are to the lack of true branch lengths.

Phylogenetic Comparative Tests

Correlated Evolution of Self-incompatibility and Polyploidy

Pagel’s (1994) method detects correlated evolution between two discrete characters by comparing the likelihood of a character-dependent model to a model that allows characters to evolve independently. If the characters are evolving independently, the transition rates between states of one character are independent of the state of the other character. For example, the gain of polyploidy would occur at the same rate, irrespective of whether the species is SI or SC ($q_{12} = q_{34}$; Fig. 1). The model of independent evolution will then have four parameters estimated ($q_{12} = q_{34}$, $q_{21} = q_{43}$, $q_{13} = q_{24}$, $q_{31} = q_{42}$), and it is compared to an unconstrained model of correlated or dependent evolution where all eight rate parameters are estimated.

We fit parameter values in these two models using the Markov chain Monte Carlo method implemented in *BayesTraits Discrete*. (The settings detailed here were also used for the *Multistate* analysis described below.) We used an exponential hyperprior that was seeded uniformly between 0 and 100. The rate deviation was set to 0.1 – 0.3 (unit branch lengths) or 4 – 8 (Grafen branch lengths) to achieve the recommended acceptance rates of 20 – 40%. The “Pis” option, describing the prior state probabilities at the root, was set to “none” so that the likelihoods of each state at the root were added together. This setting

yields the same ratio of likelihoods between different sets of rates as does the “uniform”
weighting, and it provides the correct likelihood when only one root state is logically
possible in the *Multistate* analysis. After a 50,000 generation burn-in period, the chain was
sampled every 1,000 generations over a total of 505,000 generations.

The weight of evidence used to evaluate the relative fit of the independent and
correlated models was calculated with Bayes factors (BF ; Kass and Raftery 1995), with
the marginal likelihood of each model approximated by the harmonic mean of the
likelihoods in the Markov chain, as recommended by the *BayesTraits* manual. For the test
statistic $2\ln(BF)$, with the dependent model favored, a value between 2 and 5 provides
“positive” evidence for correlated evolution, and a value of more than 10 provides “very
strong” evidence (Raftery 1996, p. 165; Pagel and Meade 2006).

Transitions to Self-Compatibility

The rate parameters describing simultaneous dual transitions (between states 1 and 4, and
between 2 and 3) are ordinarily set to zero because the probability of two separate events
in a single instant is negligible (Pagel 1994). However, dual transitions are biologically
realistic in this system because polyploidy directly breaks down incompatibility through the
formation of diploid pollen grains. To allow dual transitions, we modeled the evolution of a
single three-state character composed of compatibility and ploidy: SI-D = state 1, SC-D =
state 3, and SC-P = state 4 (see Fig. 1). There were no observations of SI-P (state 2). The
Multistate model was used to compare the magnitudes of the transition rate parameters to
determine the dominant means by which SI is lost and by which polyploidy arises.

We initially prohibit transitions from SC to SI when implementing the model because
previous work on the evolution of SI demonstrates that the rates of reversal to SI from SC
are negligible and not significantly different than zero (Igić et al. 2006). In Solanaceae,
strong negative frequency-dependent selection yields dozens of alleles at the SI locus
(S-locus). Every SI species of Solanaceae studied so far exhibits evidence of such ancient
S-locus polymorphism, with coalescent times estimated at approximately 40-50 million

years old (Ioerger et al. 1990; Igić et al. 2006; Paape et al. 2008). This feature of the S-locus provides powerful evidence for the continuous long-term persistence of SI since the common ancestor of all extant species display shared S-locus polymorphism until present day (Igić et al. 2006). We incorporated these data in our model by setting the reversal transition rates, q_{31} and q_{41} to zero, but we also present results where reversals are allowed, for comparison. We also initially prohibit transitions from polyploidy to diploidy because polyploidy has long been thought to be a character whose evolution is exceptionally asymmetrical (Stebbins 1971, 1980; Bull and Charnov 1985; Meyers and Levin 2006). With the advent of genomic and phylogenetic methods to infer ancient polyploidization, numerous studies have found evidence for polyploid events within angiosperm lineages (reviewed in Soltis et al. 2009). However, within the Solanaceae, genomic evidence indicates there has been no duplication since the family diverged from the Rubiaceae (Lin et al. 2005; Wu et al. 2006) ca. 85 mya (Wikstrom et al. 2001). To implement irreversibility of polyploidization, we set the rates q_{43} and q_{41} to zero, but we again present results where reversals are allowed, for comparison.

Loss of SI can occur in diploid or polyploid lineages. In order to compare the frequencies of these two events, we used the *Multistate* model to compare the rate of SI loss in diploids (q_{13}), presumably occurring by mutations in genes that regulate or encode for components of SI pathway, to the rate of SI loss effected directly by polyploidization (q_{14}). The posterior distribution of the difference between these two rates allows a simple assessment of the magnitude of their difference.

Transitions to Polyploidy

Under the assumptions of irreversibility of self-compatibility and polyploidy, the root of the tree must have been in state SI-D. Comparing the magnitudes of the different means by which polyploidy can be reached from that ancestral SI-D state is more complicated than directly comparing the transition rates q_{14} and q_{34} from the *Multistate* model because a state 1-to-3 transition must precede a state 3-to-4 transition. We therefore developed a

method to compare the transition probabilities of the two routes to self-compatible
 polyploidy: pathway 14 is polyploidization from self-incompatibility (SI-D directly to
 SC-P), and pathway 134 is loss of self-incompatibility followed by polyploidization from
 self-compatibility (SI-D to SC-D to SC-P). Let $W_{14}(t)$ and $W_{134}(t)$ be the probabilities of
 transitions from SI-D to SC-P via pathway 14 and pathway 134 , respectively, after time t
 has elapsed. Derivation and explicit forms of $W_{14}(t)$ and $W_{134}(t)$ are shown in the
 Appendix. From the perspective of an extant SC-P species, the values of $W_{14}(t)$ and
 $W_{134}(t)$ reveal which pathway was more likely as a function of an SI-D ancestor's age, t ,
 which could range from very recent to the age of the tree.

Provided that none of the forward rates are zero, for very small elapsed times, pathway
 14 will be more likely than pathway 134 because only a single event is needed. For very
 large elapsed times and assuming irreversibility, the ratio of pathway 14 to 134
 probabilities is q_{14}/q_{13} because it is the first step away from SI-D that determines the
 pathway taken. At intermediate times, the relative strengths of the two pathways may
 change with the time available for transitions to take place, depending on the rate values.
 In order to compare the relative importance of pathways taken to SC-P, we computed,
 across the posterior rate distribution, the two pathway probabilities and their differences
 for time intervals ranging from 0 at the tips to the maximum, root depth of the tree.

An alternative to our pathway analysis might be to use stochastic character mapping
 to infer the sequence of states traversed as a lineage works its way from SI-D to SC-P. We
 prefer our approach because the pathway probabilities can be computed analytically and
 because it is not subject to the additional layer of model uncertainty added by the choice
 of ancestral state reconstruction method (Pagel 1999).

RESULTS

We collected ploidy information for 917 species, of which 75% are diploid, 20% polyploid,
 and 5% polymorphic. The distribution of haploid chromosome numbers is given in Fig. 2.
 The proportion of polyploids observed here is similar to a previous estimate for herbaceous

dicots (26.2%; Otto and Whitton 2000). Polyploid species are found in all genera sampled except *Petunia*, *Jaltomata*, *Datura*, and *Calibrachoa*. They are especially common in *Solanum* section *Etuberosum* and *Nicotiana* section *Suavolentes*.

We collected breeding system information for 550 species, of which 56% are SC, 35% SI, 4% are polymorphic for SI and SC, and 5% are dioecious. SI is spread throughout the family, including *Solanum*, *Nicotiana*, and *Physalis*, whereas *Jaltomata*, *Capsicum*, and *Datura* are primarily SC and *Petunia* and *Calibrachoa* are primarily SI. The distribution of ISI for Solanaceae species with sufficient data to calculate ISI is given in Fig. 3. The bimodal distribution suggests that, despite some exceptions, classification of breeding system as binary character appears a reasonable approximation of the continuous empirical distribution of ISI.

Character data for both self-(in)compatibility and ploidy is known for 408 species; 40.0% are SI-D, 0.3% are SI-P, 44.0% are SC-D, and 15.7% are SC-P (see Table 1). The single instance of SI-P is an allopolyploid hybrid of *Solanum oplocense* Hawkes x *Solanum gourlayii* Hawkes (Camadro and Peloquin 1981). The number of SI-P is significantly under-represented ($\chi^2 = 48.6$, $df = 1$; $p < 0.01$). The proportion of SI-P remains significantly underrepresented when polymorphisms are encoded as a separate category ($\chi^2 = 65.9$, $df = 4$; $p < 0.01$) in a 3×3 table (not shown).

Phylogenetic Distribution of Character States

Of approximately 98 genera and 2716 species described in the family (Bohs 2007), our phylogenetic analyses were based on the dataset containing 19 genera and 266 species for which character states and phylogenetic placement were available. Species in the genera *Solanum* and *Nicotiana* are relatively well sampled (124/1328, 48/108), whereas sampling in the genus *Lycianthes* (1/200) is poor, and *Cestrum* (0/175) is altogether absent. Of the 266 species included in analyses, 38.4% are SI-D, 0% SI-P, 44.7% SC-D, and 16.9% SC-P. These proportions are not significantly different from the larger set of species used in the non-phylogenetic analysis ($\chi^2 = 0.8$; $df = 1$; $p = 0.4$). The association between SC and P

remains significant for the subset of species placed on our phylogenetic tree ($\chi^2 = 33.0$;
 386 $df = 1$; $p < 0.01$).

Phylogenetic Comparative Tests

Correlated Evolution of Self-incompatibility and Polyploidy

For the *Discrete* character analysis on the unit branch length tree, the *log* of the harmonic
 390 mean of the likelihoods was -225.4 for the independent model with four estimated rates
 and -206.2 for the dependent model with eight rates. Comparing the two models therefore
 392 gives $2 \ln(BF) = 38.4$, providing very strong support for correlated evolution between SC
 and polyploidy within Solanaceae. On the Grafen branch length tree, the log harmonic
 394 means were -231.4 and -216.8 for the independent and dependent models, respectively,
 giving $2 \ln(BF) = 29.2$ and, again, very strong support for correlated evolution.

Transitions to Self-Compatibility

The *Multistate* analyses performed to reveal how often polyploidy breaks down SI find that
 398 the loss of SI occurs more often through mutations in diploids than through
 polyploidization (Table 2). This conclusion is robust to the branch length assumption and
 400 to the irreversibility assumption ($q_{13} > q_{14}$ with probability 1.0 for the restricted model
 with either type of branch lengths; for the reversible model that probability is 0.994 with
 402 unit branch lengths and 0.993 with Grafen branch lengths).

Transitions to Polyploidy

Using the unit branch length tree, the probability of the single step pathway to SC
 404 polyploids, $W_{14}(t)$, is greater over time intervals $t < 10.5$, and the two step pathway
 probability, $W_{134}(t)$, is greater for $t > 10.5$ up to the maximum depth of 17 (Fig. 5A). Both
 406 pathways therefore contribute to the evolution of SC polyploids over the time spanned by
 this tree. To assess the significance of the difference between the two pathway probabilities,
 408 we computed $W_{14}(t) - W_{134}(t)$ across the posterior rate distribution for values of time

ranging from 0 to 17. Fig. 6A shows probability contours that illustrate the probability associated with pathway 14 or pathway 134 being more likely on the unit branch length tree. For example, at $t = 5.7$ there is a 90% chance (or greater, for $t < 5.7$) that the evolution of SC polyploidy is more likely to occur through pathway 14. As elapsed time increases, pathway 14 becomes less and less likely until at the maximum node depth of the tree, $t = 17$, there is a 75% chance that SC polyploidy is more likely to occur through pathway 134.

Using instead the Grafen branch lengths affects the inference of the relative strength of the pathways leading to the evolution of SC polyploids. In this tree, the single step pathway 14 dominates initially, but only for very short elapsed times ($t < 0.1$; Fig. 5B). Fig. 6B illustrates the probability associated with pathway 14 or pathway 134 being more likely. For example, at $t \leq 0.04$ there is a 90% chance that the evolution of SC polyploidy is more likely to occur through pathway 14. At an elapsed time of $t = 0.09$, both pathways contribute equally to the evolution of SC polyploidy. However for the majority of time across the Grafen branch length tree ($t = 0.17$ to 1), pathway 134 is the dominant route leading to SC polyploidy.

DISCUSSION

Phylogenetic Comparative Tests

A strong association exists between polyploidy and SC in Solanaceae. While it is perhaps not surprising, given the mechanism of SI breakdown upon polyploidization within RSI systems (Livermore and Johnstone 1940; Stout and Chandler 1942; Pandey 1968; Chawla et al. 1997; Entani et al. 1999), our result provides strong comparative evidence from this family for a causal association, and it therefore predicts similar patterns for all angiosperm groups that share a common genetic basis for SI. In an earlier broad comparative study of the relationship between compatibility and ploidy, Mable (2004) did not find significant evidence for correlated evolution within families with different SI systems, despite recognizing a significant trend within RSI families. The absence of a correlation in Mable's

(2004) broader study suggests that a causal relationship between SC and polyploidy may
 438 be driving the association within RSI families, but that in general, SC may not be a strict
 requirement for polyploid establishment. Examining the rate parameters leading to SC
 440 and/or polyploidy in other families with non-RSI systems could determine to what extent
 mate limitation and inbreeding depression contribute to an association between SC and
 442 polyploidy. If there is strong selection within newly established polyploids for SC due to
 mate limitation, the rate from SI-P to SC-P is expected to be larger than the rate from
 444 SI-D to SC-D. Conversely, if the alleviation of inbreeding depression is the primary factor
 leading to a correlation between SC and P, the rate from SC-D to SC-P is expected to be
 446 larger than the rate from SI-D to SI-P.

Statistical phylogenetic methods measure lineage transition rates, which depend on the
 448 availability of individual transitions (the mutation rate) and selective processes that act to
 fix these mutations within populations. Our analysis shows that SC lineages arise more
 450 often within diploids than as a byproduct of polyploidization ($q_{13} > q_{14}$), which could be a
 result of differences in the mutational opportunity for character change or selection acting
 452 on those mutations. Generally, estimates from mutational studies suggest that the
 mutation rate of SI breakdown within diploids is approximately equal to or lower than the
 454 rate of polyploidization; this suggests that selection, not mutational opportunity, may be
 the primary factor causing our observed rate difference. Estimates of autopolyploid
 456 formation are on the order of the genic mutation rate ($\mu_P = 3 \times 10^{-5}$) (Ramsey and
 Schemske 1998). The mutation rate to SC has been previously inferred from incompatible
 458 pollen tube growth studies (Lewis 1979). These studies often cannot separate the
 mutations that arise with an increase in ploidy from those that arise without an increase in
 460 ploidy (de Nettancourt 1977). We use a conservative assumption, that none of the
 breakdowns of SI detected in these studies are due to polyploid pollen grains. The
 462 estimated rate of breakdown of SI within diploids in *Oenothera organensis*, *Prunus avium*,
Trifolium repens, *T. pratense*, *Nicotiana glauca* and *Petunia sp.* ranges from
 464 $\mu_{SC} = 0.02 \times 10^{-5}$ to 1×10^{-5} per pollen grain (Lewis 1979; de Nettancourt 1977). Given

that an average number of pollen grains per flower is ca. 5×10^5 within a SI diploid, the
 466 number of SC diploid gametes is approximately comparable to the number of SC polyploid
 gametes ($5 \times 10^5 \times \mu_{SC} = 0.1$ to 5 for SC diploid pollen, and $5 \times 10^5 \times \mu_P = 15$ for SC
 468 polyploid pollen). Although both measures are associated with a high estimation error,
 taken together with the transition rate estimates they suggest a remarkably clear result: a
 470 more severe selection pressure restricts the ability of newly established polyploids to fix
 within populations (Ramsey and Schemske 2002; Levin and Miller 2005).

Because polyploidization of diploid GSI individuals almost invariably causes a direct
 472 transition to SC in one step (SC-D to SI-P), use of the common statistical phylogenetic
 models for measuring correlated evolution of discrete characters (*BayesTraits Discrete*)
 474 would be inappropriate (Pagel 1994). Consequently, we use a widely employed alternative
 evolutionary model (*BayesTraits Multistate*), which allows appropriate single-step
 476 transitions (Fig. 1). We also develop a new extension that enables assessment of the
 relative contribution of multiple rates to the evolution of self-compatible polyploidy. Our
 478 results on the unit branch length tree provide considerable support for the direct pathway
 from SI-D to SC-P, thought to precede the evolution of gender dimorphism in the genus
 480 *Lycium*, and many others (Miller and Venable 2000). At the same time, it appears that
 Brunet and Liston (2001) expressed a valid concern when they highlighted the importance
 482 of comparing the relative magnitude of the possible pathways to SC-P, the inferred
 stepping stone to dioecy. In fact, our results employing the Grafen branch lengths, likely to
 484 bear more resemblance to the true tree than the unit branch length assumption, show
 much greater support for the two-step pathway, from SI-D to SC-D to SC-P, over all but
 486 the shortest time intervals. An extension of the approach we use to compare sequential rate
 pathways could also be employed to directly assess their relative contribution in the
 488 evolution of gender dimorphism in *Lycium*, as well as other taxa.

Our *Multistate* model omits SI-P species because they do not occur in the phylogenetic
 490 dataset. In the full dataset, including those with no certain placement on the phylogeny,
 we found the description of one likely occurrence of SI-P. Camadro and Peloquin (1981)
 492

describe an allopolyploid hybrid, *Solanum oplocense* Hawkes x *Solanum gourlayii* Hawkes,
494 whose expression of self-(in)compatibility depends on which parent was the pollen or ovule
donor. Interestingly, it is possible that this exception may not break the genetic rule. If
496 one of the parent species of the allopolyploid hybrid were SI and another SC, with the
S-locus sufficiently degraded, the hybrid may have expressed only one functional allele of
498 the S-locus (J. Kohn, pers. comm.). Non-haploid pollen grains homozygous at the S-locus
may be rejected via a normal SI response Lewis (1947) .

500 Despite the extremely rare occurrence of SI-P in our collection of data from
Solanaceae, a few well-studied examples are known from another family with RSI.
502 Self-incompatible Rosaceae express a system that is homologous to one found in Solanaceae
(Igić and Kohn 2001), and members of the family show a strong association between SC
504 and polyploidy (Dickinson et al. 2007). However, the genus *Prunus* presents noted
exceptions. While *Prunus pseudocerasus* (Huang et al. 2008) shows the expected SI
506 breakdown in polyploids, the likely recent allopolyploids *P. cerasus* (Hauck et al. 2002),
P. spinosa (Nunes et al. 2006) and *P. fruticosa* (Pruski 2007) each retain functioning SI.
508 Alterations in the pollen-S gene of *Prunus cerasus* are suggested to be responsible for the
maintenance of SI in polyploids (Hauck et al. 2006). The source of evidence in support of
510 this hypothesis is the differential SI response to deletion of the pollen-S gene in *Prunus* and
the Solanaceae. In *Prunus avium*, a SI diploid, deletions in the haplotype-specific region of
512 the pollen-S gene result in SC (Sonneveld et al. 2005). However, in Solanaceae, studies of
irradiated pollen find instances of pollen-S gene duplications, but no deletions, suggesting
514 that many deletions in the pollen-S gene incapacitate pollen (Golz et al. 1999, 2001). As
emphasized in Sonneveld et al. (2005), transgenic pollen-S gene knock-outs and
516 gain-of-function mutants could provide more definitive evidence regarding the idiosyncrasy
of the S-locus in *Prunus*, and the exact mechanism responsible for the maintenance of SI in
518 polyploids. Additional evidence stems from sequence diversity differences where the
inferred pollen gene of the S-locus in *Prunus* exhibits a higher degree of sequence diversity
520 than the pollen gene in *Antirrhinum* and *Petunia* (Ikeda et al. 2004; Kao and Tsukamoto

2004). Inferences involving the male-expressed gene are made more difficult by the enormous size of the F-box gene family, to which the pollen gene function of RSI belongs. The gene family contains several hundred members, including closely linked paralogous copies at the S-locus (Kuroda et al. 2002). Many questions remain about the exact identity, mode of evolution, and mechanism of action of pollen-expressed genes in RSI, as well as differences in expression between distantly related families (Newbigin et al. 2008). Additional mutational studies, which measure both breeding system and ploidy states within the same individuals, are necessary to determine more precisely the likelihood of occurrence of SI polyploids in Solanaceae and other families.

Ancient Polyploidy

Two separate studies, using similar methods, both find an ancient round of polyploidization within the Solanaceae. Schlueter et al. (2004) and Blanc and Wolfe (2004) use expressed sequence tags data to infer paralogous genes and the synonymous distance between them in order to estimate a distribution of synonymous distances. The expectation that a large scale duplication event would produce detectable peaks of synonymous distances can be used to estimate the temporal divergence approximating the date of such an event (Lynch and Conery 2000, 2003). Schlueter et al. (2004) report finding two apparent large scale duplication events, one ca. 52 mya, and another more recent duplication, with an uncertain time due to the large standard deviation around the synonymous distance peak. Using the results from Schlueter et al. (2004), Soltis et al. (2009) argued that the genome duplication could have occurred in the lineage leading to the Solanoideae based on the fact that more “basal” lineages in the Solanaceae have lower chromosome numbers (e.g., Cestroideae $x=7-12$; Penas et al. 2006) whereas the more “derived” Solanoideae is characterized by a basal chromosome number $x=12$ (Olmstead et al. 2008). In a separate study, Blanc and Wolfe (2004) infer a large scale duplication event 18-23 mya. Despite the wide acceptance of these and similar results, it remains unclear how frequently the employed methods falsely infer whole genome duplications, in part because accurate null models are difficult

to construct. Our comparative data, as well as genomic analyses obtained by Doganlar et al. (2002) and Wu et al. (2006), directly contradict the findings of whole genome duplication events in the recent ancestry of Solanaceae.

Given the strong association between SC and P, and evidence for irreversibility of the transition to SC in the Solanaceae (Ioerger et al. 1990; Miller and Venable 2000; Igić et al. 2004), we argue that ancient polyploidy is, in fact, highly unlikely to have occurred within Solanaceae. First, within Solanaceae, species in a group often termed “x=12” display a remarkably conserved pattern of synteny (Olmstead and Palmer 1992). The highly conserved karyotype evolution in a subset of that group, containing tomato, potato, eggplant, and pepper, requires the inference of only a few dozen rearrangements in karyotype evolution to explain the extant species patterns (Bonierbale et al. 1988; Livingstone et al. 1999; Doganlar et al. 2002). It is therefore extremely unlikely that any large scale or whole-genome duplications took place in this subset of species (ca. 1500 species, with a 15 my-old common ancestor). Furthermore, Wu et al. (2006) used a set of 2869 conserved gene-based markers, derived from single-copy genes shared by euasterid species, to construct comparative genetic maps of coffee and tomato, with a much older common ancestor (ca. 85 my). They found a strong pattern of conserved synteny between these distantly related species, with absence of “networks of synteny” expected to occur with whole genome duplication followed by selective gene loss. This finding is inconsistent with a proposed polyploidization event in the Solanaceae, which is supposed to have occurred around 20 mya (Blanc and Wolfe 2004; Schlueter et al. 2004). Consequently, we urge caution in interpreting the ancient duplication events from fitted K_s distributions between paralogous genes. The uncertainty in the null distribution of K_s values may be underestimated in part because the extent to which dysploidy affects the distribution is unclear, and assumptions regarding the operation of the molecular clock may be violated in an unpredictable manner.

Our results should be at least somewhat tempered, however, because of the known weaknesses in our data and models. Promising novel approaches (Smith et al. 2009) allow rapid construction of large-scale datasets with estimates of divergence times, which could allow us to more meaningfully estimate transition rates in units of time, instead of making the branch length assumptions as we did here. Such approaches are likely to yield better tests, especially when combined with our method for assessing relative pathway contributions, which are time-dependent. In addition, breeding systems are thought to affect diversification rate, and failure to incorporate differences in diversification rates may lead to incorrect inferences of trait evolution (Maddison 2006; Goldberg and Igić 2008). A multistate version of the model that allows for simultaneous inference of speciation and extinction along with transition rates (Maddison et al. 2007), used in conjunction with a tree with branch lengths proportional to time, will be required to assess this effect. Finally, the models currently in use do not allow for temporal or clade-specific rate heterogeneity, which provide a clear direction for further progress.

It may be that neither polyploidization nor breakdown of SI occurs at a constant family-wide rate, independent of unmeasured traits, for example, and it is unknown how robust these models are to such violations. Nevertheless, we offer what is likely a sound first approximation, which unites a large dataset of breeding system and ploidy, models informed by genetic and genomic data, and improved general methods for evaluation of the relative importance of different pathways in comparative phylogenetic studies.

ACKNOWLEDGEMENTS

We thank R. FitzJohn, J. Kohn, A. Raduski, and R. Ree for providing ideas and comments that improved the manuscript. A. Meade provided help with the operation of *BayesTraits*. A. Raduski provided the ISI distribution data and programming assistance. This work was supported by a grant from the National Science Foundation award DEB-0919089.

We wish to compute the transition probabilities of the two pathways from SI-D to SC-P:
 602 polyploidization from self-incompatibility (SI-D directly to SC-P, pathway 14), and loss of
 SI followed by polyploidization from SC (SI-D to SC-D to SC-P, pathway 134).

Consider the three states shown in black in Fig. 1, but insert an artificial division in
 the SC-P state so that it can be treated as two different states: state $4a$ contains the
 members of state 4 that arrived by pathway 14 , and state $4b$ contains those that arrived by
 pathway 134 (Fig. A1). The implied transition rate matrix (for a row vector of states
 ordered 1, 3, $4a$, $4b$) is:

$$\mathbf{Q} = \begin{pmatrix} -q_{13} - q_{14} & q_{13} & q_{14} & 0 \\ q_{31} & -q_{31} - q_{34} & 0 & q_{34} \\ q_{41} & 0 & -q_{41} & 0 \\ 0 & q_{43} & 0 & -q_{43} \end{pmatrix} \quad (\text{A1})$$

604 The probabilities of changing from one state to any other state after time t are given
 by the elements of the matrix $\mathbf{P} = \exp(\mathbf{Q}t)$. The pathway probabilities of interest are
 606 $W_{14}(t) = P_{14a}(t)$ and $W_{134}(t) = P_{14b}(t)$. In the general case where the reversal rates q_{31} , q_{41} ,
 and q_{43} are non-zero, our interpretation of a pathway is the last route taken to SC-P by
 608 time t , but landing in state $4a$ does not preclude having previously been in state $4b$, and
 vice versa.

In the special case for this system of no reverse transitions ($q_{31} = q_{41} = q_{43} = 0$), the
 transition probabilities of interest are:

$$W_{14}(t) = P_{14a}(t) = \frac{q_{14}}{q_{13} + q_{14}} (1 - e^{-(q_{13} + q_{14})t}) \quad (\text{A2})$$

and

$$W_{134}(t) = P_{14b}(t) = \frac{q_{13}}{q_{13} + q_{14}} \left(1 - \frac{1}{q_{13} + q_{14} - q_{34}} [(q_{13} + q_{14})e^{-q_{34}t} - q_{34}e^{-(q_{13}+q_{14})t}] \right) \quad (\text{A3a})$$

unless $q_{14} = q_{34} - q_{13}$, in which case

$$W_{134}(t) = P_{14b}(t) = \frac{q_{13}}{q_{34}} (1 - e^{-q_{34}t} [1 + q_{34}t]) . \quad (\text{A3b})$$

LITERATURE CITED

Acosta, M. C., A. del V. Ordonez, A. A. Cocucci, and E. A. Moscone, 2006. Chromosome reports in South American Nicotianeae (Solanaceae) with particular reference to *Nierembergia*. *Annals of the Missouri Botanical Garden* 93:634–646.

Ando, T., S. Iida, H. Kokubun, Y. Ueda, and E. Marchesi, 1998. Distribution of infraspecific taxa of *Petunia axillaris* (Solanaceae) in Uruguay as revealed by discriminant analyses. *Acta Phytotaxon Geobot.* 45:95–109.

Ando, T., H. Kokubun, H. Watanabe, N. Tanaka, T. Yukawa, G. Hashimoto, E. Marchesi, E. Suarez, and I. L. Basualdo, 2005. Phylogenetic analysis of *Petunia sensu* Jussieu (Solanaceae) using chloroplast DNA RFLP. *Annals of Botany* 96:289–297.

Badr, A., S. F. Khalifa, Aboel-Atta, and Abou-El-Enain, 1997. Chromosome criteria and taxonomic relationships in the Solanaceae. *Cytologia* 62:103–113.

Barker, M. S., N. C. Kane, M. Matvienko, A. Kozik, R. W. Michelmore, S. J. Knapp, and L. H. Riesberg, 2008. Multiple paleopolyploidizations during the evolution of the Compositae reveal parallel patterns of duplicate gene retention after millions of years. *Molecular Biology and Evolution* 25:2445–2455.

Barringer, B. C., 2007. Polyploidy and self-fertilization in flowering plants. *American Journal of Botany* 94:1527–1533.

- 628 Bawa, K. S., 1974. Breeding systems of tree species of a lowland tropical community.
Evolution 28:85–92.
- 630 Blanc, G., A. Barakat, R. Guyot, R. Cooke, and M. Delseny, 2000. Extensive duplication
and reshuffling in the *Arabidopsis* genome. Plant Cell 12:1093–1101.
- 632 Blanc, G., K. Hokamp, and K. H. Wolfe, 2003. A recent polyploidy superimposed on older
large-scale duplications in the *Arabidopsis* genome. Genome Research 13:137–144.
- 634 Blanc, G. and K. H. Wolfe, 2004. Widespread paleopolyploidy in model plant species
inferred from age distributions of duplicate genes. The Plant Cell 16:1667–1678.
- 636 Bohs, L., 2000. Insights into the *Witheringia solanacea* (Solanaceae) complex in Costa
Rica. I. Breeding systems and crossing studies. Biotropica 32:70–79.
- 638 ———, 2007. Phylogeny of the *Cyphomandra* clade of the genus *Solanum* (Solanaceae)
based on ITS sequence data. Taxon 56:1012–1026.
- 640 Bonierbale, M., R. Plaisted, and S. Tanksley, 1988. RFLP maps based on a common set of
clones reveal modes of chromosomal evolution in potato and tomato. Genetics
642 120:1095–1103.
- Bowers, J. E., B. A. Chapman, J. Rong, and A. H. Paterson, 2003. Unravelling angiosperm
644 genome evolution by phylogenetic analysis of chromosomal duplication events. Nature
422:433–438.
- 646 Brewbaker, J. L., 1953. Oppositional allelism in diploid and autotetraploid *Trifolium*
hybridum l. Genetics 38:444–455.
- 648 ———, 1954. Incompatibility in an autotetraploid *Trifolium repens*. I. Competition and
self-compatibility. Genetics 39:307–316.
- 650 ———, 1958. Self-compatibility in tetraploid strains of *Trifolium hybridum*. Hereditas
44:547–553.

- 652 ———, 1959. Biology of the angiosperm pollen grain. *Indian Journal of Genetics and*
Plant Breeding 19:121–133.
- 654 Brunet, J. and A. Liston, 2001. Technical comment on polyploidy and gender dimorphism.
Science 291:1441–1442.
- 656 Bull, J. J. and E. L. Charnov, 1985. On irreversible evolution. *Evolution* 39:1149–1155.
- Byrne, K. P. and G. Blanc, 2006. Computational analyses of ancient polyploidy. *Current*
658 *Bioinformatics* 1:131–146.
- Camadro, E. and S. J. Peloquin, 1981. Cross-incompatibility between two sympatric
660 polyploid *Solanum* species. *Theoretical and Applied Genetics* 60:65–70.
- Charlesworth, D., 1985. Distribution of dioecy and self-incompatibility in angiosperms. Pp.
662 237–268, *in* P. J. Greenwood, P. H. Harvey, and M. Slatkin, eds. *Evolution: Essays in*
Honour of John Maynard Smith. Cambridge University Press, Cambridge.
- 664 Chawla, B., R. Bernatzky, W. Liang, and M. Marcotrigiano, 1997. Breakdown of
self-incompatibility in tetraploid *Lycopersicon peruvianum*: inheritance and expression of
666 S-related proteins. *Theoretical and Applied Genetics* 95:992–996.
- Clarkson, J. J., S. Knapp, V. F. Garcia, R. G. Olmstead, A. R. Leitch, and M. W. Chase,
668 2004. Phylogenetic relationships in *Nicotiana* (Solanaceae) inferred from multiple plastid
DNA regions. *Molecular Phylogenetics and Evolution* 33:75–90.
- 670 Correll, D. S., 1962. *The Potato and Its Wild Relatives*. Texas Research Foundation,
Renner, Texas.
- 672 Crane, M. B. and D. Lewis, 1942. Genetical studies in pears. III. Incompatibility and
sterility. *Journal of Genetics* 43:31–42.
- 674 Darlington, C. D. and A. P. Wylie, 1956. *Chromosome Atlas of Flowering Plants*. G. Allen
& Unwin, New York.

- 676 Dickinson, T. A., E. Lo, and N. Talent, 2007. Polyploidy, reproductive biology, and
Rosaceae: understanding evolution and making classifications. *Plant Systematics and*
678 *Evolution* 266:59–78.
- Doganlar, S., A. Frary, C. M. Daunay, R. N. Lester, and S. D. Tanksley, 2002. A
680 comparative genetic linkage map of eggplant (*Solanum melongena*) and its implication to
genome evolution in the Solanaceae. *Genetics* 161:1697–1711.
- 682 East, E. and A. Mangelsdorf, 1925. A new interpretation of the heredity behaviour of
self-sterile plants. *Proceedings of the National Academy of Science, USA* 11:166–171.
- 684 Entani, T., S. Takayama, M. Iwano, H. Shiba, F.-S. Che, and A. Isogai, 1999. Relationship
between polyploidy and pollen self-incompatibility phenotype in *Petunia hybrida* Vilm.
686 *Bioscience, Biotechnology, and Agrochemistry* 63:1882–1888.
- Felsenstein, J., 1985. Phylogenies and the comparative method. *The American Naturalist*
688 125:1–15.
- Gaiser, L., 1926. A list of chromosome numbers in angiosperms. *Genetica* 8:401–484.
- 690 Goldberg, E. E. and B. Igić, 2008. On phylogenetic tests of irreversible evolution.
Evolution 62:2727–2741.
- 692 Goldblatt, P., 1980. Polyploidy in angiosperms: monocotyledons. Plenum Press, New York.
- Goldblatt, P. and D. E. Johnson, 1979. Index to plant chromosome numbers (IPCN).
694 Missouri Botanical Garden, St. Louis. URL
<http://mobot.mobot.org/W3T/Search/ipcn.html>.
- 696 Golz, J. F., H.-Y. Oh, V. Su, M. Kusaba, and E. Newbigin, 2001. Genetic analysis of
Nicotiana pollen-part mutants is consistent with the presence of the *S*-ribonuclease
698 inhibitor at the *S* locus. *Proceedings of the National Academy of Science, USA*
98:15372–15376.

- 700 Golz, J. F., V. Su, A. E. Clarke, and E. Newbigin, 1999. A molecular description of
mutations affecting the pollen component of the *Nicotiana glauca* *S* locus. *Genetics*
702 152:1123–1135.
- Goodspeed, T. H., 1954. The genus *Nicotiana*, vol. 16. *Chronica Botanica*, An
704 International Collection of Studies in the Method and History of Biology and
Agriculture, Waltham, Mass.
- 706 Grafen, A., 1989. The phylogenetic regression. *Philosophical Transactions of the Royal*
Society of London. Series B, Biological Sciences 326:119–157.
- 708 Grant, C., 1956. The influence of breeding habit on the outcome of natural hybridization
in plants. *The American Naturalist* 90:319–322.
- 710 Grant, V., 1963. *The Origin of Adaptations*. Columbia University Press, New York.
———, 1971. *Plant Speciation*. Columbia University Press, New York and London.
- 712 Halverson, K., S. B. Heard, J. D. Nason, and J. O. S. III, 2008. Origins, distribution, and
local co-occurrence of polyploid cytotypes in *Solidago altissima* (Asteraceae). *American*
714 *Journal of Botany* 95:50–58.
- Hauck, N. R., H. Yamane, R. Tao, and A. F. Iezzoni, 2002. Self-compatibility and
716 incompatibility in tetraploid sour cherry (*Prunus cerasus* L.). *Sexual Plant*
Reproduction 15:39–46.
- 718 ———, 2006. Accumulation of nonfunctional *S*-haplotypes results in the breakdown of
gametophytic self-incompatibility in tetraploid prunus. *Genetics* 172:1191–1198.
- 720 Hawkes, J. G. and J. P. Hjerting, 1969. *The Potatoes of Argentina, Brazil, Paraguay, and*
Uruguay. A Biosystematic Study. Oxford University Press, Great Britain.
- 722 Hermesen, J. G. T. and A. J. E. D. Boer, 1971. The effect of colchicine treatment on
Solanum acaule and *S. bulbocastanum*; a complete analysis of ploidy chimeras in *S.*
724 *bulbocastanum*. *Euphytica* 20:171–180.

Hijmans, R. J., T. Gavrilenko, S. Stephenson, J. Bamberg, A. Salas, and D. M. Spooner,
726 2007. Geographical and environmental range expansion through polyploidy in wild
potatoes (*Solanum* section *Petota*). *Global Ecology and Biogeography* 16:485–495.

728 Hua, Z.-H., A. Fields, and T.-h. Kao, 2008. Biochemical models for S-RNase-based
self-incompatibility. *Molecular Plant* 1:575–585.

730 Huang, S.-X., H.-Q. Wu, Y.-R. Li, J. Wu, S.-J. Zhang, W. Heng, and S.-L. Zhang, 2008.
Competitive interaction between two functional *S*-haplotypes confer self-compatibility on
732 tetraploid Chinese cherry (*Prunus pseudocerasus* Lindl. CV. Nanjing Chuisi). *Plant Cell*
Reports 27:1075–1085.

734 Husband, B. C., B. Ozimec, S. L. Martin, and L. Pollock, 2008. Mating consequences of
polyploid evolution in flowering plants: Current trends and insights from synthetic
736 polyploids. *International Journal of Plant Sciences* 169:195–206.

Igić, B., L. Bohs, and J. R. Kohn, 2004. Historical inferences from the self-incompatibility
738 locus. *New Phytologist* 161:97–105.

———, 2006. Ancient polymorphism reveals unidirectional breeding system shifts.

740 *Proceedings of the National Academy of Sciences* 103:1359–1363.

Igić, B. and J. R. Kohn, 2001. Evolutionary relationships among self-incompatibility
742 RNases. *Proceedings of the National Academy of Sciences* 98:13167–13171.

———, 2006. The distribution of plant mating systems: Study bias against obligately
744 outcrossing species. *Evolution* 60:1098–1103.

Igić, B., R. Lande, and J. R. Kohn, 2008. Loss of self-incompatibility and its evolutionary
746 consequences. *Int J Plant Sci* 169:93–104.

Ikeda, K., B. Igić, K. Ushijima, H. Yamane, N. R. Hauck, R. Nakano, H. Sassa, A. F.

748 Iezzoni, J. R. Kohn, and R. Tao, 2004. Primary structural features of the *S*
haplotype-specific F-box protein, SFB in *Prunus*. *Sexual Plant Reproduction* 16:235–243.

- 750 Ioerger, T. R., A. G. Clark, and T.-H. Kao, 1990. Polymorphism at the self-incompatibility
locus in Solanaceae predates speciation. Proceedings of the National Academy of
752 Sciences of the United States of America 87:9372–9735.
- Kao, T. H. and T. Tsukamoto, 2004. The molecular and genetic bases of S-RNase-based
754 self-incompatibility. The Plant Cell 16:S72–S83.
- Kass, R. E. and A. E. Raftery, 1995. Bayes factors. Journal of the American Statistical
756 Association 90:773–795.
- Kirch, H. H., H. Uhrig, F. Lottspeich, F. Salamini, and R. D. Thompson, 1989.
758 Characterization of proteins associated with self-incompatibility in *Solanum tuberosum*.
Theoretical and Applied Genetics 78:581–588.
- 760 Ku, H. M., T. Vision, J. Liu, and S. D. Tanksley, 2000. Comparing sequenced segments of
the tomato and *Arabidopsis* genomes: large-scale duplication followed by selective gene
762 loss creates a network of synteny. Proceedings of the National Academy of Science, USA
97:9121–9126.
- 764 Kuroda, H., N. Takahashi, H. Shimada, M. Seki, K. Shinozaki, and M. Matsui, 2002.
Classification and expression analysis of *Arabidopsis* F-box-containing protein genes.
766 Plant Cell and Physiology 43:1073–85.
- Levin, D. A., 1983. Polyploidy and the novelty of flowering plants. The American
768 Naturalist 122:1–25.
- Levin, R. A. and J. S. Miller, 2005. Relationships within tribe Lycieae (Solanaceae):
770 Paraphyly of *Lycium* and multiple origins of gender dimorphism. American Journal of
Botany 92:2044–2053.
- 772 Levin, R. A., N. R. Myers, and L. Bohs, 2006. Phylogenetic relationships among the “spiny
solanums” (*Solanum* subgenus *Leptostemonum*, Solanaceae). American Journal of
774 Botany 93:157–169.

Lewis, D., 1943. Physiology of incompatibility on plants. III. Autopolyploids. *Journal of*
776 *Genetics* 45:171–185.

———, 1947. Competition and dominance of incompatibility alleles in diploid pollen.
778 *Heredity* 1:85–108.

———, 1979. Sexual Incompatibility in Plants, *The Institute of Biology's Studies in*
780 *Biology*, vol. 110. Edward Arnold Limited.

Lewis, W. H. and Y. Suda, 1976. Diploids and polyploids from a single species population:
782 temporal adaptations. *The Journal of Heredity* 67:391–393.

Lin, C., L. A. Mueller, J. McCarthy, D. Crouzillat, and V. Petiard, 2005. Coffee and
784 tomato share common gene repertoires as revealed by deep sequencing of seed and cherry
transcripts. *Theoretical and Applied Genetics* 112:114–130.

786 Livermore, J. and F. E. Johnstone, 1940. The effect of chromosome doubling on the
crossability of *Solanum chacoense*, *Solanum jamesii* and *S. bulbocastanum* with *S.*
788 *tuberosum*. *American Potato Journal* 17:170–173.

Livingstone, K., V. Lackney, J. Blauth, R. van Wijk, and M. Jahn, 1999. Genome mapping
790 in *Capsicum* and the evolution of genome structure in the Solanaceae. *Genetics*
152:1183–1202.

792 Lloyd, D. G., 1965. Evolution of self-compatibility and racial differentiation in
Leavenworthia (Cruciferae). *Contributions from the Gray Herbarium of Harvard*
794 *University* 195:3–134.

Lynch, M. and J. S. Conery, 2000. The evolutionary fate and consequence of duplicate
796 genes. *Science* 290:1151–1155.

———, 2003. The evolutionary demography of duplicate genes. *Journal of Structural and*
798 *Functional Genomics* 3:35–44.

Mable, B. K., 2004. Polyploidy and self-compatibility: is there an association? *New*
800 *Phytologist* 162:803–811.

Mace, E. S., C. G. Gebhardt, and R. N. Lester, 1999. AFLP analysis of genetic
802 relationships in the tribe Datureae (Solanaceae). *Theoretical and Applied Genetics* Pp.
634–641.

804 Maddison, W. P., 2006. Confounding asymmetries in evolutionary diversification and
character change. *Evolution* 60:1743–1746.

806 Maddison, W. P., P. E. Midford, and S. P. Otto, 2007. Estimating a binary character's
effect on speciation and extinction. *Systematic Biology* 56:701–710.

808 Marks, G. E., 1965. Cytogenetic studies of tuberous *Solanum* species. III. Species
relationships in some South and Central American species. *New Phytologist* 64:293–306.

810 Masterson, J., 1994. Stomatal size in fossil plants: Evidence for polyploidy in majority of
angiosperms. *Science* 264:421–424.

812 McClure, B., 2009. Darwin's foundation for investigating self-incompatibility and the
progress toward a physiological model for S-RNase-based SI. *Journal of Experimental*
814 *Botany* 60:1069–1081.

Meyers, L. A. and D. A. Levin, 2006. On the abundance of polyploids in flowering plants.
816 *Evolution* 60:1198–1206.

Miller, J. S., R. A. Levin, and N. M. Feliciano, 2008. A tale of two continents: Baker's rule
818 and the maintenance of self-incompatibility in *Lycium* (Solanaceae). *Evolution*
62:1052–1065.

820 Miller, J. S. and D. L. Venable, 2000. Polyploidy and the evolution of gender dimorphism
in plants. *Science* 289:2335–2338.

- 822 Mione, T., R. C. Olmstead, R. K. Jansen, and G. J. Anderson, 1994. Systematic
implications of chloroplast DNA variation in *Jaltomata* and selected Physaloid genera
824 (Solanaceae). American Journal of Botany Pp. 912–918.
- Montero-Castro, J. C., A. Delgado-Salinas, E. D. Luna, and L. E. Eguiarte, 2006.
826 Phylogenetic analysis of *Cestrum* section *Habrothamnus* (Solanaceae) based on plastid
and nuclear DNA sequences. Systematic Botany 31:843–850.
- 828 de Nettancourt, D., 1977. Incompatibility in Angiosperms. Springer-Verlag, Berlin.
- de Nettancourt, D., U. Laneri, E. Capaccio, M. Westerhof, and R. Ecochard, 1974.
830 Polyploidy and induced mutations in plant breeding, chap. Self-compatibility in a
spontaneous tetraploid of *Lycopersicon peruvianum* Mill., Pp. 74–84. I.A.E.A., Vienna.
- 832 Newbigin, E., T. Paape, and J. R. Kohn, 2008. RNase-based self-incompatibility: puzzled
by pollen S. Plant Cell 20:2286–2292.
- 834 Nunes, M. D. S., R. A. M. Santos, S. M. Ferreira, J. Vieira, and C. P. Vieira, 2006.
Variability patterns and positively selected sites at the gametophytic self-incompatibility
836 pollen *SFB* gene in a wild self-incompatible *Prunus spinosa* (Rosaceae) population. New
Phytologist 172:577–587.
- 838 Olmstead, R. G. and L. Bohs, 2007. A summary of molecular systematic research in
Solanaceae: 1982-2006. Acta Horticulturae 745:255–268.
- 840 Olmstead, R. G., L. Bohs, H. A. Migid, E. Santiago-Valentin, V. F. Garcia, and S. M.
Collier, 2008. A molecular phylogeny of the Solanaceae. Taxon Pp. 1159–1181.
- 842 Olmstead, R. G. and J. D. Palmer, 1992. A chloroplast DNA phylogeny of the Solanaceae:
subfamilial relationships and character evolution. Annals of the Missouri Botanical
844 Garden 79:346–360.

- Olmstead, R. G. and J. A. Sweere, 1994. Combining data in phylogenetic systematics: An
846 empirical approach using three molecular data sets in the Solanaceae. *Systematic*
Biology 43:467–481.
- 848 Olmstead, R. G., J. A. Sweere, R. E. Spangler, L. Bohs, and J. D. Palmer, 1999. Phylogeny
and provisional classification of the Solanaceae DNA. Pp. 111–137, *in* M. Nee, D. E.
850 Symon, R. N. Lester, and J. P. Jesop, eds. *Solanaceae*. Royal Botanic Gardens, Kew.
- Otto, S. P. and J. Whitton, 2000. Polyploid incidence and evolution. *Annual Review of*
852 *Genetics* 34:401–437.
- Paape, T., B. Igic, S. D. Smith, R. Olmstead, L. Bohs, and J. R. Kohn, 2008. A
854 15-million-year-old genetic bottleneck. *Molecular Biology and Evolution* 25:655–663.
- Pagel, M., 1994. Detecting correlated evolution on phylogenies: a general method for the
856 comparative analysis of discrete characters. *Proceedings of the Royal Society*
B-Biological Sciences 255:37–45.
- 858 ———, 1999. The maximum likelihood approach to reconstructing ancestral character
states of discrete characters on phylogenies. *Systematic Biology* 48:612–622.
- 860 Pagel, M. and A. Meade, 2006. Bayesian analysis of correlated evolution of discrete
characters by reversible-jump Markov chain Monte Carlo. *The American Naturalist*
862 168:808–825.
- Pandey, K. K., 1968. Colchicine induced changes in the self-incompatibility behavior of
864 *Nicotiana*. *Genetica* 39:257–271.
- Paterson, A. H., J. E. Bowers, M. D. Burow, X. Draye, C. G. Elsik, C. X. Jiang, C. S.
866 Katsar, T. H. Lan, Y. R. Lin, R. Ming, and R. J. Wright, 2000. Comparative genomics
of plant chromosomes. *Plant Cell* 12:1523–1540.

- 868 Penas, M. L. L., F. E. Chiarini, G. Bernadello, and C. B. de Rojas, 2006. Karyotypes of
some species of *Cestrum*, *Sessea*, and *Vestia* (tribe Cestreae, Solanaceae). *Caryologia*
870 59:131–137.
- Peralta, I. E. and D. M. Spooner, 2001. Granule-bound starch synthase (GBSSI) gene
872 phylogeny of wild tomatoes (*Solanum* l. section *Lycopersicon* [Mill.] Wettst. subsection
Lycopersicon). *American Journal of Botany* 88:1888–1902.
- 874 Perez, F., M. T. K. Arroyo, R. Medel, and M. A. HersHKovitz, 2006. Ancestral
reconstruction of flower morphology and pollination systems in *Schizanthus*
876 (*Solanaceae*). *American Journal of Botany* 93:1029–1038.
- Prohens, J., G. J. Anderson, J. M. Blanca, J. Canizares, E. Zuriaga, and F. Nuez, 2006.
878 The implications of AFLP data for the systematics of the wild species of *Solanum*
section *Basarthurum*. *Systematic Botany* 31:208–216.
- 880 Pruski, K., 2007. Tissue culture propagation of Mongolian cherry (*Prunus fruticosa* L.)
and Nanking cherry (*Prunus tomentosa* L.). Pp. 391–407, *in* S. J. Jain and H. Haggman,
882 eds. *Protocols for Micropropagation of Woody Trees and Fruits*. Springer.
- Raftery, A. E., 1996. Hypothesis testing and model selection. Pp. 163–188, *in* W. R. Gilks,
884 S. Richardson, and D. J. Spiegelhalter, eds. *Markov chain Monte Carlo in practice*.
Chapman & Hall, London.
- 886 Ramsey, J. and D. W. Schemske, 1998. Pathways, mechanisms, and rates of polyploid
formation in flowering plants. *Annual Review of Ecology and Systematics* 29:467–501.
- 888 ———, 2002. Neopolyploidy in flowering plants. *Annual Review of Ecology and*
Systematics 33:589–639.
- 890 Rick, C. M., 1986. Reproductive isolation in the *Lycopersicon peruvianum* complex. Pp.
477–495, *in* W. G. D’Arcy, ed. *Solanaceae, biology and systematics*. Columbia University
892 Press, New York, New York, USA.

Rick, C. M. and S. D. Tanksley, 1981. Genetic variation in *Solanum pennellii*: Comparisons

with two other sympatric tomato species. *Plant Systematics and Evolution* 139:11–45.

Schlueter, J. A., P. Dixon, C. Granger, D. Grant, L. Clark, J. J. Doyle, and R. C.

Schoenmaker, 2004. Mining EST databases to resolve evolutionary events in major crop species. *Genome* 47:868–876.

Simillion, C., K. Vandepoele, M. C. E. V. Montagu, M. Zabeau, and Y. V. de Peer, 2002.

The hidden duplication past of *Arabidopsis thaliana*. *Proceedings of the National Academy of Science, USA* 99:13627–13632.

Smith, S. A., J. M. Beaulieu, and M. J. Donoghue, 2009. Mega-phylogeny approach for

comparative biology: An alternative to supertree and supermatrix approaches. *BMC Evolutionary Biology* 9.

Smith, S. D. and D. A. Baum, 2006. Phylogenetics of the florally diverse Andean clade

Iochrominae (Solanaceae). *American Journal of Botany* 93:1140–1153.

Soltis, D. E., V. A. Albert, J. Leebens-Mack, C. D. Bell, A. H. Paterson, C. Zheng,

D. Sankoff, C. W. DePamphilis, P. K. Wall, and P. S. Solits, 2009. Polyploidy and angiosperm diversification. *American Journal of Botany* 96:336–348.

Sonneveld, T., K. R. Tobutt, S. P. Vaughan, and T. P. Robbins, 2005. Loss of pollen-*S*

function in two self-compatible selections of *Prunus avium* is associated with deletion/mutation of an *S* haplotype-specific F-Box gene. *The Plant Cell* 17:37–51.

Spooner, D. M., G. J. Anderson, and R. K. Jansen, 1993. Chloroplast DNA evidence for

the interrelationships of tomatoes, potatoes, and pepinos (Solanaceae). *American Journal of Botany* 80:676–688.

Spooner, D. M., D. S. Douches, and A. C. M., 1992. Allozyme variation within *Solanum*

sect. *Petota*, ser. *Etuberosa* (Solanaceae). *American Journal of Botany* 79:467–471.

- Stebbins, G. L., 1938. Cytological characteristics associated with the different growth
918 habits in the dicotyledons. *American Journal of Botany* 25:189–198.
- , 1950. *Variation and Evolution in Plants*. Columbia University Press, New York.
- 920 ———, 1957. Self-fertilization and population variability in the higher plants. *American Naturalist* 91.
- 922 ———, 1971. *Chromosomal Evolution in Higher Plants*. Addison-Wesley, London, UK.
- , 1974. *Flowering plants: evolution above the species level*. Belknap Press of
924 Harvard University Press, Cambridge, Mass.
- , 1980. Polyploidy in plants: Unsolved problems and prospects. Pp. 495–520, *in*
926 W. H. Lewis, ed. *Polyploidy: Biological Relevance*. Plenum, New York.
- Stone, J. L., 2002. Molecular mechanisms underlying the breakdown of gametophytic
928 self-incompatibility. *The Quarterly Review of Biology* 77:17–30.
- Stout, A. B. and C. Chandler, 1942. Hereditary transmission of induced tetraploidy and
930 compatibility in fertilization. *Science* 96:257–258.
- Tu, T.-Y., H. Sun, Z.-J. Gu, and J.-P. Yue, 2005. Cytological studies on the
932 Sino-Himalayan endemic *Anisodus* and four related genera from the tribe *Hyoscyameae*
(Solanaceae) and their systematic and evolutionary implications. *Botanical Journal of*
934 *the Linnean Society* 147:457–468.
- Vieira, J., R. Santos, S. Ferreira, and C. Vieira, 2008. Inferences on the number and
936 frequency of S-pollen gene (SFB) specificities in the polyploid *Prunus spinosa*. *Heredity*
101:351–358.
- 938 Vision, T. J., D. G. Brown, and S. D. Tanksley, 2000. The origins of genomic duplications
in *Arabidopsis*. *Science* 290:2114–2117.

- 940 Walsh, B. M. and S. B. Hoot, 2001. Phylogenetic relationships of *Capsicum* (Solanaceae)
using DNA. International Journal of Plant Sciences 162:1409–1418.
- 942 Weese, T. L. and L. Bohs, 2007. A three-gene phylogeny of the genus *Solanum*
(Solanaceae). Systematic Botany 32:445–463.
- 944 Welsh, K. J. and K. C. Sink, 1981. Morphogenetic responses of *Browallia* leaf sections and
callus. Annals of Botany 48:583–590.
- 946 Wendel, J. F., 2000. Genome evolution in polyploids. Plant Molecular Biology 42:225–249.
- Whitson, M. and P. S. Manos, 2005. Untangling *Physalis* (Solanaceae) from the
948 Physaloids: A two-gene phylogeny of the Physalinae. Systematic Botany 30:216–230.
- Wikstrom, N., V. Savolainen, and M. W. Chase, 2001. Evolution of the angiosperms:
950 calibrating the family tree. Proc. Biol. Sci. 268:2211–2220.
- Wolfe, K. H., 2001. Yesterday’s polyploids and the mystery of diploidization. Nature
952 Reviews Genetics 2:333–341.
- Wu, F., L. A. Mueller, D. Crouzillat, V. Petiard, and S. D. Tanksley, 2006. Combining
954 bioinformatics and phylogenetics to identify large sets of single-copy orthologous genes
(COSII) for comparative, evolutionary and systematic studies: A test case in the
956 euasterid plant clade. Genetics 174:1407–1420.
- Yates, F., 1934. Contingency tables involving small numbers and the χ^2 test. Journal of
958 the Royal Statistical Society 1:217–235.
- Yeung, K., J. S. Miller, A. E. Savage, B. C. Husband, B. Igic, and J. R. Kohn, 2005.
960 Association of ploidy and sexual system in *Lycium californicum* (Solanaceae). Evolution
59:2048–2055.

	D		P		Total
	Obs	Exp	Obs	Exp	
SI	164	138.7	1	26.3	165
SC	179	204.3	64	38.7	244
Total	343		65		408

Table 1: Table of observed (Obs) and expected (Exp) character state values for species where both ploidy and compatibility states are known. Polymorphic data were collapsed into SI and D categories as described in text. SI-P is significantly under-represented ($\chi^2 = 48.6$, $df = 1$; $p < 0.01$).

Rates	Unit Branch Lengths		Grafen Branch Lengths	
	Restricted	Full	Restricted	Full
q_{13}	0.25	0.16	5.6	16.4
q_{31}	0	0.08	0	12.1
q_{14}	0.08	0.05	1.3	3.6
q_{41}	0	0.16	0	11.5
q_{34}	0.05	0.04	6.0	4.3
q_{43}	0	0.11	0	7.8

Table 2: Inferred median rates of character evolution from the *Multistate* analysis. Results are shown for both the restricted model of character evolution, where the evolution of ploidy and self-compatibility is irreversible, and unrestricted (full) model on the phylogenetic tree with unit or Grafen branch lengths. Under both models and branch length assignment methods, self-incompatibility is inferred to be lost more often through mutations within the diploids than through polyploidization ($q_{13} > q_{14}$ with probability ≥ 0.99 in all cases, as computed from the posterior rate distributions).




Figure 1: A schematic representation of the model of evolution for breeding system (SI vs. SC) and ploidy (D vs. P). Arrows forming the outer square represent the rates estimated in our *Discrete* model of character evolution (Pagel 1994). A model of correlated evolution estimates all eight rates independently (only the four forward rates q_{12} , q_{13} , q_{34} , and q_{24} are shown) and is compared to a model of independent evolution in which four rates are estimated (only the two forward rates $q_{12} = q_{34}$, $q_{13} = q_{24}$ are shown). The solid black arrows forming the lower left triangle represent the model of character evolution in the *Multistate* framework. SI-D = state 1, SC-D = state 3, SC-P = state 4. SI-P = state 2 is not observed in our dataset and cannot be included in the *Multistate* framework.




Figure 2: The distribution of haploid chromosome numbers in the Solanaceae for 917 species. Arrows mark the most frequently observed chromosome counts. Chromosome number found in somatic cells is indicated with “ $2n$ ”, and the number preceding “ x ” refers to the inferred ploidy. The marked numbers $2n=2x=24$, $2n=3x=36$, $2n=4x=48$ and $2n=6x=72$ refer to the diploid, triploid, tetraploid, and hexaploid cytotypes, respectively.




Figure 3: The distribution of ISI values. Both self- and cross-pollinations were performed manually, and fruit and seed set was scored for 92 species of Solanaceae. Species with ISI values greater than 0.8 are classified SI, whereas species with ISI values less than 0.8 are classified as SC. The ten species with negative ISI values (ranging from -5.5 to -0.03) were set to zero.




Figure 4: A supertree representing a phylogenetic hypothesis for 266 Solanaceae taxa for which breeding system and ploidy is known. The analyses were performed on a tree with unit and Grafen branch lengths (see text for details). Open circles denote self-incompatible diploids; dotted circles, self-compatible diploids; closed circles, self-compatible polyploids.




Figure 5: Probabilities of the direct and indirect pathways from self-incompatible diploids (state 1) to self-compatible polyploids (state 4). The pathway probabilities $W_{14}(t)$ (solid lines) and $W_{134}(t)$ (dashed lines) were computed with the median values of the posterior rate distribution under the restricted model (Table 2). Results are shown for the unit branch length tree (A; $q_{13} = 0.25$, $q_{14} = 0.08$, $q_{34} = 0.05$) and for the Grafen branch length tree (B; $q_{13} = 5.6$, $q_{14} = 1.3$, $q_{34} = 6.0$). In each case, the elapsed times plotted cover the full depth of the tree (maximum node depth is 17 for A, root age is 1 for B). With unit branch lengths, the single step pathway (14) is more probable over time spans of approximately $t \leq 10.5$ units. Over longer time intervals, the two-step pathway (134) becomes dominant. Both pathways therefore may be important contributors in the correlated evolution to self-compatible polyploids in Solanaceae. With Grafen branch lengths, the single step pathway (14) is more probable only over time spans of $t \leq 0.1$. Pathway 134 therefore appears to be dominant.




Figure 6: Contour plots of the difference in probability between the two-step pathway (134) and the one-step pathway (14). Results are shown for the unit branch length tree (A) and for the Grafen branch length tree (B). The solid contour lines represent the cumulative probability density of $\Delta W(t) = W_{134}(t) - W_{14}(t)$, computed over the posterior rate distribution; their levels are printed on the right. At each time t , there is a 10% chance that $\Delta W(t)$ is less than the value shown by the line marked 0.1, and similarly for the ten other levels shown. The left dashed line (at $t = 5.7$ in A, at $t = 0.04$ in B) marks the time at which there is a 90% probability that pathway 14 is more likely. The right dashed line (at $t = 11.3$ in A, at $t = 0.09$ in B) marks the time at which pathway 134 becomes more likely. Over time spans that are long relative to the tree age (time values near the upper limits plotted), pathway 134 dominates in the Grafen tree but both pathways contribute in the unit tree.




Figure A1: Schematic representation of the states used in the pathway probability derivation. States 1 (SI-D) and 3 (SC-D) are as for the *Multistate* analysis (Fig. 1), but state 4 (SC-P) is divided into two. Transitions to SC-P from SI-D are tracked in state 4a, and transitions to SC-P from SC-D are tracked in state 4b.