

## Week 4 at a glance

### We will be learning and practicing to:

- Translate between different representations to illustrate a concept.
  - Translating between symbolic and English versions of statements using precise mathematical language
  - Translating between truth tables (tables of values) and compound propositions
- Use precise notation to encode meaning and present arguments concisely and clearly
  - Listing the truth tables of atomic boolean functions (and, or, xor, not, if, iff)
  - Defining functions, predicates, and binary relations using multiple representations
- Know, select and apply appropriate computing knowledge and problem-solving techniques. Reason about computation and systems. Use mathematical techniques to solve problems. Determine appropriate conceptual tools to apply to new situations. Know when tools do not apply and try different approaches. Critically analyze and evaluate candidate solutions.
  - Evaluating compound propositions
  - Judging logical equivalence of compound propositions using symbolic manipulation with known equivalences, including DeMorgan's Law
  - Writing the converse, contrapositive, and inverse of a given conditional statement
  - Determining what evidence is required to establish that a quantified statement is true or false
  - Evaluating quantified statements about finite and infinite domains

### TODO:

Review quiz based on class material each day (due Friday April 26, 2024)

Start reviewing for Test 1, in class next week on Friday May 3, 2024.

# Week 4 Monday: Conditionals and Logical Equivalence

| Input |     | Output                      |                              |                           |                                  |                                        |
|-------|-----|-----------------------------|------------------------------|---------------------------|----------------------------------|----------------------------------------|
| $p$   | $q$ | Conjunction<br>$p \wedge q$ | Exclusive or<br>$p \oplus q$ | Disjunction<br>$p \vee q$ | Conditional<br>$p \rightarrow q$ | Biconditional<br>$p \leftrightarrow q$ |
| $T$   | $T$ | $T$                         | $F$                          | $T$                       | $T$                              | $T$                                    |
| $T$   | $F$ | $F$                         | $T$                          | $T$                       | $F$                              | $F$                                    |
| $F$   | $T$ | $F$                         | $T$                          | $T$                       | $T$                              | $F$                                    |
| $F$   | $F$ | $F$                         | $F$                          | $F$                       | $T$                              | $T$                                    |
|       |     | " $p$ and $q$ "             | " $p$ xor $q$ "              | " $p$ or $q$ "            | "if $p$ then $q$ "               | " $p$ if and only if $q$ "             |

The only way to make the conditional statement  $p \rightarrow q$  false is to \_\_\_\_\_

The **hypothesis** of  $p \rightarrow q$  is \_\_\_\_\_ The **antecedent** of  $p \rightarrow q$  is \_\_\_\_\_

The **conclusion** of  $p \rightarrow q$  is \_\_\_\_\_ The **consequent** of  $p \rightarrow q$  is \_\_\_\_\_

The **converse** of  $p \rightarrow q$  is \_\_\_\_\_

The **inverse** of  $p \rightarrow q$  is \_\_\_\_\_

The **contrapositive** of  $p \rightarrow q$  is \_\_\_\_\_

We can use a recursive definition to describe all compound propositions that use propositional variables from a specified collection. Here's the definition for all compound propositions whose propositional variables are in  $\{p, q\}$ .

Basis Step:  $p$  and  $q$  are each a compound proposition  
 Recursive Step: If  $x$  is a compound proposition then so is  $(\neg x)$  and if  $x$  and  $y$  are both compound propositions then so is each of  $(x \wedge y), (x \oplus y), (x \vee y), (x \rightarrow y), (x \leftrightarrow y)$

Order of operations (Precedence) for logical operators:

Negation, then conjunction / disjunction, then conditional / biconditionals.

Example:  $\neg p \vee \neg q$  means  $(\neg p) \vee (\neg q)$ .

## (Some) logical equivalences

*Can replace  $p$  and  $q$  with any compound proposition*

$$\neg(\neg p) \equiv p$$

**Double negation**

$$p \vee q \equiv q \vee p$$

$$p \wedge q \equiv q \wedge p$$

**Commutativity** Ordering of terms

$$(p \vee q) \vee r \equiv p \vee (q \vee r)$$

$$(p \wedge q) \wedge r \equiv p \wedge (q \wedge r)$$

**Associativity** Grouping of terms

$$p \wedge F \equiv F \quad p \vee T \equiv T \quad p \wedge T \equiv p \quad p \vee F \equiv p$$

**Domination** aka short circuit evaluation

$$\neg(p \wedge q) \equiv \neg p \vee \neg q$$

$$\neg(p \vee q) \equiv \neg p \wedge \neg q$$

**DeMorgan's Laws**

$$p \rightarrow q \equiv \neg p \vee q$$

$$p \rightarrow q \equiv \neg q \rightarrow \neg p$$

**Contrapositive**

$$\neg(p \rightarrow q) \equiv p \wedge \neg q$$

$$\neg(p \leftrightarrow q) \equiv p \oplus q$$

$$p \leftrightarrow q \equiv q \leftrightarrow p$$

*Extra examples:*

$p \leftrightarrow q$  is not logically equivalent to  $p \wedge q$  because \_\_\_\_\_

$p \rightarrow q$  is not logically equivalent to  $q \rightarrow p$  because \_\_\_\_\_

## Common ways to express logical operators in English:

**Negation**  $\neg p$  can be said in English as

- Not  $p$ .
- It's not the case that  $p$ .
- $p$  is false.

**Conjunction**  $p \wedge q$  can be said in English as

- $p$  and  $q$ .
- Both  $p$  and  $q$  are true.
- $p$  but  $q$ .

**Exclusive or**  $p \oplus q$  can be said in English as

- $p$  or  $q$ , but not both.
- Exactly one of  $p$  and  $q$  is true.

**Disjunction**  $p \vee q$  can be said in English as

- $p$  or  $q$ , or both.
- $p$  or  $q$  (inclusive).
- At least one of  $p$  and  $q$  is true.

**Conditional**  $p \rightarrow q$  can be said in English as

- if  $p$ , then  $q$ .
- $p$  is sufficient for  $q$ .
- $q$  when  $p$ .
- $q$  whenever  $p$ .
- $p$  implies  $q$ .
- $q$  follows from  $p$ .
- $p$  is sufficient for  $q$ .
- $q$  is necessary for  $p$ .
- $p$  only if  $q$ .

**Biconditional**

- $p$  if and only if  $q$ .
- $p$  iff  $q$ .
- If  $p$  then  $q$ , and conversely.
- $p$  is necessary and sufficient for  $q$ .

**Translation:** Express each of the following sentences as compound propositions, using the given propositions.

|                                                        |                                                       |
|--------------------------------------------------------|-------------------------------------------------------|
| “A sufficient condition for the warranty to be good is | $w$ is “the warranty is good”                         |
| that you bought the computer less than a year ago”     | $b$ is “you bought the computer less than a year ago” |

|                                                                                    |                                                      |
|------------------------------------------------------------------------------------|------------------------------------------------------|
| “Whenever the message was sent from an unknown system, it is scanned for viruses.” | $s$ is “The message is scanned for viruses”          |
|                                                                                    | $u$ is “The message was sent from an unknown system” |

|                                                                            |                                                                                    |
|----------------------------------------------------------------------------|------------------------------------------------------------------------------------|
| “I will complete my to-do list only if I put a reminder<br>in my calendar” | $d$ is “I will complete my to-do list”<br>$c$ is “I put a reminder in my calendar” |
|----------------------------------------------------------------------------|------------------------------------------------------------------------------------|

**Definition:** A collection of compound propositions is called **consistent** if there is an assignment of truth values to the propositional variables that makes each of the compound propositions true.

**Consistency:**

Whenever the system software is being upgraded, users cannot access the file system. If users can access the file system, then they can save new files. If users cannot save new files, then the system software is not being upgraded.

1. Translate to symbolic compound propositions
2. Look for some truth assignment to the propositional variables for which all the compound propositions output  $T$

## Week 4 Wednesday: Predicates and Quantifiers

**Definition:** A **predicate** is a function from a given set (domain) to  $\{T, F\}$ .

A predicate can be applied, or **evaluated** at, an element of the domain.

Usually, a predicate *describes a property* that domain elements may or may not have.

Two predicates over the same domain are **equivalent** means they evaluate to the same truth values for all possible assignments of domain elements to the input. In other words, they are equivalent means that they are equal as functions.

To define a predicate, we must specify its domain and its value at each domain element. The rule assigning truth values to domain elements can be specified using a formula, English description, in a table (if the domain is finite), or recursively (if the domain is recursively defined).

| Input<br>$x$ | Output                     |                            |              |
|--------------|----------------------------|----------------------------|--------------|
|              | $V(x)$<br>$[x]_{2c,3} > 0$ | $N(x)$<br>$[x]_{2c,3} < 0$ | $Mystery(x)$ |
| 000          | $F$                        |                            | $T$          |
| 001          | $T$                        |                            | $T$          |
| 010          | $T$                        |                            | $T$          |
| 011          | $T$                        |                            | $F$          |
| 100          | $F$                        |                            | $F$          |
| 101          | $F$                        |                            | $T$          |
| 110          | $F$                        |                            | $F$          |
| 111          | $F$                        |                            | $T$          |

The domain for each of the predicates  $V(x)$ ,  $N(x)$ ,  $Mystery(x)$  is \_\_\_\_\_.

Fill in the table of values for the predicate  $N(x)$  based on the formula given.

**Definition:** The **truth set** of a predicate is the collection of all elements in its domain where the predicate evaluates to  $T$ .

Notice that specifying the domain and the truth set is sufficient for defining a predicate.

The truth set for the predicate  $V(x)$  is \_\_\_\_\_.

The truth set for the predicate  $N(x)$  is \_\_\_\_\_.

The truth set for the predicate  $Mystery(x)$  is \_\_\_\_\_.

The **universal quantification** of predicate  $P(x)$  over domain  $U$  is the statement “ $P(x)$  for all values of  $x$  in the domain  $U$ ” and is written  $\forall x P(x)$  or  $\forall x \in U P(x)$ . When the domain is finite, universal quantification over the domain is equivalent to iterated *conjunction* (ands).

The **existential quantification** of predicate  $P(x)$  over domain  $U$  is the statement “There exists an element  $x$  in the domain  $U$  such that  $P(x)$ ” and is written  $\exists x P(x)$  for  $\exists x \in U P(x)$ . When the domain is finite, existential quantification over the domain is equivalent to iterated *disjunction* (ors).

An element for which  $P(x) = F$  is called a **counterexample** of  $\forall x P(x)$ .

An element for which  $P(x) = T$  is called a **witness** of  $\exists x P(x)$ .

Statements involving predicates and quantifiers are **logically equivalent** means they have the same truth value no matter which predicates (domains and functions) are substituted in.

**Quantifier version of De Morgan’s laws:**  $\neg \forall x P(x) \equiv \exists x (\neg P(x))$   $\neg \exists x Q(x) \equiv \forall x (\neg Q(x))$

Examples of quantifications using  $V(x), N(x), Mystery(x)$ :

**True or False:**  $\exists x ( V(x) \wedge N(x) )$

**True or False:**  $\forall x ( V(x) \rightarrow N(x) )$

**True or False:**  $\exists x ( N(x) \leftrightarrow Mystery(x) )$

Rewrite  $\neg \forall x ( V(x) \oplus Mystery(x) )$  into a logical equivalent statement.

Notice that these are examples where the predicates have *finite* domain. How would we evaluate quantifications where the domain may be infinite?



*Recall the definitions:* The set of RNA strands  $S$  is defined (recursively) by:

Basis Step:  $\mathbf{A} \in S, \mathbf{C} \in S, \mathbf{U} \in S, \mathbf{G} \in S$   
 Recursive Step: If  $s \in S$  and  $b \in B$ , then  $sb \in S$

where  $sb$  is string concatenation.

The function  $rnalen$  that computes the length of RNA strands in  $S$  is defined recursively by:

$rnalen : S \rightarrow \mathbb{Z}^+$   
 Basis Step: If  $b \in B$  then  $rnalen(b) = 1$   
 Recursive Step: If  $s \in S$  and  $b \in B$ , then  $rnalen(sb) = 1 + rnalen(s)$

The function  $basecount$  that computes the number of a given base  $b$  appearing in a RNA strand  $s$  is defined recursively by:

$basecount : S \times B \rightarrow \mathbb{N}$   
 Basis Step: If  $b_1 \in B, b_2 \in B$   $basecount( (b_1, b_2) ) = \begin{cases} 1 & \text{when } b_1 = b_2 \\ 0 & \text{when } b_1 \neq b_2 \end{cases}$   
 Recursive Step: If  $s \in S, b_1 \in B, b_2 \in B$   $basecount( (sb_1, b_2) ) = \begin{cases} 1 + basecount( (s, b_2) ) & \text{when } b_1 = b_2 \\ basecount( (s, b_2) ) & \text{when } b_1 \neq b_2 \end{cases}$

### Example predicates on $S$ , the set of RNA strands (an infinite set)

$H : S \rightarrow \{T, F\}$  where  $H(s) = T$  for all  $s$ .

Truth set of  $H$  is \_\_\_\_\_

$L_A : S \rightarrow \{T, F\}$  defined recursively by:

Basis step:  $L_A(\mathbf{A}) = T, L_A(\mathbf{C}) = L_A(\mathbf{G}) = L_A(\mathbf{U}) = F$

Recursive step: If  $s \in S$  and  $b \in B$ , then  $L_A(sb) = L_A(s)$ .

Example where  $L_A$  evaluates to  $T$  is \_\_\_\_\_

Example where  $L_A$  evaluates to  $F$  is \_\_\_\_\_

## Using functions to define predicates:

$L$  with domain  $S \times \mathbb{Z}^+$  is defined by, for  $s \in S$  and  $n \in \mathbb{Z}^+$ ,

$$L( (s, n) ) = \begin{cases} T & \text{if } \text{rntalen}(s) = n \\ F & \text{otherwise} \end{cases}$$

In other words,  $L( (s, n) )$  means  $\text{rntalen}(s) = n$

$BC$  with domain  $S \times B \times \mathbb{N}$  is defined by, for  $s \in S$  and  $b \in B$  and  $n \in \mathbb{N}$ ,

$$BC( (s, b, n) ) = \begin{cases} T & \text{if } \text{basecount}( (s, b) ) = n \\ F & \text{otherwise} \end{cases}$$

In other words,  $BC( (s, b, n) )$  means  $\text{basecount}( (s, b) ) = n$

Example where  $L$  evaluates to  $T$ : \_\_\_\_\_ Why?

Example where  $BC$  evaluates to  $T$ : \_\_\_\_\_ Why?

Example where  $L$  evaluates to  $F$ : \_\_\_\_\_ Why?

Example where  $BC$  evaluates to  $F$ : \_\_\_\_\_ Why?

$$\exists t \ BC(t) \qquad \exists (s, b, n) \in S \times B \times \mathbb{N} \ (\text{basecount}( (s, b) ) = n)$$

In English:

Witness that proves this existential quantification is true:

$$\forall t \ BC(t) \qquad \forall (s, b, n) \in S \times B \times \mathbb{N} \ (\text{basecount}( (s, b) ) = n)$$

In English:

Counterexample that proves this universal quantification is false:

# Week 4 Friday: Evaluating Nested Quantifiers

## New predicates from old

1. Define the **new** predicate with domain  $S \times B$  and rule

$$\text{basecount}( (s, b) ) = 3$$

Example domain element where predicate is  $T$ :

2. Define the **new** predicate with domain  $S \times \mathbb{N}$  and rule

$$\text{basecount}( (s, \mathbf{A}) ) = n$$

Example domain element where predicate is  $T$ :

3. Define the **new** predicate with domain  $S \times B$  and rule

$$\exists n \in \mathbb{N} (\text{basecount}( (s, b) ) = n)$$

Example domain element where predicate is  $T$ :

4. Define the **new** predicate with domain  $S$  and rule

$$\forall b \in B (\text{basecount}( (s, b) ) = 1)$$

Example domain element where predicate is  $T$ :

**Notation:** for a predicate  $P$  with domain  $X_1 \times \cdots \times X_n$  and a  $n$ -tuple  $(x_1, \dots, x_n)$  with each  $x_i \in X$ , we can write  $P(x_1, \dots, x_n)$  to mean  $P( (x_1, \dots, x_n) )$ .

## Nested quantifiers

$$\forall s \in S \forall b \in B \forall n \in \mathbb{N} (\text{basecount}( (s, b) ) = n)$$

In English:

Counterexample that proves this universal quantification is false:

$$\forall n \in \mathbb{N} \forall s \in S \forall b \in B (\text{basecount}( (s, b) ) = n)$$

In English:

Counterexample that proves this universal quantification is false:

## Alternating nested quantifiers

$$\forall s \in S \exists b \in B ( \text{basecount}( (s, b) ) = 3 )$$

In English: For each RNA strand there is a base that occurs 3 times in this strand.

Write the negation and use De Morgan's law to find a logically equivalent version where the negation is applied only to the  $BC$  predicate (not next to a quantifier).

Is the original statement **True** or **False**?

$$\exists s \in S \forall b \in B \exists n \in \mathbb{N} ( \text{basecount}( (s, b) ) = n )$$

In English: There is an RNA strand so that for each base there is some nonnegative integer that counts the number of occurrences of that base in this strand.

Write the negation and use De Morgan's law to find a logically equivalent version where the negation is applied only to the  $BC$  predicate (not next to a quantifier).

Is the original statement **True** or **False**?