
# **Problem Set 4 Exercise #19: Fraction**

Reference: Week 11 notes

**Learning objective:** Structures

Estimated completion time: 60 minutes



### **Problem statement:**

[Modified from CS1020 assignment]

Write **fraction.c** to

- (1) Create a **fraction\_t** structure type that contains 2 integer members: **numerator** and **denominator**.
- (2) (in the main function) read in values of two **fraction\_t** variables one by one. You may assume that **denominator** is always positive.
- (3) write a function **simplify()** that returns the simplified form of a **fraction\_t** variable. For example, the simplified form of  $\frac{9}{12}$  is  $\frac{3}{4}$ .
- (4) write a function **equals()** that takes two **fraction\_t** variables, checks whether they have the same value. This function returns 1 if so or 0 otherwise.
- (5) write a function add() that takes two fraction\_t variables, adds them and returns the result in the simplified form.

You may want to calculate the greatest common divisor (GCD) when simplifying a fraction.

### Sample run #1:

Enter 1st fraction: 2 20
Enter 2nd fraction: 3 30
1st fraction is 2/20
2nd fraction is 3/30
Two fractions are the same.
2/20 + 3/30 = 1/5

#### Sample run #2:

Enter 1st fraction: 0 8
Enter 2nd fraction: 6 14
1st fraction is 0/8
2nd fraction is 6/14
Two fractions are not the same.
0/8 + 6/14 = 3/7

# Sample run #3:

```
Enter 1st fraction: 1 10
Enter 2nd fraction: -5 10
1st fraction is 1/10
2nd fraction is -5/10
Two fractions are not the same.
1/10 + -5/10 = -2/5
```

# Sample run #4:

```
Enter 1st fraction: -2 5
Enter 2nd fraction: 14 6
1st fraction is -2/5
2nd fraction is 14/6
Two fractions are not the same.
-2/5 + 14/6 = 29/15
```