

STAT 4010 Bayesian Learning

TUTORIAL 10

Spring 2023

Cheuk Hin (Andy) CHENG ([Email](#) | [Homepage](#))

Di SU ([Email](#) | [Homepage](#))

1 Gibbs Sampler

We want to draw d -dimensional sample $\theta_j = (\theta_{j1}, \dots, \theta_{jd})^T$ from $\pi(\theta)$.

Algorithm 1: Gibbs Sampler

Input: (i) number of iteration J ; (ii) conditional PDF $\pi^{(k|-k)}(\cdot | \theta_{-k})$ for $k = 1, \dots, d$; and (iii) initialization PDF $\pi_{\text{initial}}(\cdot)$.

begin

 (1) Generate $\theta_0 \sim \pi_{\text{initial}}(\cdot)$

 (2) Set $\vartheta \leftarrow \theta_0$

 (3) **for** j in $\{1, \dots, J\}$ **do**

for k in $\{1, \dots, d\}$ **do**

 Generate $\theta_{jk} \sim \pi^{(k|-k)}(\cdot | \vartheta_{-k})$

 Update the k th component of ϑ as $\vartheta_k \leftarrow \theta_{jk}$.

end

end

end

Output: $\theta_{1:J}$

Theorem 1.1. *Gibbs sampler is a composition of d MH algorithm with acceptance probabilities in each step always equals to 1.*

When the full conditionals are not easy to sample from, we can make use of MH algorithm.

Algorithm 2: MH-within-Gibbs Sampler

Input: (i) number of iteration J ; (ii) proposal PDF $p^{(k)}(\theta_k | \theta_{-k})$ for $k = 1, \dots, d$; (iii) conditional PDF $\pi_u^{(k|-k)}(\cdot | \theta_{-k})$ for $k = 1, \dots, d$; and (iv) initialization PDF $\pi_{\text{initial}}(\cdot)$.

begin

(1) Generate $\theta_0 \sim \pi_{\text{initial}}(\cdot)$

(2) Set $\vartheta \leftarrow \theta_0$

(3) **for** j in $\{1, \dots, J\}$ **do**

for k in $\{1, \dots, d\}$ **do**

 Generate $\tilde{\theta}_{jk} \sim p^{(k)}(\cdot | \vartheta)$

 Generate $U_{jk} \sim \text{Unif}(0, 1)$

 Compute the acceptance probability

$$a_{jk} = \min \left\{ 1, \frac{\pi_u^{(k|-k)}(\tilde{\theta}_{jk} | \vartheta_{-k}) p_u^{(k)}(\theta_{j-1,k} | \tilde{\theta}_{jk}, \vartheta_{-k})}{\pi_u^{(k|-k)}(\theta_{j-1,k} | \vartheta_{-k}) p_u^{(k)}(\tilde{\theta}_{jk} | \theta_{j-1,k}, \vartheta_{-k})} \right\}$$

 Compute $\theta_{jk} = \tilde{\theta}_{jk} \mathbb{1}(U_{jk} \leq a_{jk}) + \theta_{j-1,k} \mathbb{1}(U_{jk} > a_{jk})$

 Update the k th component of ϑ as $\vartheta_k \leftarrow \theta_{jk}$

end

end

end

Output: $\theta_{1:J}$

Remark 1.1. Notice that the unnormalized density $\pi_u^{(k|-k)}(\tilde{\theta}_{jk} | \vartheta_{-k})$ is not conditioned on $\theta_{j-1,k}$.

2 Examples

Example 2.1. (Exercise 6.2 A6 2021) Let x_1, \dots, x_n be the numbers of reported COVID-19 cases in Hong Kong from 1 February 2020 to 10 April 2020 (i.e., $n = 70$), respectively. The dataset can be downloaded from the HKSAR government dataset ([click here](#)). Some people believe that the distribution of $x_1, \dots, x_{\tau-1}$ is different from that of x_{τ}, \dots, x_n for some $2 \leq \tau \leq n$. Suppose that the data are modeled by a change-point model as follows:

$$\begin{aligned} [x_i | \tau, \theta_1, \theta_2] &\stackrel{\text{iid}}{\sim} \begin{cases} \text{Po}(\theta_1) & \text{if } i = 1, \dots, \tau - 1; \\ \text{Po}(\theta_2) & \text{if } i = \tau, \dots, n; \end{cases} \\ \theta_1, \theta_2 &\stackrel{\text{iid}}{\sim} \text{Ga}(\alpha)/\beta \\ \tau &\sim \text{Unif}\{2, \dots, n\}, \end{aligned}$$

where $\alpha, \beta > 0$ are non-random and are suitably chosen by you. The goals of this exercise are (i) to perform statistical inference on τ, θ_1, θ_2 ; and (ii) to learn from data and give statistically grounded suggestions.

1. Derive the conditional densities of $[\tau \mid \theta_1, \theta_2, x_{1:n}]$, $[\theta_1 \mid \tau, \theta_2, x_{1:n}]$, and $[\theta_2 \mid \tau, \theta_1, x_{1:n}]$.
2. Use a suitable MCMC method to draw posterior samples of τ, θ_1, θ_2 with $J = 2^{13}$ iterations. Discard the first half as burn-in.
3. Visualize your MCMC sample produced in part 2. Comment the quality of your MCMC sample.

SOLUTION:

1. Note that

$$\begin{aligned} f(\tau \mid \theta_1, \theta_2, x_{1:n}) &\propto f(x_{1:n} \mid \tau, \theta_1, \theta_2) f(\tau \mid \theta_1, \theta_2) \\ &\propto \left(\prod_{i=1}^{\tau-1} e^{-\theta_1 \theta_2^{x_i}} \right) \left(\prod_{i=\tau}^n e^{-\theta_2 \theta_2^{x_i}} \right) \mathbb{1}(\tau \in \{2, \dots, n\}) \\ &= \exp \left\{ -\tau \theta_1 - (n - \tau) \theta_2 + \left(\sum_{i=1}^{\tau-1} x_i \right) \ln \theta_1 + \left(\sum_{i=\tau}^n x_i \right) \ln \theta_2 \right\} \mathbb{1}(\tau \in \{2, \dots, n\}) \end{aligned}$$

Similarly, we have

$$\begin{aligned} f(\theta_1 \mid \tau, \theta_2, x_{1:n}) &\propto f(x_{1:n} \mid \tau, \theta_1, \theta_2) f(\theta_1 \mid \tau, \theta_2) \\ &\propto \left(\prod_{i=1}^{\tau-1} e^{-\theta_1 \theta_2^{x_i}} \right) \theta_1^{\alpha-1} e^{-\beta \theta_1} \mathbb{1}(\theta_1 > 0) \\ &\sim \text{Ga} \left(\alpha + \sum_{i=1}^{\tau-1} x_i, \beta + \tau \right) \quad \text{and} \\ f(\theta_2 \mid \tau, \theta_1, x_{1:n}) &\sim \text{Ga} \left(\alpha + \sum_{i=\tau}^n x_i, \beta + n - \tau \right) \end{aligned}$$

2. Since we have derived all conditional densities, we can use the Gibbs sampler. While $f(\tau \mid \theta_1, \theta_2, x_{1:n})$ is not a named distribution, note that it is a PMF and so we can use the sample function. For simplicity, we take $\alpha = \beta = 1$.

```

1 # Import and extract data
2 data = read.csv("enhanced_sur_covid_19_eng.csv")
3 t0 = as.numeric(as.Date("01/02/2020", "%d/%m/%Y"))
4 t1 = as.numeric(as.Date("10/04/2020", "%d/%m/%Y"))
5 x = t = rep(NA, t1-t0+1)
6 for (i in t0:t1) {
7   x[i-t0+1] = sum(as.Date(data$Report.date, "%d/%m/%Y") == i)
8   t[i-t0+1] = as.character(as.Date("01/02/2020", "%d/%m/%Y") + i-t0)
9 }
10 names(x) = t
11
12 # Ex6.2.2: sample from posterior
13 sim_tau <- function(theta1, theta2, x, n) {
14   tau = 2:n
15   cs = cumsum(x)
16   lp = -tau*theta1 - (n-tau)*theta2 + cs[1:(n-1)]*log(theta1) + (sum(x)-cs
17     [1:(n-1)])*log(theta2)
18   p = exp(lp-max(lp))

```

```

18   sample(tau, 1, prob=p/sum(p))
19 }
20
21 ##initial proposal
22 #tau ~ Unif{2,..,n}
23 #theta1, theta2 ~iid Ga(1)/1
24
25 gibbs21 <- function(J,x,burn_frac = 0.5, keep = F, a = 1,b = 1){
26   nsim = ceiling(J/burn_frac)+1
27   n = length(x)
28   theta = array(NA,c(nsim,3))
29   colnames(theta) = c('tau','theta1','theta2')
30   ##sample from the initial proposal
31   theta[1,] = c(sample(2:n,1),rgamma(2,a,b))
32   for (j in 2:nsim) {
33     theta_use = theta[j-1,] ##current info
34     ##Update tau
35     theta_use[1] = sim_tau(theta_use[2],theta_use[3],x,n)
36     ##Update theta1
37     theta_use[2] = rgamma(1, a+sum(x[1:(theta_use[1]-1)]), b+theta_use
38       [1])
39     ##Update theta2
40     theta_use[3] = rgamma(1, a+sum(x[theta_use[1]:n]), b+n-theta_use[1])
41     ##update theta
42     theta[j,] = theta_use
43   }
44   if (keep) {
45     return(theta)
46   }else{
47     return(tail(theta,J)) ##burn the first fraction
48   }
49 }
50
51
52 J_sim = 2^13
53 theta_name = c('tau','theta1','theta2')
54 theta_sim = gibbs21(J_sim,x,keep=TRUE)
55 theta_sim = theta_sim[-1,]##remove initialization
56 iUse = (J_sim/2+1):J_sim

```

3. The plots show that the MCMC sample produced in part 2 looks stationary and converge quickly. We can further compare the auto-correlation plots which shows that the burn-in has effectively reduced the stickiness of the chain as the cross-correlations are eliminated (we omit the 3-by-3 auto-correlation plots here for compactness).

```


1 transCol = function(color, percent=50) {
2   v = col2rgb(color)
3   newCol = rgb(v[1],v[2],v[3],max=255,alpha=(1-percent/100)*255)
4   invisible(newCol)
5 }
6 par(mfrow=c(1,2), mar=c(4.5,5,3,2))
7 col = c(transCol("red", percent=50),
8         transCol("blue", percent=50),
9         transCol("green", percent=50))
10 matplot(1:200, theta_sim[1:200,], col=col, lwd=2, type="l", lty=1,
11         ylab="", xlab="Time", main="First 200 samples")
12 legend("topright", c("tau", "theta1", "theta2"), col=col, pch=15, cex=1.2)

```

```

13 matplot(iUse, theta_sim[iUse,], col=col, lwd=2, type="l", lty=1,
14         ylab="", xlab="Time", main="Used sample")
15 acf(theta_sim, mar=c(3,2.5,2,0.5))
16 acf(theta_sim[iUse,], mar=c(3,2.5,2,0.5))
17
18 ##MCMC Diagnostic
19 ##1) Density plot
20 windows(height=15,width = 15)
21 par(mfrow = c(3,3))
22 for (i in 1:3) {
23   plot(density(theta_sim[,i], kernel="epanechnikov"),main = paste('
24     Density plot for',theta_name[i]),
25         ylab = 'Density',xlab = expression(theta))
26 }
27 ##2) ACF plot
28 for (i in 1:3) {
29   acf = acf(theta_sim[,i],plot=F)
30   plot(acf,main = paste('ACF of',theta_name[i]),xlab = 'Lag')
31 }
32 ##3) Trace plot
33 for (i in 1:3) {
34   plot((J_sim-500+1):J_sim,tail(theta_sim[,i],500),type = 'l',
35        main = paste('Trace plot of',theta_name[i]),ylab = expression(
36          theta),xlab = 'j')
37 }
38 ##Cp analysis
39 cp = round(mean(theta_sim[,1]),0)
40 cp
41 n = length(x)
42 plot(1:n,x,type='l')
43 abline(v=cp,col = 'green')

```



Example 2.2 (Option pricing under double exponential model). Consider the following

model,

$$\begin{aligned} r_{1:n} \mid \lambda_0, \lambda_1, p &\stackrel{\text{iid}}{\sim} f(r \mid \lambda_0, \lambda_1, p) = p \frac{1}{\lambda_1} e^{-r/\lambda_1} \mathbb{1}(r \geq 0) + (1-p) \frac{1}{\lambda_0} e^{r/\lambda_0} \mathbb{1}(r < 0) \\ \lambda_0, \lambda_1 &\stackrel{\text{iid}}{\sim} \text{InvGamma}(\mu = 5\%, \sigma = 1\%) \\ p &\sim \text{Beta}(\mu = 0.5, \sigma = 0.1), \end{aligned}$$

where r can be thought of log return for each unit of time. Note that the common representations for the parameters of Inverse-Gamma and Beta distribution are

$$\begin{aligned} \text{InvGamma}(\mu, \sigma) &= k/Ga(h) \\ h &= \mu^2/\sigma^2 + 2 \\ k &= \mu(h-1) \\ \text{Beta}(\mu, \sigma) &= \text{Beta}(\alpha, \beta) \\ \alpha &= \frac{\mu^2(1-\mu)}{\sigma^2} - \mu \\ \beta &= \alpha(1/\mu - 1). \end{aligned}$$

Define $P_n = \sum_{i:r_i \geq 0} r_i$, $N_n = \sum_{i:r_i < 0} r_i$, n_1 be the number of positive $r_{1:n}$ and $n_0 = n - n_1$. From the data, we have $n = 100$, $n_1 = 55$, $P_n = 2.2$ and $N_n = -2.7$. Let $S_0 = 373$ and $K = 380$. Using MH-within-Gibbs sampler, estimate

$$\mathbb{E}[\max(S_T - k, 0) \mid r_{1:n}] = \mathbb{E}[\max(S_0 e^{r_{n+1}} - k, 0) \mid r_{1:n}].$$

SOLUTION: Note that the joint sampling distribution is

$$f(r_{1:n} \mid \lambda_0, \lambda_1, p) = p^{n_1} (1-p)^{n_0} \lambda_1^{-n_1} \lambda_0^{-n_0} e^{-P_n/\lambda_1} e^{N_n/\lambda_0}.$$

We first find the conditional distributions for the parameters.

$$\begin{aligned} f(\lambda_1 \mid r_{1:n}, \lambda_0, p) &\propto f(\lambda_1 \mid \lambda_0, p) f(r_{1:n} \mid \lambda_1, \lambda_0, p) \\ &= f(\lambda_1) f(r_{1:n} \mid \lambda_1, \lambda_0, p) \\ &\propto \lambda_1^{-h-1} e^{-k/\lambda_1} [p^{n_1} (1-p)^{n_0} \lambda_1^{-n_1} \lambda_0^{-n_0} e^{N_n/\lambda_1}] \\ &= p^{n_1} (1-p)^{n_0} \lambda_1^{-h-n_1-1} \lambda_0^{-n_0} e^{-(P_n+k)/\lambda_1} e^{N_n/\lambda_0} \mathbb{1}(\lambda_1 > 0). \end{aligned}$$

Similarly,

$$f(\lambda_0 \mid r_{1:n}, \lambda_1, p) \propto p^{n_1} (1-p)^{n_0} \lambda_1^{-n_1} \lambda_0^{-h-n_0-1} e^{-P_n/\lambda_1} e^{-(k-N_n)/\lambda_0} \mathbb{1}(\lambda_0 > 0).$$

Moreover,

$$f(p \mid r_{1:n}, \lambda_1, \lambda_0) \propto p^{n_1+\alpha-1} (1-p)^{n_0+\beta-1} \lambda_1^{-n_1} \lambda_0^{-n_0} e^{-P_n/\lambda_1} e^{N_n/\lambda_0} \mathbb{1}(p \in (0, 1)).$$

The discussion for the proposals are as followed.

- The conditional densities for λ_1 and λ_0 are proportional to inverse-gamma kernel. Therefore, we would set $\tilde{\lambda} \mid \lambda \sim \text{InvGamma}(\mu = \lambda, \sigma = 0.01)$.
- The conditional density for p is proportional to beta kernel. Therefore, we would set $\tilde{p} \mid p \sim \text{Beta}(\mu = p, \sigma = 0.05)$.

Note that the conditional distributions are not exactly the commonly known distributions. We implement the MH algorithm to generate samples.

```

1 library(invgamma)
2
3 log_pi_lambda1 <- function(Pn,Nn,n1,n0,lambda1,lambda0,p,h,k){
4   n1*log(p)+n0*log(1-p)-(h+n1+1)*log(lambda1)-n0*log(lambda0)-(Pn+k)/lambda1+
5     Nn/lambda0
6 }
7 log_pi_lambda0 <- function(Pn,Nn,n1,n0,lambda1,lambda0,p,h,k){
8   n1*log(p)+n0*log(1-p)-n1*log(lambda1)-(h+n0+1)*log(lambda0)-Pn/lambda1-(k-Nn
9     )/lambda0
10 }
11 log_pi_p <- function(Pn,Nn,n1,n0,lambda1,lambda0,p,a,b){
12   (n1+a-1)*log(p)+(n0+b-1)*log(1-p)-n1*log(lambda1)-n0*log(lambda0)-Pn/lambda1
13     +Nn/lambda0
14 }
15 get_invgamma_para <- function(mu,sd){
16   h = mu^2/sd^2+2
17   k = mu*(h-1)
18   c(h,k)
19 }
20
21 get_beta_para <- function(mu,sd){
22   a = mu^2*(1-mu)/sd^2-mu
23   b = a*(1/mu-1)
24   c(a,b)
25 }
26
27 ##MH
28 MH_lambda1 <- function(Pn,Nn,n1,n0,lambda1,lambda0,p,sd_lambda1 = 0.01){
29   ##represent the parameters
30   ##parameters for numerator in proposal odd
31   para = get_invgamma_para(lambda1,sd_lambda1)
32   h = para[1]
33   k = para[2]
34
35   lambda1_p = rinvgamma(1,h,k)
36   log_target_odd = log_pi_lambda1(Pn,Nn,n1,n0,lambda1 = lambda1_p,lambda0,p,h,
37     k)-log_pi_lambda1(Pn,Nn,n1,n0,lambda1 = lambda1,lambda0,p,h,k)
38
39   ##parameters for denominator in proposal odd
40   para = get_invgamma_para(lambda1_p,sd_lambda1)
41   h_d = para[1]
42   k_d = para[2]
43   log_proposal_odd = log(dinvgamma(lambda1_p,h,k)) - log(dinvgamma(lambda1,h_d
44     ,k_d))
45   accept_prob = exp(min(0,log_target_odd-log_proposal_odd))
46   u = runif(1)
47   lambda1_p*(u<=accept_prob)+lambda1*(u>accept_prob)
48 }
49
50 MH_lambda0 <- function(Pn,Nn,n1,n0,lambda1,lambda0,p,sd_lambda0 = 0.01){
51   para = get_invgamma_para(lambda0,sd_lambda0)
52   h = para[1]
53   k = para[2]

```

```

52 lambda0_p = rinvgamma(1,h,k)
53 log_target_odd = log_pi_lambda0(Pn,Nn,n1,n0,lambda1,lambda0 = lambda0_p,p,h,
    k)-log_pi_lambda0(Pn,Nn,n1,n0,lambda1,lambda0,p,h,k)
54 para = get_invgamma_para(lambda0_p,sd_lambda0)
55 h_d = para[1]
56 k_d = para[2]
57 log_proposal_odd = log(dinvgamma(lambda0_p,h,k)) - log(dinvgamma(lambda0,h_d
    ,k_d))
58 accept_prob = exp(min(0,log_target_odd-log_proposal_odd))
59 u = runif(1)
60 lambda0_p*(u<=accept_prob)+lambda0*(u>accept_prob)
61 }
62
63 MH_p <- function(Pn,Nn,n1,n0,lambda1,lambda0,p,sd_p = 0.1){
64   para = get_beta_para(p,sd_p)
65   a = para[1]
66   b = para[2]
67   p_p = rbeta(1,a,b)
68   log_target_odd = log_pi_p(Pn,Nn,n1,n0,lambda1,lambda0,p = p_p,a,b)-log_pi_p(
    Pn,Nn,n1,n0,lambda1,lambda0,p,a,b)
69   para = get_beta_para(p_p,sd_p)
70   a_d = para[1]
71   b_d = para[2]
72   log_proposal_odd = log(dbeta(p_p,a,b)) - log(dbeta(p,a_d,b_d))
73   accept_prob = exp(min(0,log_target_odd-log_proposal_odd))
74   u = runif(1)
75   p_p*(u<=accept_prob)+p*(u>accept_prob)
76 }

```

The Gibbs sampler can be implemented as followed.

```

1 gibbs <- function(J,Pn,Nn,n1,n0,burn_frac = 0.5, keep = F, ##general para
    representing Gibbs
2     sd_lambda1 = 0.01,sd_lambda0 = 0.01,sd_p = 0.1, ##para for
    step size
3     mu_lambda_ini = 0.05,sd_lambda_ini = 0.01, ##para for
    initial proposal
4     mu_p_ini = 0.5,sd_p_ini = 0.1){
5   nsim = ceiling(J/burn_frac)+1
6   theta = array(NA,c(nsim,3))
7   colnames(theta) = c('lambda1','lambda0','p')
8   ##sample from the initial proposal
9   para = get_invgamma_para(mu_lambda_ini,sd_lambda_ini)
10  h = para[1]
11  k = para[2]
12  para = get_beta_para(mu_p_ini,sd_p_ini)
13  a = para[1]
14  b = para[2]
15  theta[1,] = c(rinvgamma(2,h,k),rbeta(1,a,b))
16  for (j in 2:nsim) {
17    theta_use = theta[j-1,]
18    ##update lambda1
19    theta_use[1] = MH_lambda1(Pn,Nn,n1,n0,
20        theta_use[1],theta_use[2],theta_use[3],sd_
        lambda1)
21    ##update lambda0
22    theta_use[2] = MH_lambda0(Pn,Nn,n1,n0,
23        theta_use[1],theta_use[2],theta_use[3],sd_
        lambda0)

```



```

24     ##update p
25     theta_use[3] = MH_p(Pn,Nn,n1,n0,
26                         theta_use[1],theta_use[2],theta_use[3],sd_p)
27     theta[j,] = theta_use
28 }
29 if (keep) {
30     return(theta)
31 }else{
32     return(tail(theta,J)) ##burn the first fraction
33 }
34 }
35
36 ##sample parameters from Gibbs
37 set.seed(410)
38 J_sim = 2^12
39 n1 = 55
40 n0 = 45
41 Pn = 2.2
42 Nn = -2.7
43 theta_sim = gibbs(J_sim,Pn,Nn,n1,n0,keep=F)
44 theta_name = c('lambda1','lambda0','p')
45 colnames(theta_sim) = theta_name


```

After running Gibbs, some diagnostic can be performed.

```

1  ##MCMC Diagnostic
2  ##1) Density plot
3  windows(height=15,width = 15)
4  par(mfrow = c(3,3))
5  for (i in 1:3) {
6      plot(density(theta_sim[,i], kernel="epanechnikov"),main = paste('Density
7          plot for',theta_name[i]),
8          ylab = 'Density',xlab = expression(theta))
9  }
10 ##2) ACF plot
11 for (i in 1:3) {
12     acf = acf(theta_sim[,i],plot=F)
13     plot(acf,main = paste('ACF of',theta_name[i]),xlab = 'Lag')
14 }
15 ##3) Trace plot
16 for (i in 1:3) {
17     plot((J_sim-500+1):J_sim,tail(theta_sim[,i],500),type = 'l',
18         main = paste('Trace plot of',theta_name[i]),ylab = expression(theta),
19         xlab = 'j')

```



Finally, use the posterior samples to generate posterior predictive samples and estimate the call price. Note that $\max(A, 0) = A \cdot \mathbb{1}(A > 0)$. The estimated option price is \$6.00.

```

1 ##generate posterior predictive samples
2 set.seed(4010)
3 u = rbinom(J_sim,1,theta_sim[, 'p'])
4 r = rexp(J_sim,1/theta_sim[, 'lambda1']) * u - rexp(J_sim,1/theta_sim[, 'lambda0']) *
  (1-u)
5 ##Compute option payoff
6 S0 = 373
7 K = 380
8 c = S0 * exp(r) - K
9 c = c * (c > 0)
10 mean(c)
11 [1] 5.99987

```