Derivation of Vieta's theorem for solving quadratic equations

Danil Ivlev

February 8, 2025

1 Introduction

Most solutions to quadratic equations rely on the application of the discriminant. Although this formula proves to be an effective tool for determining roots, the question arises: is there a simpler approach? This research proposes to use the derivation of Vieta's theorem as an alternative method and aims to provide a more accessible approach to solving quadratic equations.

2 Derivation

We have a system like this

$$\begin{cases} x_1 + x_2 = r_1 \\ x_1 x_2 = r_2 \end{cases}$$

Let $\frac{x_1}{x_2} = r_2$ be true, then

$$\begin{cases} x_1 + \frac{1}{x_2} = r_1 \\ \frac{x_1}{x_2} = r_2 \end{cases}$$

Now we need to express x_1 through x_2 , then x_1 will look like this

$$x_1 = \frac{r_2}{x_2}$$

We substitute this new expression in place of x_1 in the first expression of the system and solve the equation

$$r_{1} = \frac{r_{2} + 1}{x_{2}}$$

$$r_{1}x_{2} = r_{2} + 1$$

$$x_{2} = \frac{r_{2} + 1}{r_{1}}$$
(1)

Finally, to get the formula for x_1 , we divide r_2 by x_2 and raise x_2 to degree -1, yielding the following system of formulas (here x_1 and x_2 are swapped, but it makes no difference)

$$x_1 = \frac{r_1}{r_2 + 1}$$

$$x_2 = \frac{r_2}{x_1}$$
(2)

Now we can reduce these formulas, and we get the following

$$x_{1} = \frac{-\frac{b}{a}}{\frac{c+a}{a}} = (-\frac{b}{a})(\frac{a}{c+a}) = -\frac{b}{c+a}$$

$$x_{2} = \frac{\frac{c}{a}}{-\frac{b}{c+a}} = (\frac{c}{a})(-\frac{c+a}{b}) = -\frac{c(c+a)}{ab}$$
(3)

3 Conclusion

Thus, we were able to obtain two formulas that allow us to calculate the roots of a quadratic equation easily and simply. We hope that this work will inspire further research in this area and contribute to a better understanding of quadratic equations and their properties. We thank for attention to our work and encourage discussion of the results presented.