

Solutions: Linear Algebra

ECON 441: Introduction to Mathematical Economics

Instructor: Div Bhagia

Exercise 4.2

$$1. \ A = \begin{bmatrix} 7 & -1 \\ 6 & 9 \end{bmatrix} \quad B = \begin{bmatrix} 0 & 4 \\ 3 & -2 \end{bmatrix} \quad C = \begin{bmatrix} 8 & 3 \\ 6 & 1 \end{bmatrix}$$

$$(a) \ A + B = \begin{bmatrix} 7 & 3 \\ 9 & 7 \end{bmatrix}$$

$$(b) \ C - A = \begin{bmatrix} 1 & 4 \\ 0 & -8 \end{bmatrix}$$

$$(c) \ 3A = \begin{bmatrix} 21 & -3 \\ 18 & 27 \end{bmatrix}$$

$$(d) \ 4B + 2C = \begin{bmatrix} 0 & 16 \\ 12 & -8 \end{bmatrix} + \begin{bmatrix} 16 & 6 \\ 12 & 2 \end{bmatrix} = \begin{bmatrix} 16 & 22 \\ 24 & -6 \end{bmatrix}$$

$$2. \ A = \begin{bmatrix} 2 & 8 \\ 3 & 0 \\ 5 & 1 \end{bmatrix} \quad B = \begin{bmatrix} 2 & 0 \\ 3 & 8 \end{bmatrix} \quad C = \begin{bmatrix} 7 & 2 \\ 6 & 3 \end{bmatrix}$$

- (a) AB is defined as number of columns in A is two which is equal to the number of rows in B .

$$AB = \begin{bmatrix} 2 \times 2 + 8 \times 3 & 2 \times 0 + 8 \times 8 \\ 3 \times 2 + 0 \times 3 & 3 \times 0 + 0 \times 8 \\ 5 \times 2 + 1 \times 3 & 5 \times 0 + 1 \times 8 \end{bmatrix} = \begin{bmatrix} 28 & 64 \\ 6 & 0 \\ 13 & 8 \end{bmatrix}$$

Not possible to calculate BA as B has two columns, but A has three rows.

- (b) BC and CB are both defined as both have two rows and two columns.

$$BC = \begin{bmatrix} 14 & 4 \\ 69 & 30 \end{bmatrix} \neq CB = \begin{bmatrix} 20 & 16 \\ 21 & 24 \end{bmatrix}$$

4.

$$(a) \begin{bmatrix} 0 & 2 \\ 36 & 20 \\ 16 & 3 \end{bmatrix}_{3 \times 2}$$

$$(b) \begin{bmatrix} 49 & 3 \\ 4 & 3 \end{bmatrix}_{2 \times 2}$$

$$(c) \begin{bmatrix} 3x + 5y \\ 4x + 2y - 7z \end{bmatrix}_{2 \times 1}$$

$$(d) [7a + c \quad 2b + 4c]$$

Exercise 4.4

5. (e)

$$A = \begin{bmatrix} -2 \\ 4 \\ 7 \end{bmatrix}_{3 \times 1} \quad B = \begin{bmatrix} 3 & 6 & -2 \end{bmatrix}_{1 \times 3}$$

$$C = AB = \begin{bmatrix} -6 & -12 & 4 \\ 12 & 24 & -8 \\ 21 & 42 & -14 \end{bmatrix}_{3 \times 3}$$

$$D = BA = [3 \times -2 + 6 \times 4 + -2 \times 7] = [4]_{1 \times 1}$$

7. In example 5, $x = \begin{bmatrix} x_1 \\ x_2 \end{bmatrix}$ and $A = \begin{bmatrix} a_{11} & 0 \\ 0 & a_{22} \end{bmatrix}$. In which case,

$$\begin{aligned} x^T A x &= \begin{bmatrix} x_1 & x_2 \end{bmatrix} \begin{bmatrix} a_{11} & 0 \\ 0 & a_{22} \end{bmatrix} \begin{bmatrix} x_1 \\ x_2 \end{bmatrix}_{2 \times 1} \\ &= \begin{bmatrix} a_{11}x_1 & a_{22}x_2 \end{bmatrix}_{1 \times 2} \begin{bmatrix} x_1 \\ x_2 \end{bmatrix}_{2 \times 1} \\ &= \underbrace{a_{11}x_1^2 + a_{22}x_2^2}_{\text{Weighted sum of squares}} = \sum_{i=1}^2 a_{ii}x_i^2 \end{aligned}$$

So $x^T A x$ represents a weighted sum of squares where a_{11}, a_{22} are weights.

But now what if $A = \begin{bmatrix} a_{11} & a_{12} \\ a_{21} & a_{22} \end{bmatrix}$. In this case,

$$\begin{aligned}
x^T Ax &= \begin{bmatrix} x_1 & x_2 \end{bmatrix} \begin{bmatrix} a_{11} & a_{12} \\ a_{21} & a_{22} \end{bmatrix}_{2 \times 2} \begin{bmatrix} x_1 \\ x_2 \end{bmatrix}_{2 \times 1} \\
&= \begin{bmatrix} a_{11}x_1 + a_{21}x_2 & a_{12}x_1 + a_{22}x_2 \end{bmatrix} \begin{bmatrix} x_1 \\ x_2 \end{bmatrix}_{2 \times 1} \\
&= a_{11}x_1^2 + a_{21}x_1x_2 + a_{12}x_1x_2 + a_{22}x_2^2 \\
&= a_{11}x_1^2 + (a_{21} + a_{12})x_1x_2 + a_{22}x_2^2
\end{aligned}$$

So $x^T Ax$ no longer represents a weighted sum of squares.

You can check that the associative law i.e.

$$(x^T A)x = x^T (Ax)$$

will apply in both cases (after all, its a law!) as all products are possible.

Exercise 4.5

1.

$$A = \begin{bmatrix} -1 & 5 & 7 \\ 0 & -2 & 4 \end{bmatrix}_{2 \times 3} \quad b = \begin{bmatrix} 9 \\ 6 \\ 0 \end{bmatrix}_{3 \times 1} \quad x = \begin{bmatrix} x_1 \\ x_2 \end{bmatrix}_{2 \times 1}$$

$$(a) \quad AI = \begin{bmatrix} -1 & 5 & 7 \\ 0 & -2 & 4 \end{bmatrix} \begin{bmatrix} 1 & 0 & 0 \\ 0 & 1 & 0 \\ 0 & 0 & 1 \end{bmatrix}_{3 \times 3} = \begin{bmatrix} -1 & 5 & 7 \\ 0 & -2 & 4 \end{bmatrix} = A$$

$$(b) \quad IA = \begin{bmatrix} 1 & 0 \\ 0 & 1 \end{bmatrix}_{2 \times 2} \begin{bmatrix} -1 & 5 & 7 \\ 0 & -2 & 4 \end{bmatrix} = \begin{bmatrix} -1 & 5 & 7 \\ 0 & -2 & 4 \end{bmatrix}$$

$$(c) \quad Ix = \begin{bmatrix} 1 & 0 \\ 0 & 1 \end{bmatrix}_{2 \times 2} \begin{bmatrix} x_1 \\ x_2 \end{bmatrix} = \begin{bmatrix} x_1 \\ x_2 \end{bmatrix}$$

$$(d) \ x^T I = \begin{bmatrix} x_1 & x_2 \end{bmatrix} \begin{bmatrix} 1 & 0 \\ 0 & 1 \end{bmatrix}_{2 \times 2} = \begin{bmatrix} x_1 & x_2 \end{bmatrix}$$

4. Let's start with a 2×2 diagonal matrix

$$\begin{bmatrix} a_{11} & 0 \\ 0 & a_{22} \end{bmatrix} \begin{bmatrix} a_{11} & 0 \\ 0 & a_{22} \end{bmatrix} = \begin{bmatrix} a_{11}^2 & 0 \\ 0 & a_{22}^2 \end{bmatrix}$$

$x = x^2$ for only $x = 0, 1$ so a_{11} and a_{22} can either be 0 or 1. So we can have the following 2×2 idempotent diagonal matrices:

$$\begin{bmatrix} 1 & 0 \\ 0 & 0 \end{bmatrix}, \begin{bmatrix} 1 & 0 \\ 0 & 1 \end{bmatrix}, \begin{bmatrix} 0 & 0 \\ 0 & 0 \end{bmatrix}, \begin{bmatrix} 0 & 0 \\ 0 & 1 \end{bmatrix}$$

More generally, for $n \times n$ matrix, there can be 2^n such matrices. This is because there will be n elements, each of which can take two values.

Exercise 4.6

$$2. \ A = \begin{bmatrix} 0 & 4 \\ -1 & 3 \end{bmatrix} \quad B = \begin{bmatrix} 3 & -8 \\ 0 & 1 \end{bmatrix} \quad C = \begin{bmatrix} 1 & 0 & 9 \\ 6 & 1 & 1 \end{bmatrix}$$

$$(a) \ A + B = \begin{bmatrix} 3 & -4 \\ -1 & 4 \end{bmatrix}$$

$$A^T + B^T = \begin{bmatrix} 0 & -1 \\ 4 & 3 \end{bmatrix} + \begin{bmatrix} 3 & 0 \\ -8 & 1 \end{bmatrix} = \begin{bmatrix} 3 & -1 \\ -4 & 4 \end{bmatrix}$$

$$\text{So, } (A + B)^T = A^T + B^T$$

$$(b) \ AC = \begin{bmatrix} 0 & 4 \\ -1 & 3 \end{bmatrix}_{2 \times 2} \begin{bmatrix} 1 & 0 & 9 \\ 6 & 1 & 1 \end{bmatrix}_{2 \times 3} = \begin{bmatrix} 24 & 4 & 4 \\ 17 & 3 & -6 \end{bmatrix}_{2 \times 3} \quad C^T A^T = \begin{bmatrix} 1 & 6 \\ 0 & 1 \\ 9 & 1 \end{bmatrix}_{3 \times 2} \begin{bmatrix} 0 & -1 \\ 4 & 3 \end{bmatrix}_{2 \times 2} =$$

$$\begin{bmatrix} 24 & 17 \\ 4 & 3 \\ 4 & -6 \end{bmatrix}_{3 \times 2} \quad \text{So, } (AC)^T = C^T A^T.$$

6. $A = I - X (X^T X)^{-1} X^T$

- (a.) Say the dimension of X is $m \times n$. Then the dimension of $X_{n \times m}^T X_{m \times n}$ is $n \times n$. So the dimension of $(X^T X)^{-1}$ is also $n \times n$. This implies that the dimension of $X_{m \times n} (X^T X)^{-1}_{n \times n} X_{n \times m}^T$ is $m \times m$. Hence, $X^T X$ and A must be square matrices, but X need not be square.
- (b.) To prove a matrix is idempotent, we need to show $AA = A$.

$$\begin{aligned}
 AA &= (I - X (X^T X)^{-1} X^T)(I - X (X^T X)^{-1} X^T) \\
 &= I - X (X^T X)^{-1} X^T - X (X^T X)^{-1} X^T + \underbrace{X (X^T X)^{-1} X^T X (X^T X)^{-1} X^T}_I \\
 &= I - X (X^T X)^{-1} X^T - X (X^T X)^{-1} X^T + X (X^T X)^{-1} X^T \\
 &= I - X (X^T X)^{-1} X^T = A
 \end{aligned}$$

Exercise 5.1

3. (a) Yes (c) Yes
 (b) Yes (d) No; second row = $-2 \times$ first row.
4. Yes, we get the same answer.

Exercise 5.2

1. (c)

$$\begin{aligned}
 &8 \begin{vmatrix} 0 & 1 \\ 0 & 3 \end{vmatrix} - 1 \begin{vmatrix} 4 & 1 \\ 6 & 3 \end{vmatrix} + 3 \begin{vmatrix} 4 & 0 \\ 6 & 0 \end{vmatrix} \\
 &= 8(0 - 0) - 1(12 - 6) + 3(0 - 0) \\
 &= 0 - 6 + 0 \\
 &= -6
 \end{aligned}$$

(e)

$$\begin{aligned}
& a \begin{vmatrix} c & a \\ a & b \end{vmatrix} - b \begin{vmatrix} b & a \\ c & b \end{vmatrix} + c \begin{vmatrix} b & c \\ c & a \end{vmatrix} \\
&= a(cb - a^2) - b(b^2 - ac) + c(ab - c^2) \\
&= abc - a^3 - b^3 + abc + abc - c^3 \\
&= -a^3 - b^3 - c^3 + 3abc
\end{aligned}$$

(f)

$$\begin{aligned}
& x \begin{vmatrix} y & 2 \\ -1 & 8 \end{vmatrix} - 5 \begin{vmatrix} 3 & 2 \\ 9 & 8 \end{vmatrix} + 0 \begin{vmatrix} 3 & y \\ 9 & -1 \end{vmatrix} \\
&= x(8y + 2) - 5(24 - 18) + 0(-3 - 9y) \\
&= 8xy + 2x - 30 + 0 \\
&= 8xy + 2x - 30
\end{aligned}$$

2. $|C_{13}| : 1 + 3 = 4$ is even so +
 $|C_{23}| : 2 + 3 = 5$ is odd so -
 $|C_{33}| : 3 + 3 = 6$ is even so +
 $|C_{41}| : 4 + 1 = 5$ is odd so -
 $|C_{34}| : 3 + 4 = 7$ is odd so -

3. Minor of a :

$$|M_{11}| = \begin{vmatrix} e & f \\ h & i \end{vmatrix} = ei - fh$$

Cofactor of a :

$$|C_{11}| = (-1)^{1+1} |M_{11}| = |M_{11}|$$

Minor of b :

$$|M_{12}| = \begin{vmatrix} d & f \\ g & i \end{vmatrix} = di - fg$$

Cofactor of b :

$$|C_{12}| = (-1)^{1+2} |M_{12}| = -|M_{12}|$$

Minor of f :

$$|M_{23}| = \begin{vmatrix} a & b \\ g & h \end{vmatrix} = ah - bg$$

Cofactor of f :

$$|C_{23}| = (-1)^5 |M_{23}| = -|M_{23}|$$

6. Minors of third row:

$$|M_{31}| = \begin{vmatrix} 11 & 4 \\ 2 & 7 \end{vmatrix} = 69, \quad |M_{32}| = \begin{vmatrix} 9 & 4 \\ 3 & 7 \end{vmatrix} = 51, \quad |M_{33}| = \begin{vmatrix} 9 & 11 \\ 3 & 2 \end{vmatrix} = -15$$

Cofactors: $|C_{31}| = |M_{31}|$, $|C_{32}| = -|M_{32}|$, $|C_{33}| = |M_{33}|$.

Exercise 5.3

1.

$$\begin{aligned} \begin{vmatrix} 4 & 0 & -1 \\ 2 & 1 & -7 \\ 3 & 3 & 9 \end{vmatrix} &= 4 \begin{vmatrix} 1 & -7 \\ 3 & 9 \end{vmatrix} - 0 \begin{vmatrix} 2 & -7 \\ 3 & 9 \end{vmatrix} - 1 \begin{vmatrix} 2 & 1 \\ 3 & 3 \end{vmatrix} \\ &= 4(9 + 21) - 0(18 + 21) - 1(6 - 3) \\ &= 120 - 0 - 3 = 117 \end{aligned}$$

Interchanging rows and columns:

$$\begin{aligned} \begin{vmatrix} 4 & 2 & 3 \\ 0 & 1 & 3 \\ -1 & -7 & 9 \end{vmatrix} &= 4 \begin{vmatrix} 1 & 3 \\ -7 & 9 \end{vmatrix} - 2 \begin{vmatrix} 0 & 3 \\ -1 & 9 \end{vmatrix} + 3 \begin{vmatrix} 0 & 1 \\ -1 & -7 \end{vmatrix} \\ &= 4(9 + 21) - 2(0 + 3) + 3(0 + 1) \\ &= 120 - 6 + 3 = 117 \end{aligned}$$

Interchange row 1 and 2:

$$\begin{aligned} \begin{vmatrix} 2 & 1 & -7 \\ 4 & 0 & -1 \\ 3 & 3 & 9 \end{vmatrix} &= 2 \begin{vmatrix} 0 & -1 \\ 3 & 9 \end{vmatrix} - 1 \begin{vmatrix} 4 & -1 \\ 3 & 9 \end{vmatrix} - 7 \begin{vmatrix} 4 & 0 \\ 3 & 3 \end{vmatrix} \\ &= 2(0 + 3) - 1(36 + 3) - 7(12 - 0) \\ &= 6 - 39 - 84 = -117 \end{aligned}$$

You can verify the other two properties in the same way.

4. Multiplying every element of an n th-order determinant by k multiplies each of its n rows by k . Multiplying a single row by k multiplies the determinant by k , so doing it to all n rows gives $k^n|A|$.

5. (a)

$$\begin{aligned} & 4 \begin{vmatrix} 1 & -3 \\ 1 & 0 \end{vmatrix} + 1 \begin{vmatrix} 19 & 1 \\ 7 & 1 \end{vmatrix} \\ &= 4(0 + 3) + 1(19 - 7) \\ &= 12 + 12 = 24 \end{aligned}$$

Non-singular, rank = 3.

(b)

$$\begin{aligned} & 5 \begin{vmatrix} -2 & 1 \\ 0 & 3 \end{vmatrix} + 6 \begin{vmatrix} 4 & 1 \\ 7 & 3 \end{vmatrix} \\ &= 5(-6 - 0) + 6(12 - 7) \\ &= -30 + 30 = 0 \end{aligned}$$

Singular. Rank < 3, will need to put in echelon form to find exact rank.

(c)

$$\begin{aligned} & 7 \begin{vmatrix} 1 & 4 \\ -3 & -4 \end{vmatrix} + 1 \begin{vmatrix} 1 & 4 \\ 13 & -4 \end{vmatrix} \\ &= 7(-4 + 12) + 1(-4 - 52) \\ &= 56 - 56 = 0 \end{aligned}$$

Singular. Rank < 3, will need to put in echelon form to find exact rank.

(d)

$$\begin{aligned} & -3 \begin{vmatrix} 9 & 5 \\ 8 & 6 \end{vmatrix} - 1 \begin{vmatrix} -4 & 9 \\ 10 & 8 \end{vmatrix} \\ &= -3(54 - 40) - 1(-32 - 90) \\ &= -42 + 122 = 80 \end{aligned}$$

Non-singular, rank = 3.

8. (a) False. While we can find the transpose of any matrix, the determinant is only defined for square matrices.

- (b) False. Multiplying each element of an $n \times n$ by 2 will increase the determinant by 2^n times.
- (c) A cannot be 0, but the determinant of A can. In any case, if $|A|=0$, A is singular.

Exercise 5.4

2. Note that

$$A^{-1} = \frac{1}{|A|} \text{adj } A$$

Here, $\text{adj } A = C'$ where $C = [C_{ij}]$.

(a)

$$|A| = 5 - 0 = 5$$

$$\text{adj } A = \begin{bmatrix} |C_{11}| & |C_{21}| \\ |C_{12}| & |C_{22}| \end{bmatrix} = \begin{bmatrix} 1 & -2 \\ 0 & 5 \end{bmatrix}$$

$$A^{-1} = \frac{1}{|A|} \text{adj } A = \frac{1}{5} \begin{bmatrix} 1 & -2 \\ 0 & 5 \end{bmatrix} = \begin{bmatrix} 1/5 & -2/5 \\ 0 & 1 \end{bmatrix}$$

(b)

$$|B| = -2 - 0 = -2$$

$$B^{-1} = \frac{-1}{2} \begin{bmatrix} 2 & 0 \\ -9 & -1 \end{bmatrix}$$

(c)

$$|C| = -3 - 21 = -24$$

$$C^{-1} = \frac{-1}{24} \begin{bmatrix} -1 & -7 \\ -3 & 3 \end{bmatrix}$$

(d)

$$|D| = 21 - 0 = 21$$

$$D^{-1} = \frac{1}{21} \begin{bmatrix} 3 & -6 \\ 0 & 7 \end{bmatrix}$$

3. (a) Step 1: Exchange the two diagonal elements.
Step 2: Multiply both the off-diagonal elements by -1.
- (b) Step 3: Multiply the resulting matrix from steps 1 and 2 by $1/|A|$.

4. (a)

$$\begin{aligned}
 |E| &= 1 \begin{vmatrix} 7 & 3 \\ 2 & 0 \end{vmatrix} + 1 \begin{vmatrix} 4 & -2 \\ 7 & 3 \end{vmatrix} \\
 &= 1(0 - 6) + 1(12 + 14) \\
 &= -6 + 26 = 20
 \end{aligned}$$

$$\begin{aligned}
 \text{adj } E &= \begin{bmatrix} |C_{11}| & |C_{21}| & |C_{31}| \\ |C_{12}| & |C_{22}| & |C_{32}| \\ |C_{13}| & |C_{23}| & |C_{33}| \end{bmatrix} \\
 &= \begin{bmatrix} 3 & 2 & -3 \\ -7 & 2 & 7 \\ -6 & -4 & 26 \end{bmatrix} \\
 E^{-1} &= \frac{1}{|E|} \text{adj } E
 \end{aligned}$$

(b)

$$F^{-1} = \frac{-1}{10} \begin{bmatrix} 0 & 2 & -3 \\ 10 & -6 & -1 \\ 0 & -4 & 1 \end{bmatrix}$$

(c)

$$\begin{aligned}
 |C_{11}| &= \begin{vmatrix} 0 & 1 \\ 1 & 0 \end{vmatrix} = -1 & |C_{12}| &= -\begin{vmatrix} 0 & 1 \\ 0 & 0 \end{vmatrix} = 0 & |C_{13}| &= \begin{vmatrix} 0 & 0 \\ 0 & 1 \end{vmatrix} = 0 \\
 |C_{21}| &= -\begin{vmatrix} 0 & 0 \\ 1 & 0 \end{vmatrix} = 0 & |C_{22}| &= \begin{vmatrix} 0 & 1 \\ 0 & 0 \end{vmatrix} = 0 & |C_{23}| &= -\begin{vmatrix} 1 & 0 \\ 0 & 1 \end{vmatrix} = -1 \\
 |C_{31}| &= \begin{vmatrix} 0 & 0 \\ 0 & 1 \end{vmatrix} = 0 & |C_{32}| &= -\begin{vmatrix} 1 & 0 \\ 0 & 1 \end{vmatrix} = -1 & |C_{33}| &= \begin{vmatrix} 1 & 0 \\ 0 & 0 \end{vmatrix} = 0
 \end{aligned}$$

$$G^{-1} = -1 \begin{bmatrix} -1 & 0 & 0 \\ 0 & 0 & -1 \\ 0 & -1 & 0 \end{bmatrix} = \begin{bmatrix} 1 & 0 & 0 \\ 0 & 0 & 1 \\ 0 & 1 & 0 \end{bmatrix}$$

(d)

$$H^{-1} = \begin{bmatrix} 1 & 0 & 0 \\ 0 & 1 & 0 \\ 0 & 0 & 1 \end{bmatrix}$$

7. If a matrix is its own inverse, we would need that $A^2 = I$. This is true for the identity matrix. There are other possibilities such as matrix G in exercise 4.

Exercise 5.4

6. (a)

$$A = \begin{bmatrix} 4 & 3 \\ 2 & 5 \end{bmatrix} \quad v = \begin{bmatrix} x \\ y \end{bmatrix} \quad b = \begin{bmatrix} 28 \\ 42 \end{bmatrix}$$

$$\text{Since } Av = b \Rightarrow A^{-1}Av = A^{-1}b \Rightarrow v = A^{-1}b$$

$$|A| = 20 - 6 = 14$$

$$A^{-1} = \frac{1}{14} \begin{bmatrix} 5 & -3 \\ -2 & 4 \end{bmatrix}$$

$$\begin{aligned} v = A^{-1}b &= \frac{1}{14} \begin{bmatrix} 5 & -3 \\ -2 & 4 \end{bmatrix} \begin{bmatrix} 28 \\ 42 \end{bmatrix} \\ &= \frac{1}{14} \begin{bmatrix} 140 - 126 \\ -56 + 168 \end{bmatrix} = \frac{1}{14} \begin{bmatrix} 14 \\ 112 \end{bmatrix} = \begin{bmatrix} 1 \\ 8 \end{bmatrix} \end{aligned}$$

So $x = 1$ and $y = 8$.

- (b)

$$A = \begin{bmatrix} 4 & 1 & -5 \\ -2 & 3 & 1 \\ 3 & -1 & 4 \end{bmatrix} \quad x = \begin{bmatrix} x_1 \\ x_2 \\ x_3 \end{bmatrix} \quad b = \begin{bmatrix} 8 \\ 12 \\ 5 \end{bmatrix}$$

$$\begin{aligned} |A| &= 4 \begin{vmatrix} 3 & 1 \\ -1 & 4 \end{vmatrix} - 1 \begin{vmatrix} -2 & 1 \\ 3 & 4 \end{vmatrix} - 5 \begin{vmatrix} -2 & 3 \\ 3 & -1 \end{vmatrix} \\ &= 4(12 + 1) - 1(-8 - 3) - 5(2 - 9) \\ &= 52 + 11 + 35 = 98 \end{aligned}$$

$$\text{adj } A = \begin{bmatrix} 13 & 1 & 16 \\ 11 & 31 & 6 \\ -7 & 7 & 14 \end{bmatrix}$$

$$x = A^{-1}b = \frac{1}{98} \begin{bmatrix} 13 & 1 & 16 \\ 11 & 31 & 6 \\ -7 & 7 & 14 \end{bmatrix}_{3 \times 3} \begin{bmatrix} 8 \\ 12 \\ 5 \end{bmatrix}_{3 \times 1} = \frac{1}{98} \begin{bmatrix} 196 \\ 490 \\ 98 \end{bmatrix}_{3 \times 1} = \begin{bmatrix} 2 \\ 5 \\ 1 \end{bmatrix}$$

Exercise 5.5

$$1 \text{ \& } 2. \text{ (a) } A = \begin{bmatrix} 3 & -2 \\ 2 & 1 \end{bmatrix} \rightarrow |A| = 7 \quad b = \begin{bmatrix} 6 \\ 11 \end{bmatrix}$$

By Cramer's Rule:

$$A_1 = \begin{bmatrix} 6 & -2 \\ 11 & 1 \end{bmatrix} \rightarrow |A_1| = 28$$

$$A_2 = \begin{bmatrix} 3 & 6 \\ 2 & 11 \end{bmatrix} \rightarrow |A_2| = 21$$

$$x_1^* = \frac{|A_1|}{|A|} = \frac{28}{7} = 4$$

$$x_2^* = \frac{|A_2|}{|A|} = \frac{21}{7} = 3$$

Taking the inverse:

$$A^{-1} = \frac{1}{|A|} \text{adj} = \frac{1}{7} \begin{bmatrix} 1 & 2 \\ -2 & 3 \end{bmatrix}$$

$$\begin{bmatrix} x_1^* \\ x_2^* \end{bmatrix} = A^{-1}b = \frac{1}{7} \begin{bmatrix} 1 & 2 \\ -2 & 3 \end{bmatrix} \begin{bmatrix} 6 \\ 11 \end{bmatrix} = \frac{1}{7} \begin{bmatrix} 28 \\ 21 \end{bmatrix} = \begin{bmatrix} 4 \\ 3 \end{bmatrix}$$

(b)

$$A = \begin{bmatrix} -1 & 3 \\ 4 & -1 \end{bmatrix} \rightarrow |A| = -11 \quad b = \begin{bmatrix} -3 \\ 12 \end{bmatrix}$$

By Cramer's rule:

$$|A_1| = \begin{vmatrix} -3 & 3 \\ 12 & -1 \end{vmatrix} = -33 \rightarrow x_1^* = \frac{-33}{-11} = 3$$

$$|A_2| = \begin{vmatrix} -1 & -3 \\ 4 & 12 \end{vmatrix} = 0 \rightarrow x_2^* = \frac{0}{-11} = 0$$

By taking the inverse:

$$\begin{bmatrix} x_1^* \\ x_2^* \end{bmatrix} = \frac{-1}{11} \begin{bmatrix} -1 & -3 \\ -4 & -1 \end{bmatrix} \begin{bmatrix} -3 \\ 12 \end{bmatrix} = \frac{-1}{11} \begin{bmatrix} -33 \\ 0 \end{bmatrix} = \begin{bmatrix} 3 \\ 0 \end{bmatrix}$$

(c)

$$\begin{aligned} |A| &= \begin{vmatrix} 8 & -7 \\ 1 & 1 \end{vmatrix} = 15 \\ |A_1| &= \begin{vmatrix} 9 & -7 \\ 3 & 1 \end{vmatrix} = 30 \rightarrow x_1^* = \frac{30}{15} = 2 \\ |A_2| &= \begin{vmatrix} 8 & 9 \\ 1 & 3 \end{vmatrix} = 15 \rightarrow x_2^* = \frac{15}{15} = 1 \end{aligned}$$

Alternatively,

$$\begin{bmatrix} x_1^* \\ x_2^* \end{bmatrix} = \frac{1}{15} \begin{bmatrix} 1 & 7 \\ -1 & 8 \end{bmatrix} \begin{bmatrix} 9 \\ 3 \end{bmatrix} = \begin{bmatrix} 2 \\ 1 \end{bmatrix}$$

(d)

$$\begin{aligned} |A| &= \begin{vmatrix} 5 & 9 \\ 7 & -3 \end{vmatrix} = -78 \\ |A_1| &= \begin{vmatrix} 14 & 9 \\ 4 & -3 \end{vmatrix} = -78 \rightarrow x_1^* = 1 \\ |A_2| &= \begin{vmatrix} 5 & 14 \\ 7 & 4 \end{vmatrix} = -78 \rightarrow x_2^* = 1 \end{aligned}$$

Alternatively,

$$\begin{bmatrix} x_1^* \\ x_2^* \end{bmatrix} = \frac{-1}{78} \begin{bmatrix} -3 & -9 \\ -7 & 5 \end{bmatrix} \begin{bmatrix} 14 \\ 4 \end{bmatrix} = \begin{bmatrix} 1 \\ 1 \end{bmatrix}$$

3. (a)

$$\begin{aligned} |A| &= \begin{vmatrix} 8 & -1 & 0 \\ 0 & 2 & 5 \\ 2 & 0 & 3 \end{vmatrix} \\ &= 8 \begin{vmatrix} 2 & 5 \\ 0 & 3 \end{vmatrix} - 1 \begin{vmatrix} 0 & 5 \\ 2 & 3 \end{vmatrix} = 48 - 10 = 38 \end{aligned}$$

$$\begin{aligned}
 |A_1| &= \begin{vmatrix} 16 & -1 & 0 \\ 5 & 2 & 5 \\ 7 & 0 & 3 \end{vmatrix} \\
 &= 16 \begin{vmatrix} 2 & 5 \\ 0 & 3 \end{vmatrix} - (-1) \begin{vmatrix} 5 & 5 \\ 7 & 3 \end{vmatrix} = 96 - 20 = 76 \\
 x_1^* &= \frac{76}{38} = 2
 \end{aligned}$$

$$\begin{aligned}
 |A_2| &= \begin{vmatrix} 8 & 16 & 0 \\ 0 & 5 & 5 \\ 2 & 7 & 3 \end{vmatrix} \\
 &= 8 \begin{vmatrix} 5 & 5 \\ 7 & 3 \end{vmatrix} - 16 \begin{vmatrix} 0 & 5 \\ 2 & 3 \end{vmatrix} \\
 &= 8 \times -20 - 16 \times -10 = -160 + 160 = 0 \\
 x_2^* &= 0
 \end{aligned}$$

$$\begin{aligned}
 |A_3| &= \begin{vmatrix} 8 & -1 & 16 \\ 0 & 2 & 5 \\ 2 & 0 & 7 \end{vmatrix} \\
 &= 8 \begin{vmatrix} 2 & 5 \\ 0 & 7 \end{vmatrix} + 1 \begin{vmatrix} 0 & 5 \\ 2 & 7 \end{vmatrix} + 16 \begin{vmatrix} 0 & 2 \\ 2 & 0 \end{vmatrix} \\
 &= 112 - 10 - 64 = 38 \rightarrow x_3^* = 1
 \end{aligned}$$

(d)

$$\begin{aligned}
 |A| &= \begin{vmatrix} -1 & 1 & 1 \\ 1 & -1 & 1 \\ 1 & 1 & -1 \end{vmatrix} \\
 &= 1 \begin{vmatrix} -1 & 1 \\ 1 & -1 \end{vmatrix} - 1 \begin{vmatrix} 1 & 1 \\ 1 & -1 \end{vmatrix} + 1 \begin{vmatrix} 1 & -1 \\ 1 & 1 \end{vmatrix} \\
 &= 0 + 2 + 2 = 4
 \end{aligned}$$

$$\begin{aligned}
|A_1| &= \begin{vmatrix} a & 1 & 1 \\ b & -1 & 1 \\ c & 1 & -1 \end{vmatrix} \\
&= a \begin{vmatrix} -1 & 1 \\ 1 & -1 \end{vmatrix} - 1 \begin{vmatrix} b & 1 \\ c & -1 \end{vmatrix} + 1 \begin{vmatrix} b & -1 \\ c & 1 \end{vmatrix} \\
&= 0 - (-b - c) + (b + c) = 2(b + c) & |A_2| &= \begin{vmatrix} -1 & a & 1 \\ 1 & b & 1 \\ 1 & c & -1 \end{vmatrix} \\
&= -1 \begin{vmatrix} b & 1 \\ c & -1 \end{vmatrix} - a \begin{vmatrix} 1 & 1 \\ 1 & -1 \end{vmatrix} + 1 \begin{vmatrix} 1 & b \\ 1 & c \end{vmatrix} \\
&= -1(-b - c) - a(-1 - 1) + 1(c - b) \\
&= b + c - 2a + c - b = 2(a + c) & |A_3| &= \begin{vmatrix} -1 & 1 & a \\ 1 & -1 & b \\ 1 & 1 & c \end{vmatrix} \\
&= -1 \begin{vmatrix} -1 & b \\ 1 & c \end{vmatrix} - 1 \begin{vmatrix} 1 & b \\ 1 & c \end{vmatrix} + a \begin{vmatrix} 1 & -1 \\ 1 & 1 \end{vmatrix} \\
&= -1(-c - b) - 1(c - b) + 2a \\
&= c + b - c + b + 2a = 2(a + b)
\end{aligned}$$

$$x^* = \frac{b+c}{2} \quad y^* = \frac{a+c}{2} \quad z^* = \frac{a+b}{2}$$