# Solutions: Calculus

ECON 441: Introduction to Mathematical Economics — Instructor: Div Bhagia

## Exercise 6.2

2. $y = 5x^2 - 4x$

   (a)
   $$\begin{aligned}\frac{\Delta y}{\Delta x} &= \frac{f(x+\Delta x) - f(x)}{\Delta x} \\ &= \frac{5(x+\Delta x)^2 - 4(x+\Delta x) - \left(5x^2 - 4x\right)}{\Delta x} \\ &= \frac{5\left(x^2 + 2x\Delta x + (\Delta x)^2\right) - 4x - 4\Delta x - 5x^2 + 4x}{\Delta x} \\ &= \frac{10x\Delta x + 5(\Delta x)^2 - 4\Delta x}{\Delta x} = 10x - 4 + 5\Delta x\end{aligned}$$

   (b)
   $$\frac{dy}{dx} = \lim_{\Delta x \to 0} \frac{\Delta y}{\Delta x} = \lim_{\Delta x \to 0} (10x - 4 + 5\Delta x) = 10x - 4$$

   (c) $f'(2) = 10 \times 2 - 4 = 16$ and $f'(3) = 10 \times 3 - 4 = 26$.

3. $y = 5x - 2$

   (a)
   $$\frac{\Delta y}{\Delta x} = \frac{5(x+\Delta x) - 2 - (5x - 2)}{\Delta x} = \frac{5\Delta x}{\Delta x} = 5$$
   It is a constant function.

   (b) No: the difference quotient equals 5 whatever the size of $\Delta x$, so its limit as $\Delta x \to 0$ is also 5, i.e. $dy/dx = 5$.

## Exercise 7.1

3. (a) $f'(x) = 18$, so $f'(1) = 18$ and $f'(2) = 18$.

   (b) $f'(x) = 3cx^2$, so $f'(1) = 3c$ and $f'(2) = 12c$.

   (c) $f'(x) = 10x^{-3}$, so $f'(1) = 10$ and $f'(2) = \frac{10}{8} = \frac{5}{4}$.

   (d) $f'(x) = x^{1/3} = \sqrt[3]{x}$, so $f'(1) = 1$ and $f'(2) = \sqrt[3]{2}$.

   (e) $f'(w) = 2w^{-2/3}$, so $f'(1) = 2$ and $f'(2) = 2 \cdot 2^{-2/3} = 2^{1/3}$.

(f) $f'(w) = \frac{1}{2}w^{-7/6}$, so $f'(1) = \frac{1}{2}$ and $f'(2) = \frac{1}{2} \cdot 2^{-7/6} = 2^{-13/6}$.

Exercise 7.2

3. (d) Let $f(x) = ax - b$ and $g(x) = cx^2$, so $f'(x) = a$ and $g'(x) = 2cx$.

$$\begin{aligned}\frac{d}{dx}[f(x)g(x)] &= f'(x)g(x) + f(x)g'(x)\\ &= acx^2 + (ax - b)2cx\\ &= acx^2 + 2acx^2 - 2bcx\\ &= 3acx^2 - 2bcx\end{aligned}$$

(e) Let $f(x) = (2 - 3x)(1 + x)$ and $g(x) = x + 2$, so $g'(x) = 1$. Using the product rule on $f(x)$ itself:

$$f'(x) = -3(1 + x) + (2 - 3x)(1) = -(6x + 1)$$

Then

$$\begin{aligned}\frac{d}{dx}[f(x)g(x)] &= f'(x)g(x) + f(x)g'(x)\\ &= -(6x + 1)(x + 2) + (2 - 3x)(1 + x)\\ &= -6x^2 - 13x - 2 + 2 - x - 3x^2\\ &= -9x^2 - 14x = -x(9x + 14)\end{aligned}$$

7. (a) With $f(x) = x^2 + 3$, $f'(x) = 2x$, and $g(x) = x$, $g'(x) = 1$:

$$\begin{aligned}\frac{d}{dx}\frac{f(x)}{g(x)} &= \frac{f'(x)g(x) - f(x)g'(x)}{g(x)^2}\\ &= \frac{(2x)x - \left(x^2 + 3\right) \cdot 1}{x^2} = \frac{x^2 - 3}{x^2}\end{aligned}$$

(b)

$$\frac{d}{dx}\frac{x + 9}{x} = \frac{1 \cdot x - 1 \cdot (x + 9)}{x^2} = \frac{-9}{x^2}$$

(c)

$$\frac{d}{dx}\frac{6x}{x + 5} = \frac{6(x + 5) - 1 \cdot 6x}{(x + 5)^2} = \frac{30}{(x + 5)^2}$$

(d)

$$\begin{aligned}\frac{d}{dx}\frac{ax^2+b}{cx+d} &= \frac{2ax(cx+d)-\left(ax^2+b\right)c}{(cx+d)^2}\\ &= \frac{2acx^2+2adx-acx^2-bc}{(cx+d)^2}\\ &= \frac{acx^2+2adx-bc}{(cx+d)^2}\end{aligned}$$

8. $f(x) = ax + b$

(a) $f'(x) = a$

(b) $\frac{d}{dx}\left(ax^2+bx\right) = 2ax + b$

(c)

$$\frac{d}{dx}\frac{1}{ax+b} = \frac{0\cdot(ax+b)-a\cdot 1}{(ax+b)^2} = \frac{-a}{(ax+b)^2}$$

(d)

$$\frac{d}{dx}\frac{ax+b}{x} = \frac{ax-(ax+b)\cdot 1}{x^2} = \frac{-b}{x^2}$$

Exercise 7.3

1.

$$\frac{dy}{du} = 3u^2+2 \qquad \frac{du}{dx} = -2x$$

$$\frac{dy}{dx} = \frac{dy}{du}\cdot\frac{du}{dx} = \left(3u^2+2\right)(-2x) = -2x\left[3\left(5-x^2\right)^2+2\right]$$

2.

$$\begin{aligned}\frac{dw}{dx} &= \frac{dw}{dy}\cdot\frac{dy}{dx}\\ &= 2ay(2bx+c)\\ &= 2a\left(bx^2+cx\right)(2bx+c)\end{aligned}$$

3. (a) Let $z = 3x^2 - 13$, so $y = z^3$:

$$\frac{dy}{dx} = \frac{dy}{dz}\cdot\frac{dz}{dx} = 3z^2(6x) = 18x\left(3x^2-13\right)^2$$

(b) Let $z = 7x^3 - 5$, so $y = z^9$:

$$\frac{dy}{dx} = \frac{dy}{dz} \cdot \frac{dz}{dx} = 9z^8 \left(21x^2\right) = 189x^2 \left(7x^3 - 5\right)^8$$

(c) Let $z = ax + b$, so $y = z^5$:

$$\frac{dy}{dx} = \frac{dy}{dz} \cdot \frac{dz}{dx} = 5z^4(a) = 5a(ax + b)^4$$

4. Chain rule, with $z = 16x + 3$ and $y = z^{-2}$:

$$\frac{dy}{dx} = \frac{dy}{dz} \cdot \frac{dz}{dx} = -2z^{-3}(16) = \frac{-32}{(16x + 3)^3}$$

Quotient rule, where the derivative of the denominator is $2(16x+3) \cdot 16 = 32(16x+3)$:


$$\frac{dy}{dx} = \frac{0 \cdot (16x + 3)^2 - 1 \cdot 32(16x + 3)}{(16x + 3)^4} = \frac{-32}{(16x + 3)^3}$$

The answers are identical.

5. The inverse function is $x = \frac{y-21}{7}$.

$$\frac{dy}{dx} = 7 \qquad \frac{dx}{dy} = \frac{1}{7} = \frac{1}{dy/dx}$$

so the inverse-function rule holds. To graph the inverse with its own independent variable on the horizontal axis, write it as $y = (x - 21)/7$. The two lines are mirror images of each other across the 45-degree line $y = x$:




6. (a)

$$\frac{dy}{dx} = -6x^5 < 0 \text{ for } x > 0$$

So the function is strictly decreasing, hence strictly monotonic, and

$$\frac{dx}{dy} = \frac{1}{dy/dx} = \frac{-1}{6x^5}$$

(b)

$$\frac{dy}{dx} = 20x^4 + 3x^2 + 3 > 0 \text{ for all } x$$

So the function is strictly increasing, hence strictly monotonic, and

$$\frac{dx}{dy} = \frac{1}{20x^4 + 3x^2 + 3}$$

## Exercise 10.5

1. (a) $\frac{dy}{dt} = 2e^{2t+4}$

(b) $\frac{dy}{dt} = -9e^{1-9t}$

(c) $\frac{dy}{dt} = 2te^{t^2+1}$

(d) $\frac{dy}{dt} = -10te^{2-t^2}$

(e) $\dfrac{dy}{dx} = (2ax + b)e^{ax^2+bx+c}$

(f) $\dfrac{dy}{dx} = x\dfrac{d}{dx}e^x + e^x\dfrac{d}{dx}x = xe^x + e^x = (x+1)e^x$

(g) $\dfrac{dy}{dx} = x^2\left(2e^{2x}\right) + 2xe^{2x} = 2x(x+1)e^{2x}$

(h) $\dfrac{dy}{dx} = a\left(xbe^{bx+c} + e^{bx+c}\right) = a(bx+1)e^{bx+c}$

3. (a) $\dfrac{dy}{dt} = \dfrac{35t^4}{7t^5} = \dfrac{5}{t}$

(b) $\dfrac{dy}{dt} = \dfrac{10at^9}{at^{10}} = \dfrac{10}{t}$

(c) $\dfrac{dy}{dt} = \dfrac{1}{t+19}$

(d) $\dfrac{dy}{dt} = 5 \cdot \dfrac{2(t+1)}{(t+1)^2} = \dfrac{10}{t+1}$

(e) $\dfrac{dy}{dx} = \dfrac{1}{x} - \dfrac{1}{1+x} = \dfrac{1}{x(1+x)}$

(f) $\dfrac{dy}{dx} = \dfrac{d}{dx}\left[\ln x + 8\ln(1-x)\right] = \dfrac{1}{x} - \dfrac{8}{1-x} = \dfrac{1-9x}{x(1-x)}$

(g) $\dfrac{dy}{dx} = \dfrac{d}{dx}\left[\ln 2x - \ln(1+x)\right] = \dfrac{2}{2x} - \dfrac{1}{1+x} = \dfrac{1}{x(1+x)}$

(h) $\dfrac{dy}{dx} = 5x^4 \cdot \dfrac{2x}{x^2} + 20x^3\ln x^2 = 10x^3\left(1 + 2\ln x^2\right) = 10x^3(1 + 4\ln x)$

7. (a) Since $\ln y = \ln 3x - \ln(x+2) - \ln(x+4)$,

$$\frac{1}{y}\frac{dy}{dx} = \frac{1}{x} - \frac{1}{x+2} - \frac{1}{x+4} = \frac{8-x^2}{x(x+2)(x+4)}$$

Hence,

$$\frac{dy}{dx} = \frac{8-x^2}{x(x+2)(x+4)} \cdot \frac{3x}{(x+2)(x+4)} = \frac{3\left(8-x^2\right)}{(x+2)^2(x+4)^2}$$

(b) Since $\ln y = \ln\left(x^2+3\right) + x^2 + 1$,

$$\frac{1}{y}\frac{dy}{dx} = \frac{2x}{x^2+3} + 2x = \frac{2x\left(x^2+4\right)}{x^2+3}$$

Hence,

$$\frac{dy}{dx} = \frac{2x\left(x^2+4\right)}{x^2+3} \cdot \left(x^2+3\right)e^{x^2+1} = 2x\left(x^2+4\right)e^{x^2+1}$$

Exercise 7.4

1. (a)

$$\frac{\partial y}{\partial x_1} = 6x_1^2 - 22x_1x_2 \qquad \frac{\partial y}{\partial x_2} = -11x_1^2 + 6x_2$$

(d)

$$\frac{\partial y}{\partial x_1} = \frac{5\left(x_2 - 2\right) - \left(5x_1 + 3\right) \cdot 0}{\left(x_2 - 2\right)^2} = \frac{5}{x_2 - 2}$$
$$\frac{\partial y}{\partial x_2} = \frac{0 \cdot \left(x_2 - 2\right) - 1 \cdot \left(5x_1 + 3\right)}{\left(x_2 - 2\right)^2} = \frac{-\left(5x_1 + 3\right)}{\left(x_2 - 2\right)^2}$$

2. (a)

$$f_x = 2x + 5y \qquad f_y = 5x - 3y^2$$

(b)

$$f_x = (2x)(x - 2) + \left(x^2 - 3y\right) \cdot 1 = 3x^2 - 4x - 3y$$
$$f_y = -3(x - 2) + \left(x^2 - 3y\right) \cdot 0 = -3x + 6$$

3. For (a), $f_x(1, 2) = 2(1) + 5(2) = 12$.
   For (b), $f_x(1, 2) = 3(1)^2 - 4(1) - 3(2) = -7$.

5. (a)

$$U_1 = \frac{\partial U}{\partial x_1} = 2\left(x_1 + 2\right)\left(x_2 + 3\right)^3$$
$$U_2 = \frac{\partial U}{\partial x_2} = 3\left(x_1 + 2\right)^2\left(x_2 + 3\right)^2$$

(b)

$$U_1(3, 3) = 2(3 + 2)(3 + 3)^3 = 2 \times 5 \times 216 = 2160$$

7. (a)

$$f_x = 2x \qquad f_y = 3y^2 \qquad f_z = 4z^3 \qquad \nabla f(x, y, z) = \begin{bmatrix} 2x \\ 3y^2 \\ 4z^3 \end{bmatrix}$$

(b)

$$f_x = yz \qquad f_y = xz \qquad f_z = xy \qquad \nabla f(x, y, z) = \begin{bmatrix} yz \\ xz \\ xy \end{bmatrix}$$

Exercise 8.1

1. (a) Since $y = -x^3 - 3x$,

$$dy = \left(-3x^2 - 3\right) dx = -3\left(x^2 + 1\right) dx$$

4. (a) Writing $Q = kP^{-n}$, $\dfrac{dQ}{dP} = -nkP^{-n-1}$, so

$$\varepsilon_d = \frac{dQ}{dP} \cdot \frac{P}{Q} = \frac{-nkP^{-n-1} \cdot P}{kP^{-n}} = -n$$

No: the elasticity is the constant $-n$, whatever the price.

(b) With $n = 1$, $Q = k/P$, so $PQ = k$: the demand curve is a rectangular hyperbola, downward sloping and approaching both axes. The point elasticity is $\varepsilon_d = -1$ everywhere (unit elastic).

6. (a) At $P = 20$ and $Y = 5{,}000$, $Q = 100 - 40 + 100 = 160$, so

$$\varepsilon_P = \frac{\partial Q}{\partial P} \cdot \frac{P}{Q} = -2 \cdot \frac{20}{160} = -0.25$$

(b)

$$\varepsilon_Y = \frac{\partial Q}{\partial Y} \cdot \frac{Y}{Q} = 0.02 \cdot \frac{5000}{160} = 0.625$$

Exercise 8.2

3. (a)

$$\frac{\partial y}{\partial x_1} = \frac{1 \cdot (x_1 + x_2) - x_1 \cdot 1}{(x_1 + x_2)^2} = \frac{x_2}{(x_1 + x_2)^2}$$

$$\frac{\partial y}{\partial x_2} = \frac{-x_1}{(x_1 + x_2)^2}$$

$$dy = \frac{x_2\, dx_1 - x_1\, dx_2}{(x_1 + x_2)^2}$$

4.

$$\varepsilon_{QP} = \frac{\partial Q}{\partial P} \cdot \frac{P}{Q} = \frac{2bP \cdot P}{Q} = \frac{2bP^2}{a + bP^2 + R^{1/2}}$$

$$\varepsilon_{QR} = \frac{\partial Q}{\partial R} \cdot \frac{R}{Q} = \frac{1}{2}R^{-1/2} \cdot \frac{R}{Q} = \frac{R^{1/2}}{2\left(a + bP^2 + R^{1/2}\right)}$$

5. Write $Q = a + bP^2 + R^{1/2}$ and assume $Q > 0$. For $\varepsilon_{QP} = 2bP^2/Q$:

$$\frac{\partial \varepsilon_{QP}}{\partial P} = \frac{4bP\,Q - 2bP^2 \cdot 2bP}{Q^2} = \frac{4bP\left(a + R^{1/2}\right)}{Q^2}$$

$$\frac{\partial \varepsilon_{QP}}{\partial R} = \frac{-bP^2 R^{-1/2}}{Q^2} < 0$$

So $\varepsilon_{QP}$ is strictly decreasing in $R$. In $P$ it is strictly increasing if $a + R^{1/2} > 0$ and strictly decreasing if $a + R^{1/2} < 0$: the sign does not depend on $P$, so it is strictly monotonic in $P$ either way.

For $\varepsilon_{QR} = R^{1/2}/(2Q)$:

$$\frac{\partial \varepsilon_{QR}}{\partial P} = \frac{-bP\,R^{1/2}}{Q^2} < 0$$

$$\frac{\partial \varepsilon_{QR}}{\partial R} = \frac{R^{-1/2}\,Q - 1}{4Q^2} = \frac{R^{-1/2}\left(a + bP^2\right)}{4Q^2}$$

So $\varepsilon_{QR}$ is strictly decreasing in $P$. In $R$ it is strictly increasing if $a + bP^2 > 0$ and strictly decreasing if $a + bP^2 < 0$: again the sign does not depend on $R$, so it is strictly monotonic in $R$.

6.

$$\varepsilon_{XP} = \frac{\partial X}{\partial P} \cdot \frac{P}{X} = \frac{-2P^{-3} \cdot P}{Y_f^{1/2} + P^{-2}} = \frac{-2}{Y_f^{1/2}P^2 + 1} < 0$$

7. (b)

$$\begin{aligned} dU &= U_x\,dx + U_y\,dy \\ &= 14xy^3\,dx + 21x^2y^2\,dy \\ &= 7xy^2\,(2y\,dx + 3x\,dy) \end{aligned}$$

(f)

$$\begin{aligned} dU &= 3(x - 3y)^2\,dx + 3(x - 3y)^2(-3)\,dy \\ &= 3(x - 3y)^2\,(dx - 3\,dy) \end{aligned}$$

Exercise 8.4

2. (a)

$$\begin{aligned}\frac{dz}{dt} &= z_x\frac{dx}{dt} + z_y\frac{dy}{dt}\\ &= (2x-8y)(3) + \left(-8x-3y^2\right)(-1)\\ &= 14x - 24y + 3y^2\\ &= 42t - 24(1-t) + 3(1-t)^2 = 3t^2 + 60t - 21\end{aligned}$$

(b)

$$\begin{aligned}\frac{dz}{dt} &= z_u\frac{du}{dt} + z_v\frac{dv}{dt} + z_t\\ &= 7(4t) + t\cdot 1 + v\\ &= 28t + t + t + 1 = 30t + 1\end{aligned}$$

(c)

$$\frac{dz}{dt} = f_x\frac{dx}{dt} + f_y\frac{dy}{dt} + f_t = bf_x + kf_y + f_t$$

4. (a)

$$\begin{aligned}\frac{\partial W}{\partial u} &= W_x\frac{\partial x}{\partial u} + W_y\frac{\partial y}{\partial u} + W_u\\ &= (2ax+by)\alpha + bx\gamma + c\\ &= (2a\alpha + b\gamma)(\alpha u + \beta v) + \alpha b\gamma u + c\end{aligned}$$

$$\begin{aligned}\frac{\partial W}{\partial v} &= W_x\frac{\partial x}{\partial v} + W_y\frac{\partial y}{\partial v}\\ &= (2ax+by)\beta + bx\cdot 0\\ &= \beta\left[(2a\alpha + b\gamma)u + 2a\beta v\right]\end{aligned}$$

(b)

$$\frac{\partial W}{\partial u} = f_1\frac{\partial x_1}{\partial u} + f_2\frac{\partial x_2}{\partial u} = 10u\,f_1 + f_2$$

$$\frac{\partial W}{\partial v} = f_1\frac{\partial x_1}{\partial v} + f_2\frac{\partial x_2}{\partial v} = 3f_1 - 12v^2 f_2$$

Exercise 8.5

1. (a) With $F_x = -6$ and $F_y = 1$, the implicit-function rule gives

$$\frac{dy}{dx} = -\frac{F_x}{F_y} = -\frac{-6}{1} = 6$$

Equivalently, solving for $y = 6x - 7$ and differentiating gives $dy/dx = 6$.

(b) With $F_x = 12$ and $F_y = 3$,

$$\frac{dy}{dx} = -\frac{F_x}{F_y} = -\frac{12}{3} = -4$$

(c) With $F_x = 2x + 6$ and $F_y = -1$,

$$\frac{dy}{dx} = -\frac{F_x}{F_y} = -\frac{2x+6}{-1} = 2x + 6$$

2. (d)

$$F_x = 18x^2 \qquad F_y = -3$$

$$\frac{dy}{dx} = -\frac{F_x}{F_y} = -\frac{18x^2}{-3} = 6x^2$$

3. (a)

$$F_x = 2xy^3 + yz \qquad F_y = 3x^2y^2 + xz \qquad F_z = 2z + xy$$

$$\frac{\partial y}{\partial x} = -\frac{F_x}{F_y} = -\frac{2xy^3 + yz}{3x^2y^2 + xz}$$

$$\frac{\partial y}{\partial z} = -\frac{F_z}{F_y} = -\frac{2z + xy}{3x^2y^2 + xz}$$