

Linear Algebra

ECON 441: Introduction to Mathematical Economics

Instructor: Div Bhagia

1 Introduction to Matrices

A matrix is a rectangular array of numbers, parameters, or vectors.

Example. $A = \begin{bmatrix} 2 & 3 & 1 \\ -1 & 4 & 6 \end{bmatrix}$

Dimensions of matrix:

- Number of rows (m)
- Number of columns (n)

A matrix with m rows and n columns is referred to as an $m \times n$ matrix.

It's common to denote the dimensions of a matrix as a subscript. So we can write the 2×3 matrix A in the example above as:

$$A = \begin{bmatrix} 2 & 3 & 1 \\ -1 & 4 & 6 \end{bmatrix}_{2 \times 3}$$

More generally a $m \times n$ matrix can be written as:

$$A = \begin{bmatrix} a_{11} & a_{12} & a_{13} & \dots & a_{1n} \\ a_{21} & a_{22} & a_{23} & \dots & a_{2n} \\ \vdots & \vdots & \vdots & \vdots & \vdots \\ a_{m1} & a_{m2} & a_{m3} & \dots & a_{mn} \end{bmatrix}$$

Or more compactly as:

$$A = [a_{ij}] \quad i = 1, 2, \dots, m; j = 1, 2, \dots, n$$

A square matrix is matrix that has equal number of rows and columns. For example,

$$A = \begin{bmatrix} a_{11} & a_{12} & a_{13} \\ a_{21} & a_{22} & a_{23} \\ a_{31} & a_{32} & a_{33} \end{bmatrix}_{3 \times 3}$$

Two matrices are equal if all their elements are identical. So $A = B$ if and only if $a_{ij} = b_{ij}$ for all i, j . Example.

$$\begin{bmatrix} 1 & 8 \\ 4 & -1 \end{bmatrix} \neq \begin{bmatrix} 1 & 8 \\ 4 & 2 \end{bmatrix}$$

2 Matrix Operations

2.1 Addition and Subtraction

To add two matrices, we need to add each element of one matrix to the corresponding element in the other matrix. Similarly, to take the difference between two matrices, we subtract each element of one matrix from the corresponding element in the other matrix. For this to work, we need both matrices to have the same dimension.

For two matrices of the same dimension, A and B , we have

$$A + B = [a_{ij} + b_{ij}] \quad A - B = [a_{ij} - b_{ij}]$$

In summary:

- Addition and subtraction operations are performed element-by-element
- Only matrices with the same dimension can be added or subtracted.

2.2 Scalar Multiplication

To multiply a matrix A by a scalar λ , we need to multiply each element of A by λ . So

$$\lambda A = [\lambda a_{ij}]$$

Example. Given, $A = \begin{bmatrix} 2 & 3 \\ 4 & -6 \end{bmatrix}$ and $B = \begin{bmatrix} 1 & 8 \\ -2 & 3 \end{bmatrix}$, we can find:

$$\text{a) } A + B = \begin{bmatrix} 3 & 11 \\ 2 & -3 \end{bmatrix}$$

$$\text{c) } 2A = \begin{bmatrix} 4 & 6 \\ 8 & -12 \end{bmatrix}$$

$$\text{b) } A - B = \begin{bmatrix} 1 & -5 \\ 6 & -9 \end{bmatrix}$$

$$\text{d) } A - 2B = \begin{bmatrix} 2 & 3 \\ 4 & -6 \end{bmatrix} - \begin{bmatrix} 2 & 16 \\ -4 & 6 \end{bmatrix} = \begin{bmatrix} 0 & -13 \\ 8 & -12 \end{bmatrix}$$

2.3 Matrix Multiplication

We can only multiply two matrices $A_{m \times n}$ and $B_{p \times q}$ to get AB if $n = p$. In other words, we can only multiply two matrices, if the number of columns in the first matrix is equal to the number of rows in the second matrix.

Example. $A = \begin{bmatrix} 2 & 3 & 1 \\ 4 & -6 & -2 \end{bmatrix}_{2 \times 3}$ $B = \begin{bmatrix} 1 & 8 \\ -2 & 3 \end{bmatrix}_{2 \times 2}$

Here, we cannot find AB , but can find BA .

Note: The order of multiplication matters, so even when both products are possible, AB is not necessarily equal to BA . In fact, the two will only be equal under special circumstances.

If we have two matrices $A_{m \times n}$ and $B_{n \times p}$ and we can multiply these to get $C = AB$. Then the dimension of C will be $m \times p$. To find C we take the i th row of A and the j th row of B to get the ij element of C as follows:

$$c_{ij} = a_{i1}b_{1j} + a_{i2}b_{2j} + \dots + a_{in}b_{nj} = \sum_{k=1}^n a_{ik}b_{kj}$$

Example.

$$A = \begin{bmatrix} a_{11} & a_{12} & a_{13} \\ a_{21} & a_{22} & a_{23} \end{bmatrix}_{2 \times 3} \quad B = \begin{bmatrix} b_{11} & b_{12} \\ b_{21} & b_{32} \\ b_{21} & b_{32} \end{bmatrix}_{3 \times 2}$$

$$C = AB = \begin{bmatrix} a_{11}b_{11} + a_{12}b_{21} + a_{13}b_{31} & a_{11}b_{12} + a_{12}b_{22} + a_{13}b_{32} \\ a_{21}b_{11} + a_{22}b_{21} + a_{23}b_{31} & a_{21}b_{12} + a_{22}b_{22} + a_{23}b_{32} \end{bmatrix}_{2 \times 2}$$

3 Vectors

Matrices with only one column are called column vectors.

$$x = \begin{bmatrix} x_1 \\ x_2 \\ \vdots \\ x_n \end{bmatrix}$$

Matrices with only one row are called row vectors.

$$x' = \begin{bmatrix} x_1 & x_2 & \dots & x_n \end{bmatrix}$$

Inner product of two vectors each with n elements is given by:

$$u \cdot v = u_1v_1 + u_2v_2 + \dots + u_nv_n = \sum_{i=1}^n u_i v_i$$

Example.

$$u = \begin{bmatrix} 1 \\ 5 \\ 2 \end{bmatrix} \quad v = \begin{bmatrix} 2 \\ 1 \\ 3 \end{bmatrix}$$

$$u \cdot v = 1 \cdot 2 + 5 \cdot 1 + 2 \cdot 3 = 2 + 5 + 6 = 13$$

The rows and columns of a matrix are row and column vectors, respectively.

3.1 Linear Dependence

A set of vectors is said to be linearly dependent if and only if any one of them can be expressed as a linear combination of the remaining vectors. Example.

$$v_1 = \begin{bmatrix} 1 \\ 2 \end{bmatrix} \quad v_2 = \begin{bmatrix} 2 \\ 4 \end{bmatrix}$$

Here v_1 and v_2 are linearly dependent because $v_2 = 2v_1$.

Formally, a set of m -vectors $v_1, v_2, \dots, v_n$ is linearly dependent if and only if there exists a set of scalar $k_1, k_2, \dots, k_n$ (not all zero) such that:

$$\sum_{i=1}^n k_i v_i = 0 \quad (m \times 1)$$

4 Identity and Null Matrix

An identity matrix is a square matrix that has all the elements in its principal diagonal equal to 1 and all other elements equal to 0.

A 2×2 identity matrix:

$$I_2 = \begin{bmatrix} 1 & 0 \\ 0 & 1 \end{bmatrix}$$

A 3×3 identity matrix:

$$I_3 = \begin{bmatrix} 1 & 0 & 0 \\ 0 & 1 & 0 \\ 0 & 0 & 1 \end{bmatrix}$$

Identity matrix behaves like 1 for matrices:

$$AI = IA = A$$

In particular, pre- or post-multiplying a matrix by the identity matrix returns the original matrix.

A matrix is an idempotent matrix if it remains unchanged when multiplied by itself any number of times. That is A is idempotent if and only if $A = A^k$.

A null matrix is a matrix with all elements 0.

Example.

$$\begin{bmatrix} 0 & 0 & 0 \\ 0 & 0 & 0 \end{bmatrix}$$

- $A + 0 = A$

- $A0 = 0$

5 Transpose and Inverse of a Matrix

Transpose of a matrix A is denoted by A' and is obtained by interchanging rows and columns of A .

Example.

$$A = \begin{bmatrix} 2 & 3 & 1 \\ 4 & -6 & 2 \end{bmatrix} \quad A' = \begin{bmatrix} 2 & 4 \\ 3 & -6 \\ 1 & 2 \end{bmatrix}$$

- A matrix A is said to be symmetric if $A' = A$
- A matrix A is said to be skew-symmetric if $A' = -A$
- A matrix A is said to be orthogonal if $A'A = I$

Properties of transposes:

$$\begin{aligned} (A')' &= A \\ (A + B)' &= A' + B' \\ (AB)' &= B'A' \end{aligned}$$

For a square matrix A , its inverse A^{-1} is defined as:

$$AA^{-1} = A^{-1}A = I$$

Properties of inverses:

$$\begin{aligned} (A^{-1})^{-1} &= A \\ (AB)^{-1} &= B^{-1}A^{-1} \\ (A')^{-1} &= (A^{-1})' \end{aligned}$$

6 Conditions for Nonsingularity

If a matrix's inverse exists, it's called a nonsingular matrix.

Squareness is a necessary condition for an inverse to exist, but not a sufficient condition. The sufficient condition for nonsingularity of a matrix is if all the rows or (equivalently) columns of this matrix are linearly independent.

Example.

$$A = \begin{bmatrix} 1 & 2 \\ 3 & 4 \end{bmatrix} \quad B = \begin{bmatrix} 1 & 2 \\ 2 & 4 \end{bmatrix}$$

Rows and columns of A are linearly independent, so A is nonsingular. The second row of B can be written as two times the first row of B , so B is a singular matrix.

In summary, a square matrix is nonsingular i.e. its inverse exists if and only if all its rows (or equivalently columns) are linearly independent.

7 Determinants

Determinant $|A|$ is a unique scalar associated with a square matrix A . Before we can write down the determinant for an $n \times n$ matrix, we need to define minors and cofactors.

The minor of the element a_{ij} , denoted by $|M_{ij}|$ is obtained by deleting the i th row and j th column of the matrix and taking the determinant of the resulting matrix.

Whereas, cofactor $|C_{ij}|$ is defined as:

$$|C_{ij}| = (-1)^{i+j} |M_{ij}|$$

Example.

$$A = \begin{bmatrix} a_{11} & a_{12} & a_{13} \\ a_{21} & a_{22} & a_{23} \\ a_{31} & a_{32} & a_{33} \end{bmatrix}$$

Minor for the element a_{12} :

$$|M_{12}| = \begin{vmatrix} a_{21} & a_{23} \\ a_{31} & a_{33} \end{vmatrix}$$

Cofactor for the element a_{12} :

$$|C_{12}| = (-1)^{(1+2)} |M_{12}| = -|M_{12}|$$

Determinant for an $n \times n$ matrix is given by:

$$|A| = \sum_{i=1}^n a_{ij} |C_{ij}| = \sum_{j=1}^n a_{ij} |C_{ij}|$$

The first expression corresponds to expanding with respect to the i th row, while the second expression is the expression for the determinant when expanding with respect to the j th column. In practice, we can expand with respect to any row or column.

Using the above formula we can see that the determinant for a 2×2 matrix:

$$A = \begin{bmatrix} a_{11} & a_{12} \\ a_{21} & a_{22} \end{bmatrix}$$

is given by:

$$|A| = a_{11}a_{22} - a_{12}a_{21}$$

Similarly, the determinant of a 3×3 matrix is given by:

$$\begin{aligned} |A| &= \begin{vmatrix} a_{11} & a_{12} & a_{13} \\ a_{21} & a_{22} & a_{23} \\ a_{31} & a_{32} & a_{33} \end{vmatrix} \\ &= a_{11} \begin{vmatrix} a_{22} & a_{23} \\ a_{32} & a_{33} \end{vmatrix} - a_{12} \begin{vmatrix} a_{21} & a_{23} \\ a_{31} & a_{33} \end{vmatrix} + a_{13} \begin{vmatrix} a_{21} & a_{22} \\ a_{31} & a_{32} \end{vmatrix} \end{aligned}$$

Determinant criteria for nonsingularity: The determinant of singular matrices is equal to 0.

Example.

$$\begin{aligned} \begin{vmatrix} 4 & 0 & -1 \\ 2 & 1 & -7 \\ 3 & 3 & 9 \end{vmatrix} &= 4 \begin{vmatrix} 1 & -7 \\ 3 & 9 \end{vmatrix} - 0 \begin{vmatrix} 2 & -7 \\ 3 & 9 \end{vmatrix} - 1 \begin{vmatrix} 2 & 1 \\ 3 & 3 \end{vmatrix} \\ &= 4(9 + 21) - 0(18 + 21) - 1(6 - 3) \\ &= 120 - 0 - 3 = 117 \end{aligned}$$

Properties of determinants:

1. $|A| = |A'|$

2. Interchanging rows or columns will alter the sign but not the value
3. Multiplication of any one row (or one column) by a scalar k will change the value of the determinant k -fold
4. The addition (subtraction) of a multiple of any row (or column) to (from) another row (or column) will leave the determinant unaltered
5. If one row (or column) is a multiple of another row (or column), the value of the determinant will be zero.

8 Matrix Inversion

To find the inverse of a nonsingular matrix A take the transpose of its cofactor matrix $C = [|C_{ij}|]$ to find the adjoint of A and divide it by the determinant of A .

$$A^{-1} = \frac{1}{|A|} \text{adj} A$$

Adjoint of a nonsingular $n \times n$ matrix

$$\text{adj} A = C' = \begin{bmatrix} |C_{11}| & |C_{21}| & \dots & |C_{n1}| \\ |C_{12}| & |C_{22}| & \dots & |C_{n2}| \\ \vdots & \vdots & \dots & \vdots \\ |C_{1n}| & |C_{2n}| & \dots & |C_{nn}| \end{bmatrix}$$

Example.

$$A = \begin{bmatrix} 4 & -2 & 1 \\ 7 & 3 & 0 \\ 2 & 0 & 1 \end{bmatrix}$$

We can write the cofactors of A as follows:

$$\begin{aligned}
 |C_{11}| &= \begin{vmatrix} 3 & 0 \\ 0 & 1 \end{vmatrix} = 3 & |C_{12}| &= -\begin{vmatrix} 7 & 0 \\ 2 & 1 \end{vmatrix} = -7 & |C_{13}| &= \begin{vmatrix} 7 & 3 \\ 2 & 0 \end{vmatrix} = -6 \\
 |C_{21}| &= -\begin{vmatrix} -2 & 1 \\ 0 & 1 \end{vmatrix} = 2 & |C_{22}| &= \begin{vmatrix} 4 & 1 \\ 2 & 1 \end{vmatrix} = 2 & |C_{23}| &= -\begin{vmatrix} 4 & -2 \\ 2 & 0 \end{vmatrix} = -4 \\
 |C_{31}| &= \begin{vmatrix} -2 & 1 \\ 3 & 0 \end{vmatrix} = -3 & |C_{32}| &= -\begin{vmatrix} 4 & 1 \\ 7 & 0 \end{vmatrix} = 7 & |C_{33}| &= \begin{vmatrix} 4 & -2 \\ 7 & 3 \end{vmatrix} = 26
 \end{aligned}$$

$$\text{adj } A = \begin{bmatrix} |C_{11}| & |C_{21}| & |C_{31}| \\ |C_{12}| & |C_{22}| & |C_{32}| \\ |C_{13}| & |C_{23}| & |C_{33}| \end{bmatrix} = \begin{bmatrix} 3 & 2 & -3 \\ -7 & 2 & 7 \\ -6 & -4 & 26 \end{bmatrix}$$

Now to find the determinant of $|A|$, say we expand using the second row:

$$A = a_{21}|C_{21}| + a_{22}|C_{22}| + a_{23}|C_{23}| = 7 \times 2 + 3 \times 2 + 0 \times -4 = 20$$

Note that expanding by any row or column will give us the same answer. For example, expanding by the third column:

$$A = a_{13}|C_{13}| + a_{23}|C_{23}| + a_{33}|C_{33}| = 1 \times -6 + 0 \times -4 + 1 \times 26 = 20$$

9 Solving a system of linear equations

All the linear algebra we have learned so far can be used to solve a system of linear equations.

For example, we have the following two equations:

$$q + 2p = 100$$

$$q - 3p = 20$$

We can write this as:

$$Ax = b$$

where

$$A = \begin{bmatrix} 1 & 2 \\ 1 & -3 \end{bmatrix} \quad x = \begin{bmatrix} q \\ p \end{bmatrix} \quad b = \begin{bmatrix} 100 \\ 20 \end{bmatrix}$$

If we pre-multiply the equation $Ax = b$ by A^{-1} , we get

$$A^{-1}Ax = A^{-1}b \implies x^* = A^{-1}b$$

9.1 Cramer's Rule

Cramer's rule basically gives a slightly more efficient way of solving a system of equations than finding the inverse. This method is still based on matrix inversion. In particular, the k th element of x can be solved by:

$$x_k^* = \frac{|A_k|}{|A|}$$

where A_k is a matrix formed by exchanging k th column of A by b .