

Astrophysics lab notes

Jacopo Tissino, Giorgio Mentasti

October 8, 2019

30 september 2019

Teacher: Roberto Ragazzoni, Gabriele Umbriaco et al.

roberto.ragazzoni@inaf.it

$$\text{Lab Course} = \text{Course}(\lambda) \quad (1)$$

Objective of this course: *how to build an astronomical instrument* which will push the limits of astrophysical knowledge and technology.

0.1 Orders of magnitude

Diameters: in the future 24 m, 37 m.

Resolution: 0.04 arcsec for the HST.

Collected photons: 2700 Hz for an eye at $V = 6$.

For a galaxy at redshift $z = 10$, the Lyman (?) break at 91.4 nm is shifted.

In the background: magnitude 18 (with moon), 21 (no moon) per arcsec square.

10 meters, “seeing limited”: 1 arcsec; 1 meter, “diffraction limited”: 0.1 arcsec.

Collection ratio: 100X, but the size of the background in which the unresolved source is located is also 100X.

We have a great school of *adaptive optics*.

1 October

<http://people.na.infn.it/~barbarin/MaterialeDidattico/0-corso%20Fisica%20astroparticel0-libro-longair.pdf>

Slides are found at: <http://www.thebotta.com/students/students.htm>

Username: stud, Password: stud_pd.

Bisogna fare una presentazione sui dati che ci faranno analizzare e discuterla poi. Analisi dati al pc molto su dati di alte energie e raggi X, gamma in generale.

Parla dell'importanza degli interferometri LIGO e VIRGO; dei neutrini (Kamiokande, IceCube etc.); raggi cosmici, si incazza coi fisici che usano l'eV per misurare tutto lol. Insomma, le alte energie sono tutto nella vita per un ricercatore.

2 October

What are the sources for X-ray astronomy, beyond the Sun? Many binary systems in our galaxy, there is variability on the scales of milliseconds.

Also, galactic nuclei emit X-rays; also the gas trapped in the middle of clusters of galaxies is very massive.

Diffuse radiation also exists. We do not really know where it comes from.

On top of our atmosphere there is an *ionosphere*, which reflects our radio signals: we can communicate even when we are so far that the curvature of the Earth would prohibit it.

In the sixties, some people put Geiger counters on WWII V2 rockets, and someone actually put a directional sensor: this allowed people to discover the fact that the ionosphere was ionized by X-rays from the Sun.

In the 1960s Giacconi (the fast-moving one) and Rossi (the clever one) thought of a way to build a "telescope" for X-ray astronomy: with gold foil shaped into a section of a parabola we can focus the X-rays. This would allow people to see something but the Sun, like the moon or the Crab Nebula.

Giacconi founded his own company to build the thing.

They used an *anti-coincidence* shield: this allows us to distinguish (charged) cosmic rays from X-rays. This allowed Giacconi to find evidence for X-rays from beyond the solar system: PRL, 1962.

We can use lunar occultation in order to specifically figure out which source the X-rays come from. This was used with the Crab nebula.

In 1965 people discover the fact that X-ray sources can be variable.

We can use collimators in order to select only a specific direction.

An interesting quantity to look at is the ratio of luminosities L_X/L_{optical} . For the Sun, this is of the order 10^{-6} . For Sco X-1 this is 10^3 (!)

If something is moving back and forth, we can use red and blueshift periodicity to figure out the periodicity of the system. With Kepler's law we can get a lower bound for the companion mass. This is done in the optical.

Compared to 1962 we have 10^9 better sensitivity, 10^5 better angular resolution, 10^4 better spectral resolution (now $E/(\Delta E) \sim 10^3$). Now we use X-ray CCDs, not Geiger counters.

8 October 2019

We visit <https://heasarc.gsfc.nasa.gov>

This is the archive where we will spend most of our time: one can find the data from all the current missions.

At the link http://adsabs.harvard.edu/abstract_service.html one can find Astrophysics papers, with full text sources.

There is the extragalactic database <https://ned.ipac.caltech.edu/>, and a galactic database at <http://simbad.u-strasbg.fr/simbad/>.

In <https://heasarc.gsfc.nasa.gov> one can go to Tools → Coordinate converter to find coordinates. If you look at an object and then at another it tells you what the angular distance between them is.

In Tools → energy converter you can convert energies.

Now, let us talk about *specific intensity*: it is defined as

$$I_\nu = \frac{dE_\nu}{dA d\Omega d\nu dt \cos \theta} = \frac{h\nu dN_\nu}{dA d\Omega d\nu dt \cos \theta} \quad (2)$$

We also define *flux density*, measured in $\text{erg cm}^{-2} \text{Hz}^{-1} \text{s}^{-1}$:

$$F_\nu = \int_{\text{source}} I_\nu \cos \theta d\Omega \quad (3)$$

and *flux*, measured in $\text{erg cm}^{-2} \text{s}^{-1}$:

$$F = \int_{\nu_1}^{\nu_2} F_\nu \quad (4)$$

9 October 2019