
[redacted]
Latest report 3 Jun 2026
Generated 3 Oct 2026

15 **results fell outside the range printed for them**

Reference ranges differ between laboratories and depend on age, sex and treatment.

Thyroid & Hormones
1 of 6 to watch



RFT / Kidney
1 of 12 to watch

LFT / Liver
1 of 12 to watch

Vitals

Blood pressure You recorded this	138/84 mmHg 29 Sep 2026
Pulse You recorded this	82 bpm 29 Aug 2026
Oxygen saturation You recorded this	100 % 29 Aug 2026
Temperature You recorded this	37 °C 29 Aug 2026
Weight You recorded this	81 kg 29 Aug 2026
Height You recorded this	178 cm 29 Aug 2026
Waist You recorded this	86.4 cm 2 Sep 2026

Medical history

AFI Condition · You added this	Since 2015 Ongoing
GERD Condition · You added this	Since 2014 Ongoing
Migraine Condition · You added this	Since 2010 Ongoing

Medication the patient reports taking

This is what the patient is taking, not a prescription. Nothing here was prescribed or dispensed through the platform.

PANTOJIVAN D TAB · 1 Tab · Once a day · Empty stomach	Taking now
Tab Pantojivan D · 40/30 · Once a day · Morning	Stopped

YOUR RESULTS

Hematology / Blood count

Hemoglobin ✎	15.6	g/dl	13 - 18
RBC Count ✎	**5.54 ↑**	10^6/ul	4.5 - 5.5
WBC Count ✎	6.52	10^3/ul	4.0 - 10.0

Hematology / Blood count (continued)

Platelets Count	309	10^3/ul	150 - 410
HCT	48.3	%	42 - 52
MCV	87.2	fl	80 - 96
MCH	28.1	pg	27 - 32
MCHC	32.3	g/dl	32 - 36
RDW	12.5	%	11 - 15
Neutrophils	54	%	40 - 80
Lymphocytes	36	%	20 - 40
Eosinophils	06	%	1 - 6
Monocytes	04	%	2 - 10
Platelets Appear On Smear	Adequate		no range
Malarial Parasites (Thick & Thin Smear)	Not Detected		no range
Peripheral Smear	Adequate		no range
ESR	12	mm/hr	0 - 15
Reticulocyte Count	1	%	0.5 - 2.5
S. Iron	78	ug/dl	35.0 - 150.0
TIBC	300	ug/dl	261 - 478
UIBC	222	ug/dl	111 - 343
Transferrin Saturation	26.0	%	17 - 37
Ferritin	200	ng/ml	30 - 400
Reticulocyte Index	1.1		no range
Mentzer Index	15.7		no range

Vitamins

Vitamin B12	913 ↑	pg/ml	211 - 911
Folate	3 ↓	ng/ml	> 5.9
Vitamin D3 (25 Hydroxy Cholecalciferol)	26.31 ↓	ng/ml	30 - 100

Glycemic / Pancreas

FBS	124 ↑	mg/dl	70 - 100
PPBS	93	mg/dl	80 - 140
Glycosylated Hemoglobin	6.44 ↑	%	<= 5.6
Estimated Average Glucose	138.13	mg/dl	no range
Insulin Fasting	12	mIU/ml	2.00 - 22.00
C-Peptide	1	ng/ml	0.5 - 2.0
HOMA Index (Insulin Resistance)	3.67		no range
HOMA Beta Cell Function	70.9		no range
Insulin Sensitivity	27.2		no range

Lipid Profile / Heart

Total Cholesterol	217 ↑	mg/dl	< 200
Triglyceride	178 ↑	mg/dl	< 150

Lipid Profile / Heart (continued)

HDL Cholesterol	43	mg/dl	> 40
LDL Cholesterol	151 ↑	mg/dl	< 100
Non HDL Cholesterol	174 ↑	mg/dl	< 130
VLDLCholesterol	35.6	mg/dl	6 - 38
Chol/HDL Ratio	5.05 ↑		3.5 - 5.0
LDL/HDL Ratio	3.51 ↑		2.5 - 3.5
ApoA1 (Apoliprotein A1)	2	g/l	1.04 - 2.02
ApoB (Apoliprotein B)	1	g/l	0.66 - 1.33
Lipoprotein (Lp a)	24	mg/dl	< 30

RFT / Kidney

Blood Urea	15	mg/dl	15 - 45
BUN	21	mg/dl	8 - 26
S. Creatinine	0.91	mg/dl	0.5 - 1.2
eGFR	104	ml/min/1.73 m2	> 60
BUN/Creatinine Ratio	16		12 - 20
S. Uric Acid	7.6 ↑	mg/dl	3.4 - 7.0
S. Sodium (Na)	140	mmol/L	136 - 145
S. Potassium (K)	4.30	mmol/L	3.50 - 5.10
S. Chloride	104	mmol/L	98 - 107
S. Calcium	10.2	mg/dl	8.7 - 10.7
S. Inorganic Phosphorus	3	mg/dl	2.5 - 4.5
Urine Micro Albumin	1	mg/dL	0 - 1.9

LFT / Liver

Total Bilirubin	0.36	mg/dl	0 - 1.2
Direct Bilirubin	0.14	mg/dl	0 - 0.30
Indirect Bilirubin	0.22	mg/dl	0 - 0.90
SGOT	28	u/l	0 - 40
SGPT	27	u/l	5 - 41
S. Alkaline Phosphatase	68	u/l	40 - 129
GGT	21	u/l	7 - 50
S. LDH (Lactate Dehydrogenase)	182	u/l	110.0 - 240.00
Total Protein	7.8	g/dl	6.6 - 8.7
S. Albumin	4.8	g/dl	3.5 - 4.8
Globulin	3.0	g/dl	2.3 - 3.5
A/G	1.6 ↑		1.2 - 1.5

Thyroid & Hormones

TSH	1.580	uIU/ml	0.35 - 5.50
T3	175.0	ng/dL	60.0 - 181.0
T4	12.88 ↑	ug/dl	4.50 - 12.60

Thyroid & Hormones (continued)

Free T4 ✎	1.1	ng/dL	0.9 - 1.7
PTH (Para Thyroid Hormone) ✎	61	pg/ml	15.00 - 68.30
Total PSA ✎	0.697	ng/ml	0 - 2.5

Inflammatory markers

Homocystein ✎	11	umol/l	5.00 - 15.00
CRP ✎	3	mg/l	0 - 5
High Sensitivity CRP ✎	3.53	mg/l	no range
RA Test with Titre ✎	12	IU/ml	0.00 - 14

WHAT YOUR RESULTS MEAN

Hematology / Blood count


These tests count the cells in your blood: red cells that carry oxygen, white cells that fight infection, and platelets that help a cut stop bleeding. The iron tests show whether your body has enough iron to keep making healthy red cells.

The oxygen-carrying strength of your blood

Hemoglobin

15.6 g/dl
NORMAL

Haemoglobin is the red material inside your red blood cells that picks up oxygen in your lungs and carries it everywhere else.



Range 13 - 18 · ✎ typed in

The number of red blood cells

RBC Count

5.54 10⁶/ul ↑
HIGH

How many red blood cells are packed into a measured amount of your blood.



You have more red cells than usual, or your blood is concentrated from lack of water.

WHAT TO DO

1. Read it beside haemoglobin and MCV, not alone
2. Drink water and repeat if all else is normal

Range 4.5 - 5.5 · ✎ typed in


Hematology / Blood count (continued)

Your infection-fighting cell count

WBC Count

6.52 10³/ul
NORMAL

The total number of white blood cells - the cells that fight infection - in your blood.




Range 4.0 - 10.0 · ✎ typed in

The cells that plug leaks and stop bleeding

Platelets Count

309 10³/ul
NORMAL

The number of platelets in your blood.




Range 150 - 410 · ✎ typed in

How much of your blood is red cells

HCT

48.3 %
NORMAL

If you let your blood settle, this is the share of it that is red cells rather than fluid.



Range 42 - 52 · ✎ typed in

The size of your red blood cells

MCV

87.2 fl
NORMAL

The average size of one red blood cell.



Range 80 - 96 · ✎ typed in

Hematology / Blood count (continued)

How much haemoglobin sits in each red cell

28.1 pg
NORMAL

MCH

The average amount of the red oxygen-carrying material packed into one red blood cell.



Range 27 - 32 · ✎ typed in

How tightly packed the haemoglobin is inside each cell

32.3 g/dl
NORMAL

MCHC

How concentrated the red material is inside a red blood cell - not how much, but how densely packed.




Range 32 - 36 · ✎ typed in

How uneven your red cells are in size

12.5 %
NORMAL

RDW

Whether your red blood cells are all the same size or a mixture of big and small.




Range 11 - 15 · ✎ typed in

The share of your bacteria-fighting cells

54 %
NORMAL

Neutrophils

What share of your white blood cells are neutrophils - the cells that rush to bacterial infections first.



Range 40 - 80 · ✎ typed in

Hematology / Blood count (continued)

The share of your virus-fighting cells

Lymphocytes

36 %
NORMAL

What share of your white cells are lymphocytes - the cells that handle viruses and build long-term immunity.



Range 20 - 40 · ✎ typed in

The share of your allergy and worm-fighting cells

Eosinophils

06 %
NORMAL

The share of your white cells that are eosinophils - cells that switch on for allergies and for worm infections.



Range 1 - 6 · ✎ typed in

The share of your clean-up cells

Monocytes

04 %
NORMAL

The share of your white cells that are monocytes - big cells that clear away debris and handle long-running infections.



Range 2 - 10 · ✎ typed in

Platelets Appear On Smear

Adequate
NOT COMPARED

No reference range published · ✎ typed in

Malarial Parasites (Thick & Thin Smear)

Not Detected
NOT COMPARED

No reference range published · ✎ typed in

Peripheral Smear

Adequate
NOT COMPARED

No reference range published · ✎ typed in

Hematology / Blood count (continued)

How fast your red cells settle - a rough measure of inflammation

12 mm/hr
NORMAL

ESR

Blood is left standing in a thin tube for an hour and the laboratory measures how far the red cells have sunk.



Range 0 - 15 · ✎ typed in

Reticulocyte Count

1 %
NORMAL




Range 0.5 - 2.5 · ✎ typed in

The iron circulating in your blood right now

78 ug/dl
NORMAL

S. Iron

How much iron is travelling in your blood at this moment.



Range 35.0 - 150.0 · ✎ typed in

TIBC

300 ug/dl
NORMAL



Range 261 - 478 · ✎ typed in


Hematology / Blood count (continued)

The empty seats on your iron transport

UIBC

222 ug/dl
NORMAL

Of all the seats available on your iron carrier, this is how many are still empty and waiting for iron.




Range 111 - 343 · ✎ typed in

How full your iron transport is

Transferrin Saturation

26.0 %
NORMAL

What percentage of the seats on your iron carrier actually have iron in them.




Range 17 - 37 · ✎ typed in

Your iron savings account

Ferritin

200 ng/ml
NORMAL

Ferritin is the protein your body stores iron in.



Range 30 - 400 · ✎ typed in

Reticulocyte Index

1.1
NOT COMPARED

No reference range published · ✎ typed in

Mentzer Index

15.7
NOT COMPARED

No reference range published · ✎ typed in

See a doctor if

- Haemoglobin is low along with this
- You feel breathless or unusually tired

Vitamins

These tests measure vitamins your body needs in small amounts every day. Vitamin B12 and folate help you make healthy blood and keep your nerves working, and vitamin D helps your body use calcium to build strong bones.

Vitamin B12

913 pg/ml ↑
HIGH

How much vitamin B12 is in your blood.



You have more B12 in your blood than your body needs. If you take B12 tablets or injections, or a multivitamin or health tonic, that alone explains it and it is harmless - the extra leaves in your urine and there is nothing to treat.

WHAT TO DO

1. Check what you are taking.
2. Go sooner than that if you also have weight loss you cannot explain, a swollen or painful belly, yellow eyes or skin, night sweats, or lumps in the neck, armpit or groin.

Range 211 - 911 · ✎ typed in

Folate

3 ng/ml ↓
LOW



Range > 5.9 · ✎ typed in

Vitamins (continued)

Vitamin D

Vitamin D3 (25 Hydroxy Cholecalciferol)

26.31 ng/ml ↓
LOW

How much vitamin D your body has in store.



Your vitamin D store is running low, so your body is struggling to pull calcium out of the food you eat. To keep the calcium in your blood normal it starts taking calcium out of your bones instead.

WHAT TO DO

1. Get sun on bare forearms, face and legs, and give it real time. Fair skin needs roughly 10 to 15 minutes; medium-brown skin roughly 30 to 45 minutes; dark skin longer still. Do not sit in the sun for hours chasing this - if you would need more than about an hour, a supplement is the sensible route, and it is the one your doctor will choose anyway. Sitting behind a window does nothing; glass blocks it.
2. Not in peak summer midday heat. In the hot months use mid-morning or late afternoon instead, keep water with you, and come inside at once if you feel dizzy, get a headache, stop sweating, or feel your heart racing.

Range 30 - 100 · ✎ typed in

What helps

Step out into the sun most days
a walk after lunch in the cooler months, tea on the terrace, standing outside while on a phone call.

In the hot months move that time to mid-morning or late afternoon
and keep water with you. Do not sit out in peak summer midday heat to chase a vitamin level - a supplement does the same job without the heat

Keep forearms and legs uncovered
for that time; sunscreen on that patch blocks the vitamin. Skip the sun route entirely if sun worsens a skin condition of yours or your medicine makes you sun-sensitive - use a supplement instead

Keep a glass of fortified milk
or a bowl of curd in the daily routine

Babies who are breastfed need vitamin D drops
from birth - breast milk carries very little, whatever the mother's level.

Walk
climb stairs, skip.

See a doctor if

- Chest pain, severe breathlessness at rest, or fainting - go to a hospital emergency department now
- Tingling around the mouth or in the fingers, muscle cramps, twitching, or the hands and feet going into spasm - see a doctor the same day; this can mean the calcium in your blood has fallen too low
- A fit, twitching, a hoarse or crowing sound when breathing in, or difficulty breathing - go to a hospital emergency department now
- A baby or small child with a fit, unusual floppiness, or difficulty breathing - go to a hospital emergency department now
- A high B12 when you take no supplement of any kind, especially with weight loss, a swollen belly, yellow eyes or night sweats - see a doctor in the next few days

Glycemic / Pancreas

These tests measure the sugar in your blood. Your body turns food into sugar for energy, and a hormone called insulin moves that sugar into your cells. HbA1c shows your average sugar over the last two to three months, not just on the day of the test.

FBS

124 mg/dl ↑
HIGH



Range 70 - 100 · ✎ typed in

Sugar in your blood two hours after a meal

PPBS

93 mg/dl
NORMAL

The sugar in your blood two hours after you finish eating.



Range 80 - 140 · ✎ typed in

Your average sugar over the last two to three months

Glycosylated Hemoglobin

6.44 % ↑
HIGH

Sugar sticks to the haemoglobin inside your red blood cells and stays stuck for the life of the cell.



Your sugar has been running above normal most of the time for the last two to three months - not just on the day of the test.

WHAT TO DO

1. Show this to a doctor within a week or two.
2. Check a few sugars at home Fasting and two hours after your biggest meal - and take the list with you

Range <= 5.6 · ✎ typed in

Your average sugar written in glucometer units

Estimated Average Glucose

138.13 mg/dl
NOT COMPARED

This is not a separate blood test.

No reference range published · ✎ typed in

Glycemic / Pancreas (continued)

Insulin level on an empty stomach

Insulin Fasting

12 mIU/ml
NORMAL

How much insulin your pancreas is putting out while you are fasting.



Range 2.00 - 22.00 · ✎ typed in

How much insulin your own pancreas is still making

C-Peptide

1 ng/ml
NORMAL

Every time your pancreas makes insulin it releases an equal amount of C-peptide alongside it.



Range 0.5 - 2.0 · ✎ typed in

HOMA Index (Insulin Resistance)

3.67
NOT COMPARED

No reference range published · ✎ typed in

HOMA Beta Cell Function

70.9
NOT COMPARED

No reference range published · ✎ typed in

Insulin Sensitivity

27.2
NOT COMPARED

No reference range published · ✎ typed in

What helps

- **Keep a small home record** — fasting and after the largest meal, twice a week - and take it to every appointment
- **Walk daily** — activity lowers this number more reliably than any diet trick
- **Choose ragi, jowar, bajra and whole wheat** — over white rice and maida
- **Treat gum disease** — and any long-running infection; they keep sugars up
- **Sleep seven hours and manage stress** — both raise the average quietly

Glycemic / Pancreas (continued)

Only if this is you

Do not start a walking programme if you have an ulcer, a wound, or numbness in your feet
walking on it makes it worse, and the treatment is keeping weight off it. Ask a doctor what activity is safe for you first. And if your sugar is very high and you feel unwell or have ketones, do not exercise until that has been treated - exercise makes ketones worse, not better.

If you have kidney disease
or your creatinine or eGFR is abnormal, do not follow the usual diabetes diet advice about extra dal, curd and fruit without asking your doctor or a dietitian first. Weak kidneys need a set amount of protein, no more, and dal, coconut water, orange, papaya and banana are all high in potassium - which matters especially if you take a blood-pressure tablet whose name ends in -pril or -sartan.

See a doctor if

- This number is much higher or much lower than your home readings suggest
- You have had episodes of sweating, shaking, confusion or nightmares at night
- You have anaemia, thalassaemia trait or sickle trait
- You are pregnant - this is not the test used to diagnose diabetes in pregnancy
- It has been over a year since your eyes, kidneys and feet were checked

Lipid Profile / Heart

These tests measure two kinds of fat in your blood: cholesterol, and triglycerides, which rise with sweets, alcohol and overeating. Too much LDL, the "bad" cholesterol, slowly builds up inside blood vessels, while HDL, the "good" cholesterol, helps carry it away. Some tests here also check the heart itself.

Total cholesterol

Total Cholesterol

217 mg/dl ↑
HIGH

The total amount of cholesterol — a soft, waxy fat — floating in your blood right now.



There is more fat circulating in your blood than your arteries can cope with. Nothing hurts today, and you will feel completely well — the damage takes ten to twenty years to show.

WHAT TO DO

1. Look at your LDL number first That is the one that actually causes blockages
2. Cut deep-fried snacks and bakery items (biscuits, khari, puff, cream rolls) to once a week instead of daily

Range < 200 · ✎ typed in

Lipid Profile / Heart (continued)

Blood fat (triglycerides)

Triglyceride

178 mg/dl ↑
HIGH

The fat your body makes out of extra sugar, starch and alcohol and stores away for energy.



Your body has more fat circulating than it can clear. First check one thing: did you eat, drink tea with sugar, or have a cold drink in the 10 to 12 hours before the blood was taken?

WHAT TO DO

1. Confirm the test was truly fasting If not, repeat it after 10 to 12 hours with only water, because a recent meal inflates this number and makes the calculated LDL on the same report unreliable.
2. Stop sugar in tea, cold drinks, packaged juice and sweets This alone can drop triglycerides 20 to 30 percent in a few weeks

Range < 150 · ✎ typed in

Good cholesterol (HDL)

HDL Cholesterol

43 mg/dl
NORMAL

The cholesterol being carried away from your arteries and back to the liver to be disposed of.



Range > 40 · ✎ typed in

Bad cholesterol (LDL)

LDL Cholesterol

151 mg/dl ↑
HIGH

The cholesterol being carried into your artery walls.



More cholesterol is being pushed into your artery walls than your body clears out. Over years it hardens into plaque, which narrows the artery and can crack open and form a clot — that is a heart attack.

WHAT TO DO

1. Cut the fried and bakery items — samosa, bhajiya, ganthiya, biscuits, khari, puff, cream rolls. This single change moves the number most.
2. Cook in mustard, groundnut or rice bran oil Throw out vanaspati and dalda

Range < 100 · ✎ typed in

Lipid Profile / Heart (continued)

All the harmful cholesterol added up

174 mg/dl ↑
HIGH

Non HDL Cholesterol

Your total cholesterol minus the good HDL — in other words, everything in your blood that can clog an artery, counted together.



The total load of artery-clogging cholesterol in your blood is high. If your LDL looked acceptable but this number is high, the extra is coming from triglyceride-rich particles — and that risk is real even though the LDL line looked fine.

WHAT TO DO

1. Work on both halves: cut fried and bakery food for the LDL, and cut sugar and alcohol for the triglycerides
2. Brisk walk 30 to 40 minutes on five days a week.

Range < 130 · ✎ typed in

VLDL cholesterol

35.6 mg/dl
NORMAL

VLDLCholesterol

The cholesterol sitting inside the particles that carry triglycerides around your body.



Range 6 - 38 · ✎ typed in

Cholesterol to good-cholesterol ratio

5.05 ↑
HIGH

Chol/HDL Ratio

How much of your total cholesterol is the harmful kind compared with the protective kind.



The balance is tilted the wrong way. Either there is too much harmful cholesterol, or too little protective HDL, or both.

WHAT TO DO

1. Look at which side is causing it A low HDL and a high LDL need different actions
2. For a low HDL: brisk walking daily, stop smoking, lose belly weight, cut sugar.

Range 3.5 - 5.0 · ✎ typed in

Lipid Profile / Heart (continued)

Bad-to-good cholesterol ratio

LDL/HDL Ratio

3.51 ↑
HIGH

How much artery-clogging LDL you carry for every unit of protective HDL.



You are carrying a lot of the clogging kind for the amount of protective kind you have. It can be high because LDL is high, because HDL is low, or both — and the fix is different in each case, so read those two lines rather than acting on the ratio alone.

WHAT TO DO

1. Find out which number is driving it and act on that one
2. Get HbA1c and thyroid checked

Range 2.5 - 3.5 · ✎ typed in

ApoA1 — the protein in good cholesterol

ApoA1 (Apoliprotein A1)

2 g/l
NORMAL

The main protein of HDL, the particles that pull cholesterol out of your artery walls and carry it to the liver.




Range 1.04 - 2.02 · ✎ typed in

ApoB — how many harmful particles you carry

ApoB (Apoliprotein B)

1 g/l
NORMAL

Every artery-clogging particle in your blood carries exactly one ApoB protein on its surface.



Range 0.66 - 1.33 · ✎ typed in


Lipid Profile / Heart (continued)

Lp(a) — the cholesterol particle you inherit

Lipoprotein (Lp a)

24 mg/dl
NORMAL

A sticky, clot-prone cholesterol particle whose level is set by the genes you were born with.



Range < 30 · ✎ typed in

What helps

Cook in mustard, groundnut or rice bran oil
Finish vanaspati, dalda and palm oil completely — they are the worst fats for arteries.

Sugar is the main lever here
stop sugar in tea and coffee, cold drinks, packaged juice, and daily sweets.

One bowl of oats
or chana/rajma/moong, every day — soluble fibre traps cholesterol in the gut and carries it out.

Brisk walking 30 to 40 minutes
five days a week — it improves both halves of the ratio at once.

No alcohol while the level is high
Alcohol raises triglycerides more than most people expect, and in heavy drinkers stopping can bring the number down substantially within a few weeks. But if you drink daily and heavily, do not stop suddenly on your own — sudden stopping can cause shaking, fits or confusion.

Stop smoking and tobacco
it raises HDL and protects the artery wall.

Eat one bowl of dal
rajma or chana every day, and put oats, jowar, bajra or ragi in place of some of the white rice and maida.

Only if this is you

If you take warfarin or acitrom
(a blood thinner where you have regular INR tests), do not suddenly increase or cut down green leafy vegetables — palak, methi, sarson, cabbage and broccoli are high in vitamin K, and swinging your intake moves your INR. Keep roughly the same amount week to week, and tell the doctor who manages your INR before you change your diet. The newer blood thinners — apixaban, rivaroxaban, dabigatran — are not affected by vitamin K, so if you are on one of those, eat your vegetables normally.

If you take insulin
or a diabetes tablet of the glimepiride, gliclazide or glibenclamide family, tell your doctor before you make big changes to your food or start walking every day. These changes work — which is why your dose may need to come down at the same time, or your sugar can drop too low. Warning signs of a low sugar are sweating, shaking, hunger, confusion or palpitations.

If you have kidney disease
do NOT increase dal, nuts, coconut water, banana, orange, palak, methi or beetroot on the strength of this page — these are high in potassium and the right amount for you has to be set by your kidney doctor or dietitian.

If you use isabgol (psyllium)
for fibre: one spoon in a full glass of water, drunk straight away, then another glass of water, and never right before lying down — taken dry or lying down it can swell and block the food pipe. Skip it if you have trouble swallowing or have been told you have a narrowing in the bowel.

Lipid Profile / Heart (continued)

See a doctor if

- Chest heaviness, pressure or burning when you walk fast or climb stairs that settles when you stop — see a doctor within a few days if this is a long-standing, unchanged pattern. Go to a hospital the SAME DAY if it is new, if it is coming on with less effort than before, if it happens at rest or wakes you, or if it lasts more than a few minutes.
- Severe pain in the upper stomach going straight through to your back, with vomiting — go to a hospital the same day, this can be pancreatitis
- Chest pressure, heaviness or burning on exertion that settles with rest — see a doctor within a few days if this is a long-standing, unchanged pattern. Go to a hospital the SAME DAY if it is new, if it comes on with less effort than before, if it happens at rest or wakes you, or if it lasts more than a few minutes.
- You have been told your triglycerides are very high on a properly fasting sample — that is treated to protect the pancreas, not watched
- Firm lumps on the heel tendon or knuckles, or a grey-white ring in the eye before age 45 — these point to an inherited high cholesterol that needs family testing
- You are pregnant and your triglycerides are very high

RFT / Kidney

Your kidneys clean your blood, filtering out waste and extra water as urine. These tests check how well that filter is working, and keep an eye on salts such as sodium and potassium that the kidneys hold in balance.

Urea - the waste from the protein you eat

Blood Urea

15 mg/dl
NORMAL

When your body uses protein it makes a waste called urea, and your kidneys wash it out into your urine.



Range 15 - 45 · ✎ typed in

Blood urea nitrogen - the same waste, counted differently

BUN

21 mg/dl
NORMAL

This is the same protein waste as blood urea, measured a different way.



Range 8 - 26 · ✎ typed in


RFT / Kidney (continued)

Creatinine - your kidney's report card

S. Creatinine

0.91 mg/dl
NORMAL

Creatinine is a waste product your muscles make every day at a steady rate.



Range 0.5 - 1.2 · ✎ typed in

eGFR


104 ml/min/1.73 m2
NORMAL



Range > 60 · ✎ typed in

BUN/Creatinine Ratio

16
NORMAL



Range 12 - 20 · ✎ typed in

Uric acid - the gout chemical

S. Uric Acid

7.6 mg/dl ↑
HIGH

Waste made when the body breaks down purines, which are in your own cells and in some foods.



More uric acid than the blood can comfortably carry. It may settle in a joint - classically a red, hot, agonising big toe that wakes you at night - or form kidney stones.

WHAT TO DO

1. Drink water steadily through the day unless a doctor has told you to limit fluids For example with heart failure, liver disease, advanced kidney disease or a low blood sodium
2. Of the things you can change yourself, cutting beer and spirits helps most - start there

Range 3.4 - 7.0 · ✎ typed in


RFT / Kidney (continued)

Sodium - the salt balance in your blood

S. Sodium (Na)

140 mmol/L
NORMAL

Sodium in your blood.



Range 136 - 145 · ✎ typed in

Potassium - the mineral that keeps your heartbeat steady

S. Potassium (K)

4.30 mmol/L
NORMAL

Potassium in your blood.




Range 3.50 - 5.10 · ✎ typed in

Chloride - sodium's partner in the salt balance

S. Chloride

104 mmol/L
NORMAL

Chloride travels alongside sodium in your blood.



Range 98 - 107 · ✎ typed in

Calcium - bone, nerve and muscle mineral

S. Calcium

10.2 mg/dl
NORMAL

The total calcium in your blood - the part carried on protein plus the free active part.



Range 8.7 - 10.7 · ✎ typed in

RFT / Kidney (continued)

Phosphate - calcium's partner in bone

S. Inorganic Phosphorus

3 mg/dl
NORMAL


Phosphate in your blood.



Range 2.5 - 4.5 · ✎ typed in

Urine Micro Albumin

1 mg/dL
NORMAL



Range 0 - 1.9 · ✎ typed in

What helps

Water
less alcohol especially beer, fewer sugary drinks, gradual weight loss - and keep the dairy

Cherries and vitamin C
from food help a little in some people

Be realistic
diet and alcohol changes together usually move uric acid only modestly.

See a doctor if

- A joint suddenly becomes hot, red, swollen and severely painful - get seen the same day, with or without fever. Gout and a joint infection look identical, an infection destroys the joint within days, and many people with an infected joint have no fever at all
- A rash, fever or mouth ulcers in the first weeks after starting allopurinol - stop the tablet and go to a hospital the same day, not to an appointment. This reaction is more common in people of South Asian descent than the drug's reputation suggests, and it can turn into a severe skin and organ illness that is life-threatening. Do not wait to see whether the rash settles
- Uric acid is high and creatinine is rising too
- You have had a kidney stone
- Hard lumps appearing near joints or on the ear

LFT / Liver

Your liver cleans your blood, helps digest fat and makes many of the proteins your body needs. These tests look for signs that liver cells are under strain (SGPT, SGOT) and check how well the liver is doing its work (bilirubin, albumin).

Bilirubin — the yellow pigment that causes jaundice

Total Bilirubin

0.36 mg/dl
NORMAL

Bilirubin is the yellow waste left over when old red blood cells are broken up.




Range 0 - 1.2 · ✎ typed in

Direct bilirubin — the part the liver has already processed

Direct Bilirubin

0.14 mg/dl
NORMAL

This is the share of bilirubin your liver has already packaged and is ready to send out in bile.



Range 0 - 0.30 · ✎ typed in

Indirect bilirubin — the part the liver has not processed yet

Indirect Bilirubin

0.22 mg/dl
NORMAL

This is bilirubin that has arrived at the liver but has not yet been packaged.



Range 0 - 0.90 · ✎ typed in

AST — an enzyme from the liver, and also from muscle

SGOT

28 u/l
NORMAL

SGOT is an enzyme found in liver cells but also in muscle, heart and red blood cells.



Range 0 - 40 · ✎ typed in


LFT / Liver (continued)

ALT — the liver's own enzyme

27 u/l
NORMAL

SGPT

SGPT is an enzyme that sits inside your liver cells.



Range 5 - 41 · ✎ typed in

ALP — an enzyme from your bile ducts and your bones

68 u/l
NORMAL

S. Alkaline Phosphatase

ALP comes mainly from two places — the small bile ducts inside the liver, and bone that is actively being built or repaired.



Range 40 - 129 · ✎ typed in

GGT — the enzyme that separates liver from bone, and reacts to alcohol

21 u/l
NORMAL

GGT

GGT is an enzyme concentrated in the liver and the bile ducts.




Range 7 - 50 · ✎ typed in

LDH — a general 'some tissue is damaged' marker

182 u/l
NORMAL

S. LDH (Lactate Dehydrogenase)

LDH is an enzyme present in almost every cell in the body — muscle, liver, kidney, heart and red blood cells.



Range 110.0 - 240.00 · ✎ typed in


LFT / Liver (continued)

Total protein — albumin and globulin added together

Total Protein

7.8 g/dl
NORMAL

This is all the protein in your blood — albumin plus globulin.



Range 6.6 - 8.7 · ✎ typed in

Albumin — the main protein your liver builds

S. Albumin

4.8 g/dl
NORMAL

Albumin is the main protein in your blood and your liver makes all of it.




Range 3.5 - 4.8 · ✎ typed in

Globulin — mostly your antibodies

Globulin

3.0 g/dl
NORMAL

Globulin is everything in your blood protein that is not albumin.



Range 2.3 - 3.5 · ✎ typed in

A/G ratio — albumin compared with your other blood proteins

A/G

1.6 ↑
HIGH

This is your albumin divided by your globulin.



Albumin is unusually dominant, which normally means globulin is low rather than albumin being high.

WHAT TO DO

1. Look at the actual globulin figure A high ratio matters only if the globulin is genuinely low
2. Otherwise no action is needed for a high ratio alone

Range 1.2 - 1.5 · ✎ typed in

LFT / Liver (continued)

See a doctor if

- Vomiting of blood, or black tarry stools — go to hospital now
- Swelling of the belly or legs
- Fever, night sweats or weight loss for more than two weeks
- Bone pain with tiredness

Thyroid & Hormones

The thyroid is a small gland in the front of your neck, and its hormones set how fast your body uses energy — affecting weight, heartbeat, mood and periods. TSH is the signal from a small gland under your brain that tells the thyroid how hard to work: it rises when the thyroid is slow and falls when it is overactive. Other hormone tests are grouped here too.

Thyroid stimulating hormone — the message your brain sends to your thyroid

TSH

1.580 ulU/ml
NORMAL

TSH is the instruction your brain sends to the thyroid gland in your neck, telling it how hard to work.



Range 0.35 - 5.50 · ✎ typed in

Total T3 — all the active thyroid hormone, locked and unlocked together

T3

175.0 ng/dL
NORMAL

The total amount of the active thyroid hormone T3 in your blood, most of which is bound to carrier proteins.



Range 60.0 - 181.0 · ✎ typed in

Thyroid & Hormones (continued)

Total T4 — all the main thyroid hormone in your blood, locked and unlocked together

T4

12.88 ug/dl ↑
HIGH

This counts every bit of T4 in the blood, including the large part that is stuck to carrier proteins and cannot be used.



There is a lot of thyroid hormone in the blood — either a genuinely overactive thyroid, or extra carrier protein holding more hormone that your body cannot actually use. A low TSH on the same report is what makes it real.

WHAT TO DO

1. Read it with TSH — a suppressed TSH usually makes it real, with one important exception: a biotin (hair-and-nail) supplement pushes TSH down and T3/T4 up at the same time and imitates this exact pattern. Stop biotin for at least a week and repeat the whole panel before any treatment is started
2. Ask for free T4 as well — it is not affected by pregnancy or the pill

Range 4.50 - 12.60 · ✎ typed in

Free T4 — the actual thyroid hormone available to your body

Free T4

1.1 ng/dL
NORMAL

T4 is the main hormone the thyroid makes.



Range 0.9 - 1.7 · ✎ typed in

Parathyroid hormone — the hormone that controls the calcium in your blood

PTH (Para Thyroid Hormone)

61 pg/ml
NORMAL

PTH comes from four tiny glands sitting behind the thyroid.



Range 15.00 - 68.30 · ✎ typed in

Thyroid & Hormones (continued)

Total PSA

0.697 ng/ml
NORMAL



Range 0 - 2.5 · ✎ typed in

See a doctor if

- If your thyroid is underactive: becoming very drowsy, confused, unusually cold to touch, or hard to wake — go to hospital immediately, especially in an older person or during any infection. Do not wait for an appointment
- If your thyroid is overactive: fever with a very fast or irregular heartbeat, agitation, confusion, vomiting or diarrhoea — go to hospital immediately, do not wait for an appointment
- A heartbeat that feels irregular, breathlessness when lying flat, or swelling of the ankles — get seen the same day
- TSH is abnormal along with this
- Neck swelling, weight change, racing heart or persistent tiredness

Inflammatory markers


These tests look for inflammation, the swelling and heat your body makes to fight an infection or heal an injury. It also happens when the body's defences attack the body itself by mistake. CRP and ESR go up when there is inflammation anywhere in the body, and the antibody tests help a doctor look for diseases such as rheumatoid arthritis, which makes joints swollen and painful.

Homocysteine

Homocystein

11 umol/l
NORMAL


An amino acid your body makes while processing protein.



Range 5.00 - 15.00 · ✎ typed in

CRP

3 mg/l
NORMAL



Range 0 - 5 · ✎ typed in

Inflammatory markers (continued)

hs-CRP — inflammation measured for heart risk

High Sensitivity CRP

3.53 mg/l
NOT COMPARED

The same CRP protein, but measured with a finer method that can pick up the very small amounts a healthy person carries.

No reference range published · ✎ typed in

RA factor — the older arthritis antibody test

RA Test with Titre

12 IU/ml
NORMAL

Rheumatoid factor is an antibody that attacks other antibodies.



Range 0.00 - 14 · ✎ typed in

Keep it up

No tobacco
Tobacco keeps the whole body mildly inflamed and raises the risk of rheumatoid arthritis.

Stay active
Walking 30 minutes a day lowers inflammation in the body and keeps joints moving.

Watch your waist
Fat around the belly sends out inflammation signals of its own.

Sleep 7 hours
Short sleep raises inflammation.